

Rehabilitation Services

Pelvic Health

Physical Therapy Services

Cozean Pelvic Dysfunctional Screening Scale

- I sometimes have pelvic pain (in genitals, perineum, pubic or bladder area, or pain with urination) that exceeds a 3 on a 1-10 pain scale, with 10 being the worst pain imaginable
- I can remember falling onto my tailbone, lower back, or buttocks (even in childhood)
- I sometimes experience one or more of the following urinary symptoms:
 - Accidental loss of urine
 - Feeling unable to completely empty my bladder
 - Having to void within a few minutes of a previous void
 - Pain or burning with urination
 - Difficulty starting or frequent stopping/starting of urine stream
 - I often or occasionally have to get up to urinate two or more times at night
- I sometimes have a feeling of increased pelvic pressure or the sensation of my pelvic organs slipping down or falling out
- I have a history of pain in my low back, hip, groin, or tailbone or have had sciatica
- I sometimes experience one or more of the following bowel symptoms:
 - Loss of bowel control
 - Feeling unable to completely empty my bowels
 - Straining or pain with a bowel movement
 - Difficulty initiating a bowel movement
- I sometimes experience pain or discomfort with sexual activity or intercourse
- Sexual activity increases one or more of my other symptoms
- Prolonged sitting increases my symptoms

IF YOU CHECKED 3 OR MORE BOXES, PELVIC FLOOR DYSFUNCTION IS LIKELY. YOU MAY BENEFIT FROM AN ASSESSMENT FROM A PELVIC FLOOR PHYSICAL THERAPIST.