

Community Health Needs Assessment

CHI Health Immanuel – Omaha, NE 2025

A Joint Assessment

Adopted April 2025



EXECUTIVE SUMMARY

CHNA Purpose

The purpose of this Community Health Needs Assessment (CHNA) is to identify and prioritize significant health needs in the community served by CHI Health Immanuel. The priorities identified in this report help to guide the hospital's community health improvement programs and community benefit activities, as well as its collaborative efforts with other organizations that share a mission to improve health. This CHNA report meets requirements of the Patient Protection and Affordable Care Act that not-for-profit hospitals conduct a community health needs assessment at least once every three years.

CommonSpirit Health Commitment & Mission

The hospital's commitment to engaging with the community, assessing priority needs, and helping to address them with community partners is in keeping with its mission. As CommonSpirit Health, we make the healing presence of God known in our world by improving the health of the people we serve, especially those who are vulnerable, while we advance social justice for all.


CHNA Collaborators

This is a joint assessment, and CHI Health Immanuel and CHI Health collaborated with multiple community partners on this important work, including the Douglas County Health Department, the Pottawattamie County Public Health, the Sarpy/Cass Health Department, Nebraska Medicine, Methodist Health System, Charles Drew Health Center, Inc., One World Community Health Centers, Inc., All Care Health Center, and The Wellbeing Partners. This assessment was conducted on behalf of our organizations by PRC, a nationally recognized health care consulting firm with extensive experience conducting Community Health Needs Assessments in hundreds of communities across the United States since 1994.

Community Definition

The study area, and shared community definition, for this assessment is defined as the four counties that make up the Omaha Metro Area: Douglas, Sarpy, and Cass Counties in Nebraska, and Pottawattamie County in Iowa. These counties include the following ZIP Codes: 51501, 51503, 51510, 51521, 51525, 51526, 51536, 51542, 51548, 51549, 51553, 51559, 51560, 51570, 51575, 51576, 51577, 68005, 68007,

68016, 68022, 68028, 68037, 68046, 68048, 68058, 68059, 68064, 68069, 68102, 68104, 68105, 68106, 68107, 68108, 68110, 68111, 68112, 68113, 68114, 68122, 68123, 68124, 68127, 68128, 68130, 68131, 68132, 68133, 68134, 68135, 68137, 68138, 68142, 68144, 68147, 68152, 68154, 68157, 68164, 68182, 68198, 68304, 68307, 68347, 68349, 68366, 68403, 68407, 68409, 68413, 68455, and 68463.





Assessment Process & Methods

This assessment incorporates data from multiple sources, including primary research (through the PRC Community Health Survey and PRC Online Key Informant Survey), as well as secondary research (vital statistics and other existing health-related data). It also allows for trending and comparison to benchmark data at the state and national levels.

Primary Data Collection. Primary data represent the most current information provided in this assessment. The PRC Community Health Survey provides an aggregate snapshot of the health experience, behaviors, and needs of residents in the community. The PRC Online Key Informant Survey allows key community leaders and providers in the area an opportunity to give extensive qualitative input about what they see as the most pressing issues in the populations they serve.

Secondary Data Collection. Secondary data provide information from existing data sets (e.g, public health records, census data, etc.) that complement the primary research findings.

Identifying & Prioritizing Significant Health Needs

Significant health needs for the community were identified through a review of the data collected for this assessment. These were determined after consideration of various criteria, including: standing in comparison with benchmark data (particularly national data); identified trends; the preponderance of significant findings within topic areas; the magnitude of the issue in terms of the number of persons affected; and the potential health impact of a given issue. These also take into account those issues of greatest concern to the key informants giving input to this process.

Prioritization of the health needs was determined based on a prioritization exercise conducted among providers and other community leaders (representing a cross-section of community-based agencies and organizations) as part of the Online Key Informant Survey.

This process yielded the following prioritized list of community health needs:

- MENTAL HEALTH ► Key informants identified this as a top concern. The Metro Area experiences
 a relatively high suicide rate. Survey findings revealed needs related to anxiety, stress, depression,
 overall mental health status, lack of social support, and mental health treatment.
- SOCIAL DETERMINANTS OF HEALTH ➤ Key informants identified this as a top concern in the
 community. Survey findings revealed needs related to housing, financial resilience, discrimination,
 and adverse childhood experiences.
- 3. DIABETES ► Key informants identified this as a top concern in the community. Survey findings found a high prevalence of pre-diabetes in the community.
- 4. NUTRITION, PHYSICAL ACTIVITY & WEIGHT ► Key informants identified this as a top concern in the community. Survey findings revealed a relatively high overweight/obesity prevalence, as well as needs related to a lack of physical activity and the built environment.
- 5. SUBSTANCE USE ► Key informants identified this as a top concern in the community. Existing data revealed adverse trends in alcohol-induced deaths and unintentional drug-induced deaths.
- 6. HEART DISEASE & STROKE ► Existing data show cardiovascular disease to be a leading cause of death. Survey findings revealed relatively high prevalence for both heart disease and stroke.
- 7. INFANT HEALTH & FAMILY PLANNING ► Existing data revealed needs relative to prenatal care.
- 8. INJURY & VIOLENCE ➤ Survey findings revealed needs related to fall-related injuries, as well as intimate partner violence.



Other health needs identified in this assessment include:

- DISABLING CONDITIONS
- ORAL HEALTH
- CANCER
- ACCESS TO HEALTH CARE SERVICES
- TOBACCO USE
- RESPIRATORY DISEASE


Resources Potentially Available to Meet Significant Health Needs

Measures and resources (such as programs, organizations, and facilities in the community) potentially available to address the significant health needs were identified by key informants giving input to this process. While not exhaustive, this list — which includes many potential resources — draws on the experiences and wide knowledge base of those directly serving our community.

Report Adoption, Availability & Comments

This CHNA report was adopted by the CHI Health Board of Directors in April 2025. The report is widely available to the public on the hospital's website, and a paper copy is available for inspection upon request at the Administration Office of CHI Health Immanuel. Written comments on this report can be submitted via mail to CHI Health - The McAuley Fogelstrom Center (12809 W Dodge Rd, Omaha, NE 68154 attn. Healthy Communities); electronically at: https://forms.gle/KGRq62swNdQyAehX8 or by calling Ashley Carroll, Market Director, Community and Population Health, at: (402) 343-4548.





Sponsored by

Douglas County Health Department
Pottawattamie County Public Health
Sarpy/Cass Health Department
CHI Health
Nebraska Medicine
Methodist Health System
Charles Drew Health Center, Inc.
One World Community Health Centers, Inc.
All Care Health Center

With Support From

The Wellbeing Partners



TABLE OF CONTENTS

INTRODUCTION	5
PROJECT OVERVIEW	6
Project Goals	6
Approach	7
Methodology	7
SUMMARY OF FINDINGS	14
Significant Health Needs of the Community	14
Summary Tables: Comparisons With Benchmark Data	17
COMMUNITY DESCRIPTION	31
POPULATION CHARACTERISTICS	32
Total Population	32
Urban/Rural Population	34
Age Race & Ethnicity	35 36
Linguistic Isolation	37
SOCIAL DETERMINANTS OF HEALTH	39
Poverty	39
Education	41
Employment	42
Financial Resilience	43
Housing	45
Transportation Food Access	48 49
Discrimination	52
Adverse Childhood Experiences (ACEs)	57
Key Informant Input: Social Determinants of Health	60
HEALTH STATUS	64
OVERALL HEALTH STATUS	65
MENTAL HEALTH	67
Mental Health Status	67
Depression & Anxiety	68
Stress	71
Social Support Suicide	72 73
Mental Health Treatment	75
Key Informant Input: Mental Health	77
DEATH, DISEASE & CHRONIC CONDITIONS	82
LEADING CAUSES OF DEATH	83
Distribution of Deaths by Cause	83
Age-Adjusted Death Rates for Selected Causes	83
CARDIOVASCULAR DISEASE	85
Age-Adjusted Heart Disease & Stroke Deaths	85
Prevalence of Heart Disease & Stroke	88
Key Informant Input: Heart Disease & Stroke	89



CANCER	92
Age-Adjusted Cancer Deaths	92
Cancer Incidence	95
Prevalence of Cancer Cancer Screenings	96 97
Key Informant Input: Cancer	99
RESPIRATORY DISEASE	101
Age-Adjusted Respiratory Disease Deaths	101
Prevalence of Respiratory Disease	104
Key Informant Input: Respiratory Disease	106
INJURY & VIOLENCE	107
Unintentional Injury	107
Intentional Injury (Violence) Key Informant Input: Injury & Violence	111 116
DIABETES	118
Age-Adjusted Diabetes Deaths	118
Prevalence of Diabetes	120
Age-Adjusted Kidney Disease Deaths	121
Key Informant Input: Diabetes	123
DISABLING CONDITIONS	126
Activity Limitations	126
Alzheimer's Disease Key Informant Input: Disabling Conditions	128 130
BIRTHS	132
PRENATAL CARE	133
BIRTH OUTCOMES & RISKS	135
Low-Weight Births	135
Infant Mortality	136
FAMILY PLANNING	138
Births to Adolescent Mothers Key Informant Input: Infant Health & Family Planning	138 139
	139
MODIFIABLE HEALTH RISKS	141
NUTRITION	142
Difficulty Accessing Fresh Produce	142
Sugar-Sweetened Beverages	144
PHYSICAL ACTIVITY	145
Leisure-Time Physical Activity Activity Levels	145 146
Built Environment	147
WEIGHT STATUS	150
Adult Weight Status	150
Key Informant Input: Nutrition, Physical Activity & Weight	153
SUBSTANCE USE	156
Alcohol Use	156
Drug Use	159
Alcohol & Drug Treatment Key Informant Input: Substance Use	162 162
,	102



TOBACCO USE Cigarette Smoking Use of Vaping Products Key Informant Input: Tobacco Use	166 166 169 170
SEXUAL HEALTH HIV Sexually Transmitted Infections (STIs) Key Informant Input: Sexual Health	172 172 173 174
ACCESS TO HEALTH CARE	176
HEALTH INSURANCE COVERAGE Type of Health Care Coverage Lack of Health Insurance Coverage	177 177 177
DIFFICULTIES ACCESSING HEALTH CARE Difficulties Accessing Services Barriers to Health Care Access Key Informant Input: Access to Health Care Services	179 179 180 181
PRIMARY CARE SERVICES Access to Primary Care Specific Source of Ongoing Care Utilization of Primary Care Services	183 183 184 184
EMERGENCY ROOM UTILIZATION	186
ORAL HEALTH Dental Care Key Informant Input: Oral Health	187 187 188
LOCAL RESOURCES HEALTH CARE RESOURCES & FACILITIES Federally Qualified Health Centers (FQHCs) Resources Available to Address Significant Health Needs	190 191 191 192
IMPACT OF ACTIONS TAKEN SINCE THE PRECEDING CHNA	199





INTRODUCTION

PROJECT OVERVIEW

Project Goals

This Community Health Needs Assessment — a follow-up to similar studies conducted in 2011, 2015, 2018, and 2021 — is a systematic, data-driven approach to determining the health status, behaviors, and needs of residents in Omaha metropolitan area (including Douglas, Sarpy, Cass, and Pottawattamie counties). Subsequently, this information may be used to inform decisions and guide efforts to improve community health and wellness.

A Community Health Needs Assessment provides information so that communities may identify issues of greatest concern and decide to commit resources to those areas, thereby making the greatest possible impact on community health status. This Community Health Needs Assessment will serve as a tool toward reaching three basic goals:

- To improve residents' health status, increase their life spans, and elevate their overall quality of life. A healthy community is not only one where its residents suffer little from physical and mental illness, but also one where its residents enjoy a high quality of life.
- To reduce the health disparities among residents. By gathering demographic information along with health status and behavior data, it will be possible to identify population segments that are most atrisk for various diseases and injuries. Intervention plans aimed at targeting these individuals may then be developed to combat some of the socio-economic factors that historically have had a negative impact on residents' health.
- To increase accessibility to preventive services for all community residents. More accessible preventive services will prove beneficial in accomplishing the first goal (improving health status, increasing life spans, and elevating the quality of life), as well as lowering the costs associated with caring for late-stage diseases resulting from a lack of preventive care.

This assessment was led by a coalition of local public health departments, health systems, federally qualified health centers, and community-based organizations.

SPONSORING ORGANIZATIONS Douglas County Health Department; Pottawattamie County Public Health; Sarpy/Cass Health Department; CHI Health (CHI Health Creighton University Medical Center—Bergan Mercy, CHI Health Immanuel, CHI Health Lakeside, CHI Health Mercy Council Bluffs, and CHI Health Midlands); Nebraska Medicine (Bellevue Medical Center and Nebraska Medical Center); and Methodist Health System (Methodist Hospital, Methodist Jennie Edmundson Hospital, and Methodist Women's Hospital)

SUPPORTING ORGANIZATIONS ► Charles Drew Health Center, Inc.; One World Community Health Centers, Inc.; All Care Health Center; and The Wellbeing Partners

This assessment was conducted by Professional Research Consultants, Inc. (PRC), a nationally recognized health care consulting firm with extensive experience conducting Community Health Needs Assessments in hundreds of communities across the United States since 1994.



Approach

The process for this assessment follows an approach as outlined in the Community Health Assessment Toolkit developed by the Association for Community Health Improvement™ (ACHI). In the ACHI model (at right), collaborating organizations worked through the first three steps in this process, and this assessment document and subsequent communication activities will carry the community engagement model through Step 6. Steps 7 through 9 will be undertaken by the partnering hospitals, health departments, and other organizations over the next three years, at which time the process begins again and this assessment will be updated.



Methodology

This assessment incorporates data from multiple sources, including primary research (through the PRC Community Health Survey and PRC Online Key Informant Survey), as well as secondary research (vital statistics and other existing health-related data). It also allows for trending and comparison to benchmark data at the state and national levels.

PRC Community Health Survey


Survey Instrument

The survey instrument used for this study is based largely on the Centers for Disease Control and Prevention (CDC) Behavioral Risk Factor Surveillance System (BRFSS), as well as various other public health surveys and customized questions addressing gaps in indicator data relative to health promotion and disease prevention objectives and other recognized health issues. The final survey instrument was developed by the sponsoring and supporting organizations and PRC and is similar to the previous surveys used in the region, allowing for data trending.

Community Defined for This Assessment

The study area for the survey effort (referred to as the "Metro Area" in this report) includes Douglas, Sarpy, and Cass counties in Nebraska, as well as Pottawattamie County in Iowa. For this study, Douglas County is further divided into five geographical areas (Northeast Omaha, Southeast Omaha, Northwest Omaha, Southwest Omaha, and Western Douglas County). This community definition is illustrated in the following map.





Sample Approach & Design

A precise and carefully executed methodology is critical in asserting the validity of the results gathered in the PRC Community Health Survey. Thus, to ensure the best representation of the population surveyed, a mixed-mode methodology was implemented. This included targeted surveys conducted by PRC via telephone (landline and cell phone) or through online questionnaires, as well as a community outreach component promoted by the study sponsors through social media posting and other communications.

RANDOM-SAMPLE SURVEYS (PRC) ▶ For the targeted administration, PRC administered 2,960 surveys throughout the service area.


COMMUNITY OUTREACH SURVEYS (Sponsoring and Supporting Partners) PRC also created a link to an online version of the survey, and the sponsors and supporting partners promoted this link locally in order to drive additional participation and bolster overall samples. This yielded an additional 691 surveys to the overall sample.

In all, 3,651 surveys were completed through these mechanisms, including 1,997 in Douglas County, 844 in Sarpy County, 227 in Cass County, and 583 in Pottawattamie County. Once the interviews were completed, these were weighted in proportion to the actual population distribution so as to appropriately represent the Metro Area as a whole. All administration of the surveys, data collection, and data analysis was conducted by PRC.

For statistical purposes, for questions asked of all respondents, the maximum rate of error associated with a sample size of 3,651 respondents is $\pm 1.6\%$ at the 95 percent confidence level.



Expected Error Ranges for a Sample of 3,651 Respondents at the 95 Percent Level of Confidence



- The "response rate" (the percentage of a population giving a particular response) determines the error rate associated with that response. A "95 percent level of confidence" indicates that responses would fall within the expected error range on 95 out of 100 trials.
- Examples: If 10% of the sample of 3,651 respondents answered a certain question with a "yes," it can be asserted that between 9.1% and 10.9% (10% ± 0.9%) of the total population would offer this response.
 - If 50% of respondents said "yes," one could be certain with a 95 percent level of confidence that between 48.2% and 51.8% (50% ± 1.6%) of the total population would respond "ves" if asked this guestion.


Sample Characteristics

To accurately represent the population studied, PRC strives to minimize bias through application of a proven telephone methodology and random-selection techniques. While this random sampling of the population produces a highly representative sample, it is a common and preferred practice to "weight" the raw data to improve this representativeness even further. This is accomplished by adjusting the results of a random sample to match the geographic distribution and demographic characteristics of the population surveyed (poststratification), so as to eliminate any naturally occurring bias. Specifically, once the raw data are gathered, respondents are examined by key demographic characteristics (namely sex, age, race, ethnicity, and poverty status), and a statistical application package applies weighting variables that produce a sample which more closely matches the population for these characteristics. Thus, while the integrity of each individual's responses is maintained, one respondent's responses might contribute to the whole the same weight as, for example, 1.1 respondents. Another respondent, whose demographic characteristics might have been slightly oversampled, might contribute the same weight as 0.9 respondents.

The following chart outlines the characteristics of the Metro Area sample for key demographic variables, compared to actual population characteristics revealed in census data. [Note that the sample consisted solely of area residents age 18 and older; data on children were given by proxy by the person most responsible for that child's health care needs, and these children are not represented demographically in this chart.]



Population & Survey Sample Characteristics (Metro Area, 2024)



Sources:

- US Census Bureau, 2016-2020 American Community Survey.
- 2024 PRC Community Health Survey, PRC, Inc.

Notes:

"Low Income" reflects those living under 200% of the federal poverty level, based on guidelines established by the US Department of Health & Human Services.
 All Hispanic respondents are grouped, regardless of identity with any other race group. Race reflects those who identify with a single race category, without Hispanic origin. "Diverse Races" includes those who identify as Black or African American, American Indian or Alaska Native, Asian, Native Hawaiian/Pacific Islander, or as being of multiple races, without Hispanic origin.

The sample design and the quality control procedures used in the data collection ensure that the sample is representative. Thus, the findings may be generalized to the total population of community members in the defined area with a high degree of confidence.

Online Key Informant Survey

To solicit input from key informants, those individuals who have a broad interest in the health of the community, an Online Key Informant Survey also was implemented as part of this process. A list of recommended participants was provided by the sponsoring and supporting organizations; this list included names and contact information for physicians, public health representatives, other health professionals, social service providers, and a variety of other community leaders. Potential participants were chosen because of their ability to identify primary concerns of the populations with whom they work, as well as of the community overall.

Key informants were contacted by email, introducing the purpose of the survey and providing a link to take the survey online; reminder emails were sent as needed to increase participation. In all, 118 community representatives took part in the Online Key Informant Survey, as outlined in the table that follows:

ONLINE KEY INFORMANT	SURVEY PARTICIPATION
KEY INFORMANT TYPE	NUMBER PARTICIPATING
Physicians	22
Public Health Representatives	4
Other Health Providers	42
Social Services Providers	21
Business Leaders	12
Other Community Leaders	17



Through this process, input was gathered from individuals whose organizations work with low-income, minority, or other medically underserved populations. Final participation included representatives of the organizations outlined below.

- All Care Health Center
- Board of Health
- Catholic Charities of Omaha
- Center for Holistic Development
- Charles Drew Health Center, Inc.
- CHI Health
- City of Council Bluffs
- City of Omaha Planning
- City of Papillion
- City of Plattsmouth
- Claire Memorial United Methodist Church
- College of St. Mary
- Connections Area Agency on Aging
- Douglas County
- Douglas County Board of Health
- Eastern Nebraska Office on Aging
- FAMILY, Inc.
- Family, MCH Sealant Program
- Goodwill Omaha
- Heartland Family Service
- Housing Foundation for Sarpy County
- I Be Black Girl
- Ignite Nebraska
- Immanuel Pathways Southwest Iowa (PACE)
- Intercultural Senior Center
- Iowa Child Care Resource & Referral
- Iowa West Foundation
- Lift Up Sarpy
- Methodist-EAP
- Methodist Community Health Clinic (MCHC)

- Methodist–MHS
- Methodist–MJE
- Methodist–MWH
- Methodist–NMC
- Methodist-NMH
- Methodist–NMH Admin Services
- Methodist–NMH Cancer Center
- Methodist–Kountze Commons
- Metro Area Transit
- Metropolitan Area Planning Agency
- Nebraska Medicine
- New Life Family Alliance
- No More Empty Pots
- Omaha for Us
- OneWorld Community Health Center
- Pottawattamie County
- Region 6
- ReImagine Omaha
- Santa Monica House
- Sarpy County
- Sarpy/Cass Health
- Serenity Dental
- Southwest Iowa MHDS
- Stephen Center
- UnitedHealth care
- University of Nebraska Medical Center College of Public Health
- Visiting Nurse Association
- Women's Fund of Omaha
- YMCA



In the online survey, key informants were asked to rate the degree to which various health issues are a problem in their own community. Follow-up questions asked them to describe why they identify problem areas as such and how these might better be addressed. Results of their ratings, as well as their verbatim comments, are included throughout this report as they relate to the various other data presented.

Public Health, Vital Statistics & Other Data

A variety of existing (secondary) data sources was consulted to complement the research quality of this Community Health Needs Assessment. Data for the Metro Area were obtained from the following sources (specific citations are included with the graphs throughout this report):

- Center for Applied Research and Engagement Systems (CARES), University of Missouri Extension, SparkMap (sparkmap.org)
- Centers for Disease Control & Prevention, Office of Infectious Disease, National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention
- Centers for Disease Control & Prevention, Office of Public Health Science Services, National Center for Health Statistics
- National Cancer Institute, State Cancer Profiles
- US Census Bureau, American Community Survey
- US Census Bureau, County Business Patterns
- US Census Bureau, Decennial Census
- US Department of Agriculture, Economic Research Service
- US Department of Health & Human Services
- US Department of Health & Human Services, Health Resources and Services Administration (HRSA)
- US Department of Justice, Federal Bureau of Investigation
- US Department of Labor, Bureau of Labor Statistics

Benchmark Comparisons

Trending

Similar surveys were administered in the Metro Area in 2011, 2015, 2018, and 2021 by PRC. Trending data, as revealed by comparison to prior survey results, are provided throughout this report whenever available. Historical data for secondary data indicators are also included for the purposes of trending.

Nebraska & Iowa Data

State-level findings are provided where available as an additional benchmark against which to compare local findings. For survey indicators, these are taken from the most recently published data from the CDC's Behavioral Risk Factor Surveillance System (BRFSS). For other indicators, these draw from vital statistics, census, and other existing data sources.

National Data

National survey data, which are also provided in comparison charts, are taken from the 2023 PRC National Health Survey; these data may be generalized to the US population with a high degree of confidence. National-level findings (from various existing resources) are also provided for comparison of secondary data indicators.



Healthy People 2030 Objectives

Healthy People provides 10-year, measurable public health objectives — and tools to help track progress toward achieving them. Healthy People identifies public health priorities to help individuals, organizations, and communities across the United States improve health and well-being. Healthy People 2030, the initiative's fifth iteration, builds on knowledge gained over the first four decades.



The Healthy People 2030 framework was based on recommendations made by the Secretary's Advisory Committee on National Health Promotion and Disease Prevention Objectives for 2030. After receiving feedback from individuals and organizations and input from subject matter experts, the US Department of Health and Human Services (HHS) approved the framework which helped guide the selection of Healthy People 2030 objectives.

Determining Significance

Differences noted in this report represent those determined to be significant. For survey-derived indicators (which are subject to sampling error), statistical significance is determined based on confidence intervals (at the 95 percent confidence level), using question-specific samples and response rates. For the purpose of this report, "significance" of secondary data indicators (which do not carry sampling error but might be subject to reporting error) is determined by a 15% variation from the comparative measure.

Information Gaps

While this assessment is quite comprehensive, it cannot measure all possible aspects of health in the community, nor can it adequately represent all possible populations of interest. It must be recognized that these information gaps might in some ways limit the ability to assess all of the community's health needs.

For example, certain population groups — such as the homeless, institutionalized persons, or those who only speak a language other than English or Spanish — are not represented in the survey data. Other population groups — for example, pregnant women, undocumented residents, and members of certain racial/ethnic or immigrant groups — while included in the overall findings, might not be individually identifiable or might not comprise a large-enough sample for independent analyses.

In terms of content, this assessment was designed to provide a comprehensive and broad picture of the health of the overall community. However, there are certainly medical conditions that are not specifically addressed.

Public Comment

Participating hospitals and health systems made their prior Community Health Needs Assessment (CHNA) reports publicly available through their respective websites; through that mechanism, they requested from the public written comments and feedback regarding the CHNA and implementation strategies. At the time of this writing, none had not received any written comments. However, through population surveys and key informant feedback for this assessment, input from the broader community was considered and taken into account when identifying and prioritizing the significant health needs of the community. Participating hospitals will continue to use their websites as tools to solicit public comments and ensure that these comments are considered in the development of future CHNAs.



SUMMARY OF FINDINGS

Significant Health Needs of the Community

The following "Areas of Opportunity" represent the significant health needs of the community, based on the information gathered through this Community Health Needs Assessment. From these data, opportunities for health improvement exist in the area with regard to the following health issues (see also the summary tables presented in the following section).

The Areas of Opportunity were determined after consideration of various criteria, including: standing in comparison with benchmark data (particularly national data); identified trends; the preponderance of significant findings within topic areas; the magnitude of the issue in terms of the number of persons affected; and the potential health impact of a given issue. These also take into account those issues of greatest concern to the key informants giving input to this process.

AREAS OF OPPORTU	NITY IDENTIFIED THROUGH THIS ASSESSMENT
ACCESS TO HEALTH CARE SERVICES	 Barriers to Access Inconvenient Office Hours Cost of Prescriptions Cost of Physician Visits Appointment Availability Difficulty Finding a Physician Lack of Transportation Culture/Language Skipping/Stretching Prescriptions Emergency Room Utilization
CANCER	 Leading Cause of Death Prostate Cancer Deaths Cervical Cancer Screening
DIABETES	 Prevalence of Borderline/Pre-Diabetes Key Informants: <i>Diabetes</i> ranked as a top concern.
DISABLING CONDITIONS	Activity LimitationsAlzheimer's Disease Deaths
HEART DISEASE & STROKE	 Leading Cause of Death Heart Disease Prevalence Stroke Prevalence
INFANT HEALTH & FAMILY PLANNING	■ Prenatal Care



-continued on the following page-

AREA	S OF OPPORTUNITY (continued)
INJURY & VIOLENCE	Falls [45+]Intimate Partner Violence
MENTAL HEALTH	 "Fair/Poor" Mental Health Diagnosed Depression Symptoms of Chronic Depression Stress Current Anxiety & Depression Suicide Deaths Lack of Social Support Receiving Treatment for Mental Health Difficulty Obtaining Mental Health Services Key Informants: Mental Health ranked as a top concern.
NUTRITION, PHYSICAL ACTIVITY & WEIGHT	 Leisure-Time Physical Activity Meeting Physical Activity Guidelines Built Environment Overweight & Obesity Key Informants: Nutrition, Physical Activity & Weight ranked as a top concern.
ORAL HEALTH	Regular Dental Care
RESPIRATORY DISEASE	Asthma Prevalence
SOCIAL DETERMINANTS OF HEALTH	 Financial Resilience Housing Insecurity Housing Conditions Loss of Utilities Discrimination Adverse Childhood Experiences (ACEs) Key Informants: Social Determinants of Health ranked as a top concern.
SUBSTANCE USE	 Alcohol-Induced Deaths Unintentional Drug-Induced Deaths Key Informants: Substance Use ranked as a top concern.
TOBACCO USE	 Use of Vaping Products



Community Feedback on Prioritization of Health Needs

Prioritization of the health needs identified in this assessment ("Areas of Opportunity" above) was determined based on a prioritization exercise conducted among providers and other community leaders (representing a cross-section of community-based agencies and organizations) as part of the Online Key Informant Survey.

In this process, these key informants were asked to rate the severity of a variety of health issues in the community. Insofar as these health issues were identified through the data above and/or were identified as top concerns among key informants, their ranking of these issues informed the following priorities:

- 1. Mental Health
- 2. Social Determinants of Health
- 3. Diabetes
- 4. Nutrition, Physical Activity & Weight
- 5. Substance Use
- 6. Heart Disease & Stroke
- 7. Infant Health & Family Planning
- 8. Injury & Violence
- 9. Disabling Conditions
- 10. Oral Health
- 11. Cancer
- 12. Access to Health Care Services
- 13. Tobacco Use
- 14. Respiratory Diseases

Hospital Implementation Strategy

Sponsoring hospitals will use the information from this Community Health Needs Assessment to develop Implementation Strategies to address the significant health needs in the community. While the hospitals will likely not implement strategies for all of the health issues listed above, the results of this prioritization exercise will be used to inform the development of action plans to guide community health improvement efforts in the coming years.

Note: An evaluation of the hospital's past activities to address the needs identified in prior CHNAs can be found at the end of this report.



Summary Tables: Comparisons With Benchmark Data

Reading the Summary Tables

- In the following tables, Metro Area results are shown in the larger, gray column.
- The group of columns furthest to the left provide comparisons among the five subareas within Douglas County, identifying differences for each as "better than" (*), "worse than" (*), or "similar to" (*) the combined opposing areas of the county.
- The second grouping of columns [to the left of the Metro Area column] provide comparisons among the four counties assessed, identifying differences for each as "better than" (⑤), "worse than" (⑥), or "similar to" (⑥) the combined opposing counties.
- The columns to the right of the Metro Area column provide trending, as well as comparisons between local data and any available state and national findings, and Healthy People 2030 objectives. Again, symbols indicate whether the Metro Area compares favorably (♠), unfavorably (♠), or comparably (△) to these external data.

Note that blank table cells signify that data are not available or are not reliable for that area and/or for that indicator.

Tip: Indicator labels beginning with a "%" symbol are taken from the PRC Community Health Survey; the remaining indicators are taken from secondary data sources.

TREND SUMMARY

(Current vs. Baseline Data)

SURVEY DATA INDICATORS:

Trends for survey-derived indicators represent significant changes since 2011 (or earliest available data). Note that survey data reflect the ZIP Codedefined Metro Area.

OTHER (SECONDARY) DATA INDICATORS:

Trends for other indicators (e.g., public health data) represent point-to-point changes between the most current reporting period and the earliest presented in this report (typically representing the span of roughly a decade). Local secondary data reflect county-level data.



	DISF	PARITY WI	THIN DOL	JGLAS CC	UNTY	DISPARI	TY AMONO	G METRO (COUNTIES	Matua		METRO vs. BENCHMARKS						
SOCIAL DETERMINANTS	NE Omaha	SE Omaha	NW Omaha	SW Omaha	Western Douglas	Douglas County	Sarpy County	Cass County	Pott. County	Metro Area	vs. NE	vs. IA	vs. US	vs. HP2030	TREND			
Linguistically Isolated Population (Percent)						3.1	£ 1.2	0.3	£ 1.5	2.5	2.6	1.9	3.9					
Population in Poverty (Percent)						11.0	5.5	5.6	11.3	9.7	<i>≦</i> 10.4	<i>≦</i> 11.1	12.5	8.0				
Children in Poverty (Percent)						13.2	5.8	5.4	15.6	11.6	12.0	13.0	16.7	8.0				
No High School Diploma (Age 25+, Percent)						8.6	4.3	4.6	9.4	7.7	8.1	7.0	10.9					
Unemployment Rate (Age 16+, Percent)						2.6	<i>≦</i> 2.2	<i>€</i> 3	2.9	2.5	2.3	2.9	3.6		4.0			
% Unable to Pay Cash for a \$400 Emergency Expense	31.8	38.4	19.6	17.9	15.9	24.9	19.5	<i>≦</i> 20.3	31.1	24.3		-	34.0		18.7			
% Worry/Stress Over Rent/Mortgage in Past Year	38.3	39.5	<i>≦</i> 32.4	29.9	26.4	33.8	29.0	<i>≦</i> ≳ 28.9	<i>≦</i> 32.3	32.5			45.8		20.1			
% Unhealthy/Unsafe Housing Conditions	21.8	21.2	9.3	10.0	6.4	14.1	8.9	<i>€</i> 3 10.1	<i>≦</i> ≒ 15.8	13.1			16.4		6.1			
% Went Without Electricity, Water, or Heat	12.4	14.9	£ 12.5	6.4	£	11.2	6.9	8.3	<i>≦</i> 3 12.2	10.3					5.2			
Population With Low Food Access (Percent)						11.9	36.7	18.4	30.6	19.3	£ 21.9	20.0	22.2					
% Worried About Food in the Past Year	38.6	37.7	20.6	19.6	17.1	27.0	19.8	<i>₽</i> 22.1	<i>≦</i> 3 29.0	25.6			40.7		18.8			
% Ran Out of Food in the Past Year	30.6	33.4	15.2	15.9	10.7	21.7	15.3	<i>€</i> 3 17.8	24.7	20.6			32.6					

	DISF	PARITY WI	THIN DOL	JGLAS CC	UNTY	DISPARI	TY AMONO	METRO (COUNTIES			METRO	O vs. BENC	CHMARKS	
SOCIAL DETERMINANTS (continued)	NE Omaha	SE Omaha	NW Omaha	SW Omaha	Western Douglas	Douglas County	Sarpy County	Cass County	Pott. County	Metro Area	vs. NE	vs. IA	vs. US	vs. HP2030	TREND
% Food Insecure	41.2	42.2	21.9	20.8	18.7	29.1	21.8	<i>≦</i> 3.8	31.9	27.8			43.3		
% Treated With Less Respect Than Others	35.4	<i>≦</i> 32.0	<i>≦</i> 25.9	<i>⊆</i> 25.7	25.4	<i>≅</i> 28.9	<i>≅</i> 25.5	<i>∕</i> ≃ 26.4	<i>∕</i> ⊆ 29.5	28.2					25.1
% Receive Poorer Treatment at Restaurants/Stores	17.2	£	6.7	9.2	<i>₹</i> 3	10.8	8.0	6.7	<i>≦</i> 3 10.6	10.1					7.7
% Treated as Less Intelligent	<i>≦</i> 20.9	23.5	13.2	<i>≦</i> 17.2	<i>≦</i> 3 15.7	<i>€</i> 3 18.0	<i>≦</i> 3 17.0	£	<i>€</i> 3 19.6	17.8					13.3
% Threatened or Harassed	8.4	9.0	8.3	<i>€</i> 3 8.9	5.2	8.4	5.0	4.7	<i>€</i> 3 6.9	7.4					4.8
% Disagree That the Community Welcomes All Races	£	£	£ 11.3	£	<i>₽</i> 3 12.6	12.1	7.3	£	<i>≦</i> 3 10.4	11.0					<i>≦</i> 3 11.3
% Treated as Someone to Fear	11.9	<i>≦</i> 10.6	<i>€</i> 8.2	<i>₹</i> 3	4.1	<i>₽</i> 3	6.8	9.0	<i>≦</i> 3 10.2	8.7					<i>₹</i> 3 7.4
% 4+ Adverse Childhood Experiences (High ACEs Score)	30.4	<i>≦</i> 3 28.6	<i>≨</i> 3 24.4	20.4	<i>≨</i> ≳ 22.9	<i>≨</i> ≳ 25.1	19.9	<i>≦</i> 23.7	29.5	24.4			£ 25.5		15.1
						r areas combine ample sizes are						better	similar	worse	
			THIN DOU	JGLAS CC	UNTY	DISPARI	TY AMONO	METRO C	COUNTIES	Metro		METRO	O vs. BENC	CHMARKS	
OVERALL HEALTH	NE Omaha	SE Omaha	NW Omaha	SW Omaha	Western Douglas	Douglas County	Sarpy County	Cass County	Pott. County	Area	vs. NE	vs. IA	vs. US	vs. HP2030	TREND
% "Fair/Poor" Overall Health										16.3					
	17.5	20.6	15.7	9.1	12.1	14.9	16.5	11.8	24.9		15.0	16.2	15.7		12.7

better similar

worse

	DISF	PARITY WI	THIN DOL	JGLAS CC	UNTY	DISPARI	TY AMONO	G METRO (COUNTIES		METRO vs. BENCHMARKS						
ACCESS TO HEALTH CARE	NE Omaha	SE Omaha	NW Omaha	SW Omaha	Western Douglas	Douglas County	Sarpy County	Cass County	Pott. County	Metro Area	vs. NE	vs. IA	vs. US	vs. HP2030	TREND		
% [Age 18-64] Lack Health Insurance	<i>≦</i> 3 11.1	16.5	4.7	5.2	4.6	8.5	<i>∕</i> ≤ 7.0	<i>∕</i> ≏ 7.6	4.9	7.8	10.8	6.7	<i>≊</i> 8.1	<i>₹</i> 3 7.6	12.1		
% Difficulty Accessing Health Care in Past Year (Composite)	<i>≦</i> 3.6	<i>€</i> 3 42.8	<i>€</i> 3 42.6	<i>€</i> 3.5	<i>≅</i> 39.1	<i>€</i> 3 42.8	<i>€</i> 3 41.7	<i>≦</i> 39.4	<i>≦</i> 39.8	42.2			52.5		33.4		
% Cost Prevented Physician Visit in Past Year	43.0 23 18.6	42.0 23 19.9	42.0 23	43.3 2 14.9	11.5	42.0 2 16.9	15.1	29.4 25.0	29.0 26 16.0	16.4	10.2	7.2	21.6		14.5		
% Cost Prevented Getting Prescription in Past Year	18.8	18.2	18.8	16.3	11.6	17.5	12.4	13.9	18.1	16.4	10.2	1.2	20.2		14.3		
% Difficulty Getting Appointment in Past Year	21.8	21.5	20.5	22.3	20.9	21.5	20.3	17.7	20.9	21.1			33.4		10.5		
% Inconvenient Hrs Prevented Dr Visit in Past Year	19.0	16.2	20.7	15.5	15.5	£3.7 ± 17.7	25.5 17.9	19.1	£3 18.0	17.9			22.9		12.5		
% Difficulty Finding Physician in Past Year	16.0	16.7	£ 11.1	9.2	5.5	<i>≦</i> 3 12.2	£ 11.1	10.2	11.8	11.9			22.0		6.6		
% Transportation Hindered Dr Visit in Past Year	16.7	17.8	7.8	6.1	5.0	10.8	5.7	4.8	<i>≦</i> 3	9.7			18.3		4.7		
% Language/Culture Prevented Care in Past Year	3.7	4.4	£	1.6	<i>₽</i> 3 1.7	2.6	0.7	0.5	£	2.0			5.0		0.9		
% Stretched Prescription to Save Cost in Past Year	21.2	<i>≦</i> 18.5	<i>≦</i> 18.4	<i>≅</i> 15.4	11.4	<i>☆</i>	£ 15.3	<i>≦</i> 3.8	<i>≦</i> 17.5	17.0			<i>≦</i> 3 19.4		13.6		
% Treated Worse Than Other Races (Health Care)	8.5	5.6	3.7	2.7	3.0	4.8	2.1	£	4.0	4.1			6.1		4.3		
Primary Care Doctors per 100,000			•		0.0	149.7	<i>≦</i> 3 52.5	7.5	<i>€</i> 3 48.0	114.1	98.3	<i>≦</i> 109.7	113.2				

	DISF	PARITY WI	THIN DOL	JGLAS CC	UNTY	DISPARI	TY AMONO	G METRO C	COUNTIES			METRO vs. BENCHMARKS				
ACCESS TO HEALTH CARE (continued)	NE Omaha	SE Omaha	NW Omaha	SW Omaha	Western Douglas	Douglas County	Sarpy County	Cass County	Pott. County	Metro Area	vs. NE	vs. IA	vs. US	vs. HP2030	TREND	
% Have a Specific Source of Ongoing Care	71.5	<i>₹</i> 3 72.1	<i>₹</i> 3 77.4	7 9.8	<i>₹</i> 3 78.6	<i>₹</i> 3 76.0	<i>₹</i> 3 77.2	<i>₹</i> 76.8	<i>∕</i> ≈ 78.6	76.6			6 9.9	84.0	66.1	
% Routine Checkup in Past Year	<i>₹</i> 70.4	67.2	<i>∕</i> ≤ 72.5	<i>∕</i> ≤ 73.5	<i>∕</i> ≤3 74.7	<i>₹</i> 3 71.5	<i>∕</i> ⊆ 70.5	∕≘ 71.1	<i>∕</i> ⊆ 72.8	71.5	74.7	78.3	65.3		66.8	
% Two or More ER Visits in Past Year	18.3	£	<i>⊊</i> 3 9.5	8.3	<i>€</i> 3 9.0	<i>≦</i> 3 11.8	<i>≦</i> 11.2	10.3	<i>≦</i> 13.1	11.8			15.6		4.9	

	쏨	937
etter	similar	wors

	DISF	PARITY WI	THIN DOL	JGLAS CC	UNTY	DISPARI	TY AMONG	G METRO (COUNTIES		METRO vs. BENCHMARKS					
CANCER	NE Omaha	SE Omaha	NW Omaha	SW Omaha	Western Douglas	Douglas County	Sarpy County	Cass County	Pott. County	Metro Area	vs. NE	vs. IA	vs. US	vs. HP2030	TREND	
Cancer Deaths per 100,000 (Age-Adjusted)										154.6						
						154.1	144.9	147.1	174.7		148.5	151.3	146.5	122.7	178.5	
Lung Cancer Deaths per 100,000 (Age-Adjusted)										34.3	给		含			
(go : squares)											31.8	36.3	33.4	25.1		
Female Breast Cancer Deaths per 100,000 (Age-Adjusted)										19.6		É				
											20.8	17.9	19.4	15.3		
Prostate Cancer Deaths per 100,000 (Age-Adjusted)										21.9	给	会	***	***		
											18.7	20.2	18.5	16.9		
Colorectal Cancer Deaths per 100,000 (Age-Adjusted)										13.4	给					
											14.9	13.9	13.1	8.9		
Cancer Incidence per 100,000 (Age-Adjusted)						会				483.1						
						484.5	473.2	490.4	489.0		459.1	486.8	442.3			

	DISF	PARITY W	ITHIN DOL	JGLAS CO	UNTY	DISPARI	TY AMONO	G METRO (COUNTIES			METRO	O vs. BENC	CHMARKS	
CANCER (continued)	NE Omaha	SE Omaha	NW Omaha	SW Omaha	Western Douglas	Douglas County	Sarpy County	Cass County	Pott. County	Metro Area	vs. NE	vs. IA	vs. US	vs. HP2030	TREND
Lung Cancer Incidence per 100,000 (Age-Adjusted)										60.1					
						59.2	52.6	66.7	72.9		52.3	60.7	54.0		,
Female Breast Cancer Incidence per 100,000 (Age-Adjusted)							ớ	给		140.9					
						144.5	141.0	124.7	127.5		131.0	134.7	127.0		
Prostate Cancer Incidence per 100,000 (Age-Adjusted)								会		128.0					
						131.2	130.9	120.5	111.3		124.8	120.4	110.5		
Colorectal Cancer Incidence per 100,000 (Age-Adjusted)										39.0		É			
						38.0	36.6	39.3	47.1		40.5	40.7	36.5		
% Cancer						给				8.8					
	8.5	8.2	8.3	9.6	4.9	8.4	7.9	13.2	11.7		11.1	12.3	7.4		9.2
% [Women 50-74] Breast Cancer Screening										82.2					
	77.9	78.1	83.2	85.4	92.5	82.9	82.4	78.2	79.7		76.9	79.6	64.0	80.5	82.3
% [Women 21-65] Cervical Cancer Screening										73.5					
	75.2	68.0	75.6	75.3	74.0	73.9	73.2	61.9	74.6				75.4	84.3	86.7
% [Age 50-75] Colorectal Cancer Screening		给	给			给				78.9					
	75.6	71.4	81.4	82.2	74.6	78.4	80.5	83.8	77.4		68.3	72.0	71.5	74.4	75.3

	DISF	PARITY WI	THIN DOL	JGLAS CC	UNTY	DISPARI	TY AMONO	G METRO (COUNTIES			METR	O vs. BENC	CHMARKS	
DIABETES	NE Omaha	SE Omaha	NW Omaha	SW Omaha	Western Douglas	Douglas County	Sarpy County	Cass County	Pott. County	Metro Area	vs. NE	vs. IA	vs. US	vs. HP2030	TREND
Diabetes Deaths per 100,000 (Age-Adjusted)							给			25.9					
						16.0	28.0	24.2	31.1		25.1	22.3	22.6		22.7
% Diabetes/High Blood Sugar						给				11.2					给
	15.4	10.7	10.0	9.5	7.4	11.0	11.9	10.1	11.6		10.8	11.6	12.8		10.6
% Borderline/Pre-Diabetes	给		给			给	给		会	12.1					
	13.2	13.3	13.3	9.2	11.3	12.0	11.2	12.1	14.2				15.0		8.8
Kidney Disease Deaths per 100,000 (Age- Adjusted)						绘			会	10.2					
						11.4	7.9		9.9		10.3	9.7	12.8		11.6

better

similar worse

	DISF	PARITY W	ITHIN DOL	JGLAS CC	UNTY	DISPARI	TY AMON	G METRO (COUNTIES			METRO	O vs. BENC	CHMARKS	
DISABLING CONDITIONS	NE Omaha	SE Omaha	NW Omaha	SW Omaha	Western Douglas	Douglas County	Sarpy County	Cass County	Pott. County	Metro Area	vs. NE	vs. IA	vs. US	vs. HP2030	TREND
% Activity Limitations										29.6					***
	34.1	29.6	24.0	29.2	23.0	28.5	29.1	27.0	37.0				27.5		18.4
Alzheimer's Disease Deaths per 100,000 (Age- Adjusted)						会	ح		会	38.5	\$600	\$400	***		
						37.7	41.7	31.7	39.7		30.0	30.9	30.9		28.1

Note: In the section above, each subarea is compared against all other areas combined. Throughout these tables, a blank or empty cell indicates that data are not available for this indicator or that sample sizes are too small to provide meaningful results.

better similar

	DISF	PARITY WI	THIN DOL	JGLAS CC	UNTY	DISPARI	TY AMONO	G METRO (COUNTIES			METR	O vs. BENO	CHMARKS	
HEART DISEASE & STROKE	NE Omaha	SE Omaha	NW Omaha	SW Omaha	Western Douglas	Douglas County	Sarpy County	Cass County	Pott. County	Metro Area	vs. NE	vs. IA	vs. US	vs. HP2030	TREND
Heart Disease Deaths per 100,000 (Age-Adjusted)						£ 132.6	<i>≅</i> 130.9	<i>∕</i> ≘ 167.7	180.3	139.7	£ 144.8	170.3	164.4	<i>≦</i> 127.4	£ 151.3
% Heart Disease	<i>≦</i> 10.7	<i>€</i> 3 8.1	<i>€</i> 3 6.9	<i>€</i> 3 6.9	£	<i>€</i> 8.2	<i>€</i> 3 6.9	<i>₹</i> 3 7.9	10.6	8.2	6.6	6.7	10.3		5.2
Stroke Deaths per 100,000 (Age-Adjusted)						<i>≦</i> ≘ 36.8	<i>≦</i> 31.4	<i>⊆</i> 29.1	<i>∽</i> 32.3	34.8	<i>≦</i> 33.0	<i>≦</i> 32.3	<i>≦</i> 37.6	<i>∽</i> 33.4	<i>∽</i> 38.2
% Stroke	7.1	<i>≦</i> 3.4	3.6	2.5	1.6	3.8	1.8	<i>≦</i> ≘ 3.3	<i>∽</i> 3.2	3.3	2.6	<i>≦</i> 3.1	5.4		2.3

petter similar wors

better similar worse

	DISF	PARITY WI	THIN DOL	JGLAS CC	UNTY	DISPARI	TY AMONG	G METRO (COUNTIES			METR	O vs. BENC	CHMARKS	
INFANT HEALTH & FAMILY PLANNING	NE Omaha	SE Omaha	NW Omaha	SW Omaha	Western Douglas	Douglas County	Sarpy County	Cass County	Pott. County	Metro Area	vs. NE	vs. IA	vs. US	vs. HP2030	TREND
No Prenatal Care in First 6 Months (Percent of Births)						5.7	3.3			5.1	4.9	4.3	6.1		4.0
Teen Births per 1,000 Females 15-19						18.4	9.5	10.9	20.0	16.5	£ 16.0	14.4	£ 16.6		
Low Birthweight (Percent of Births)						9.0	<i>₹</i> 3 7.0	<i>₹</i> 3 7.0	<i>€</i> 3 8.0	8.2	7.5	6.8	8.3		
Infant Deaths per 1,000 Births						<i>€</i> 3 6.1	3.8		7.5	5.8	5.4	4.8	£ 5.5	5.0	5.2

Note: In the section above, each subarea is compared against all other areas combined. Throughout these tables, a blank or empty cell indicates that data are not available for this indicator or that sample sizes are too small to provide meaningful results.

better similar worse

	DISF	PARITY W	ITHIN DOL	JGLAS CC	UNTY	DISPARI	TY AMONO	G METRO C	COUNTIES			METR	O vs. BENC	CHMARKS	
INJURY & VIOLENCE	NE Omaha	SE Omaha	NW Omaha	SW Omaha	Western Douglas	Douglas County	Sarpy County	Cass County	Pott. County	Metro Area	vs. NE	vs. IA	vs. US	vs. HP2030	TREND
Unintentional Injury Deaths per 100,000 (Age-Adjusted)						☆	£		£3	36.5	£	***	*		£3
Motor Vehicle Crash Deaths per 100,000 (Age-Adjusted)						35.6 23 8.4	36.0 2 9.4	46.4	40.4	9.5	40.3	42.9 2 10.5	51.6 11.4	43.2	32.5
[65+] Fall-Related Deaths per 100,000 (Age-Adjusted)						59.3	80.0		63.6	63.6	67.8	87.4	67.1	63.4	
% [Age 45+] Fell in the Past Year	<i>☎</i> 37.0	<i>≦</i> 33.9	<i>≦</i> 35.7	<i>∽</i> 34.7	<i>≦</i> 34.8	<i>≦</i> 35.3	34.6	<i>≦</i> 31.8	<i>≦</i> 37.2	35.3					30.1
Homicide Deaths per 100,000 (Age-Adjusted)										4.4	3.0	3.0	6.1	5.5	6.2
Violent Crimes per 100,000						493.5	94.7	108.6	249.8	369.3	286.4	283.0	416.0		
% Neighborhood Is "Slightly/Not At All Safe"	42.0	39.6	14.4	8.9	4.7	22.6	8.2	5.3	<i>≅</i> 21.8	18.9					<i>≦</i> 3 17.4
% Victim of Intimate Partner Violence	24.7	24.3	16.2	18.0	<i>€</i> 3 18.6	20.2	15.7	£ 16.3	26.5	19.8			<i>€</i> 3 20.3		12.0

£

etter si

worse

	DISF	PARITY WI	THIN DOL	JGLAS CC	DUNTY	DISPARI	TY AMONG	METRO (COUNTIES			METR	O vs. BENC	CHMARKS	
MENTAL HEALTH	NE Omaha	SE Omaha	NW Omaha	SW Omaha	Western Douglas	Douglas County	Sarpy County	Cass County	Pott. County	Metro Area	vs. NE	vs. IA	vs. US	vs. HP2030	TREND
% "Fair/Poor" Mental Health						给		给		22.7					
	23.2	27.5	22.9	20.6	17.5	22.7	20.9	20.4	26.2				24.4		9.0
% Diagnosed Depression	给									32.3					
	33.6	35.9	30.9	34.6	29.4	33.3	28.3	29.0	34.4		17.0	18.5	30.8		19.5
% Symptoms of Chronic Depression			给	给					给	41.8					** **********************************
	47.4	49.4	39.3	40.2	38.3	43.0	37.1	35.0	44.9				46.7		25.1
% Typical Day Is "Extremely/Very" Stressful		会	含	给		会				18.1					
	19.1	16.9	17.4	19.7	16.2	18.2	16.4	13.5	22.1				21.1		11.5
Suicide Deaths per 100,000 (Age-Adjusted)						会	给			14.1			给		
						13.8	12.4		19.0		14.8	16.7	13.9	12.8	10.1
% Have Someone to Turn to All/Most of the Time										74.6					
	70.7	59.7	76.3	79.8	81.6	73.5	79.1	77.1	72.9						86.1
% Recent Anxiety				给						25.9					
	29.7	27.4	25.1	24.6	23.0	26.2	24.4	17.7	28.9						20.0
% Recent Depression				给						18.9					
	25.1	20.1	17.4	20.0	15.9	20.1	14.9	10.6	21.2						15.1
% Moderate to Severe Anxiety/Depression (PHQ-4 Score of 6+)										20.0					
(1119-4 30016 0101)	25.8	20.4	20.3	19.6	17.1	21.1	15.6	12.4	24.2						15.6
Mental Health Providers per 100,000										214.5					
						280.6	72.4	33.8	143.1		184.5	136.7	183.8		
% Receiving Mental Health Treatment										28.6					
	28.2	25.8	31.7	29.9	22.2	28.7	27.3	23.2	32.1				21.9		14.4

	DISP	ARITY WI	THIN DOU	IGLAS CO	UNTY	DISPARI	TY AMONG	METRO C	COUNTIES			METRO	O vs. BENC	HMARKS	
MENTAL HEALTH (continued)	NE Omaha	SE Omaha	NW Omaha	SW Omaha	Western Douglas	Douglas County	Sarpy County	Cass County	Pott. County	Metro Area	vs. NE	vs. IA	vs. US	vs. HP2030	TREND
% Unable to Get Mental Health Services in Past Year	16.6	£	£	£	*	10.0	70	£	£	11.8			£		2.7
	16.6	14.1	11.0	12.2	7.0	12.8	7.9	9.6	13.8				13.2		2.7

3,45		940
etter	similar	worse

	DISF	PARITY WI	THIN DOL	IGLAS CC	UNTY	DISPARI	TY AMONG	G METRO (COUNTIES			METR	O vs. BENC	HMARKS	
NUTRITION, PHYSICAL ACTIVITY & WEIGHT	NE Omaha	SE Omaha	NW Omaha	SW Omaha	Western Douglas	Douglas County	Sarpy County	Cass County	Pott. County	Metro Area	vs. NE	vs. IA	vs. US	vs. HP2030	TREND
% "Very/Somewhat" Difficult to Buy Fresh Produce	29.5	29.5	<i>≦</i> 24.0	17.4	16.6	<i>⊆</i> 23.8	\$\$ 19.4	<i>∕</i> ≘ 26.0	30.1	23.7			30.0		<i>≦</i> 22.8
% 7+ Sugar-Sweetened Drinks in Past Week	30.9	33.8	<i>≦</i> 3.4	22.3	<i>≦</i> 3 24.2	<i>≦</i> 3	£ 25.3	<i>≦</i> 3 28.2	31.1	26.8					28.3
% No Leisure-Time Physical Activity	£ 26.3	32.3	£ 22.5	<i>€</i> 3.2	£3 21.2	25.2	<i>₽</i> 27.1	<i>≦</i> 30.3	32.0	26.5	24.7	£ 25.9	30.2	21.8	16.7
% Meet Physical Activity Guidelines	21.8	<i>≦</i> 3 25.8	<i>≦</i> 3 28.8	31.5	33.8	27.9	<i>≅</i> 24.7	<i>≦</i> 21.1	22.1	26.4	20.9	20.1	30.3	29.7	22.0
Recreation/Fitness Facilities per 100,000						20.4			<i>≦</i> 13.9	17.5	14.2	12.1	14.8		
% Lack of Sidewalks/Poor Sidewalks Prevent Exercise	33.2	25.7	16.1	15.0	<i>€</i> 3 16.8	<i>≦</i> 3 21.2	12.5	40.6	26.3	20.6					<i>€</i> 3 20.1
% Lack of Trails/Poor Quality Trails Prevent Exercise	26.6	<i>≦</i> 21.7	14.7	13.9	13.6	18.2	12.5	23.9	<i>≦</i> 3 18.6	17.2					12.9
% Heavy Neighborhood Traffic Prevents Exercise	29.1	29.4	<i>△</i> 19.9	16.6	12.7	22.1	9.7	6.7	<i>≦</i> 3 20.3	18.9					16.7

	DISF	PARITY WI	THIN DOL	JGLAS CC	UNTY	DISPARI	TY AMONO	G METRO (COUNTIES	/		METR	O vs. BENO	CHMARKS	
NUTRITION, PHYSICAL ACTIVITY & WEIGHT (continued)	NE Omaha	SE Omaha	NW Omaha	SW Omaha	Western Douglas	Douglas County	Sarpy County	Cass County	Pott. County	Metro Area	vs. NE	vs. IA	vs. US	vs. HP2030	TREND
% Lack of Street Lights/Poor Street Lights Prevent Exercise	17.6	17.6	10.0	7.0	<i>€</i> 3 9.5	£ 12.1	7.0	21.3	17.6	12.0					9.4
% Crime Prevents Exercise in the Neighborhood	34.2	24.8	12.0	7.8	5.1	17.4	4.1	3.9	<i>≅</i> 17.0	14.2					11.0
% Overweight (BMI 25+)	<i>€</i> 3 69.7	<i>∕</i> ≤ 73.7	<i>∕</i> ≤ 70.1	<i>€</i> 3 67.2	<i>∕</i> ≤ 70.4	69.8	75.4	<i>∕</i> ≘ 70.1	80.1	72.2	70.4	<i>∕</i> ≃ 71.2	63.3		67.5
% Obese (BMI 30+)	<i>≦</i> 39.8	<i>≦</i> 36.7	<i>≨</i> 35.9	<i>≦</i> 32.6	<i>≅</i> 33.3	35.7	41.9	46.1	45.4	38.4	35.3	<i>≅</i> 37.4	33.9	36.0	30.3
% Have Received Professional Advice to Lose Weight	<i>≦</i> 24.3	<i>⊆</i> 25.9	<i>⊆</i> 26.8	<i>≅</i> 27.7	<i>≦</i> 24.5	<i>≘</i> 26.2	<i>≘</i> 22.8	<i>⊆</i> 24.1	<i>€</i> 3 25.4	25.3					<i>≦</i> 3 24.7

	给	100
etter	similar	wors

	DISPARITY WITHIN DOUGLAS COUNTY				DISPARITY AMONG METRO COUNTIES						METRO vs. BENCHMARKS					
ORAL HEALTH	NE Omaha	SE Omaha	NW Omaha	SW Omaha	Western Douglas	Douglas County	Sarpy County	Cass County	Pott. County	Metro Area	vs. NE	vs. IA	vs. US	vs. HP2030	TREND	
% Dental Visit in Past Year	60.5	57.7	<i>€</i> 3 68.7	71.2	78.8	<i>€</i> 3 66.4	74.1	<i>€</i> 3.5	61.8	67.4	€ 66.2	<i>€</i> 3 68.3	56.5	45.0	70.4	
	Note: In the section above, each subarea is compared against all other areas combined. Throughout these tables, a blank or empty												_			

lote: In the section above, each subarea is compared against all other areas combined. Throughout these tables, a blank or empty cell indicates that data are not available for this indicator or that sample sizes are too small to provide meaningful results.

	DISPARITY WITHIN DOUGLAS COUNTY				DISPARITY AMONG METRO COUNTIES					METRO vs. BENCHMARKS						
RESPIRATORY DISEASE	NE Omaha	SE Omaha	NW Omaha	SW Omaha	Western Douglas	Douglas County	Sarpy County	Cass County	Pott. County	Metro Area	vs. NE	vs. IA	vs. US	vs. HP2030	TREND	
Lung Disease Deaths per 100,000 (Age-Adjusted)										44.8						
. ,						45.4	36.6	49.0	52.3		45.7	42.3	38.1		50.4	
Pneumonia/Influenza Deaths per 100,000 (Age-Adjusted)						ح			***	13.7						
						13.2	12.9		18.7		14.2	13.8	13.4		14.7	
% Asthma	***									12.5	****	***				
	18.2	13.7	10.7	11.1	8.5	12.8	10.9	9.2	14.8		8.1	9.7	17.9		8.6	
% COPD (Lung Disease)						给				5.9						
	7.6	5.9	5.5	5.0	4.3	5.8	3.3	6.3	11.3		5.6	6.6	11.0		7.4	

better similar wo

worse

	DISPARITY WITHIN DOUGLAS COUNTY				DISPARITY AMONG METRO COUNTIES						METRO				
SEXUAL HEALTH	NE Omaha	SE Omaha	NW Omaha	SW Omaha	Western Douglas	Douglas County	Sarpy County	Cass County	Pott. County	Metro Area	vs. NE	vs. IA	vs. US	vs. HP2030	TREND
HIV Prevalence per 100,000						83.2	24.9	26.4	152.4	76.3	147.1	114.2	382.2		
Chlamydia Incidence per 100,000						653.2	406.9	174.0	<i>≦</i> 522.0	572.1	453.1	<i>€</i> 3 489.2	495.5		
Gonorrhea Incidence per 100,000						289.9	118.9	37.0	225.1	238.8	156.0	200.5	214.0		

Note: In the section above, each subarea is compared against all other areas combined. Throughout these tables, a blank or empty cell indicates that data are not available for this indicator or that sample sizes are too small to provide meaningful results.

£

\$47.

better similar

worse

	DISF	PARITY WI	THIN DOL	IGLAS CC	UNTY	DISPARI	TY AMONO	Metro		METRO vs. BENCHMARKS					
SUBSTANCE USE	NE Omaha	SE Omaha	NW Omaha	SW Omaha	Western Douglas	Douglas County	Sarpy County	Cass County	Pott. County	Area	vs. NE	vs. IA	vs. US	vs. HP2030	TREND
Alcohol-Induced Deaths per 100,000 (Age-Adjusted)						18.6	9.8		<i>≦</i> 3.7	15.8	12.0	9.9	11.9		8.5
% Excessive Drinking										22.7					
	26.5	24.0	21.6	25.8	27.9	24.7	18.3	21.3	19.8		20.5	22.6	34.3		26.0
Unintentional Drug-Induced Deaths per 100,000 (Age-Adjusted)										8.9	\$200				
						9.4	7.4		9.5		7.4	9.4	21.0		7.3
% Used a Prescription Opioid in Past Year									*** *********************************	13.4					
	17.2	9.8	14.5	11.4	12.4	13.2	11.5	17.5	16.9				15.1		18.1
% Ever Sought Help for Alcohol or Drug Problem			***		\$000 m		\$100	***		7.2					
	9.3	13.0	5.7	7.1	4.4	8.1	4.2	4.3	8.6				6.8		3.9

	23	\$45.
better	similar	worse


DISPARITY WITHIN DOUGLAS COUNTY **DISPARITY AMONG METRO COUNTIES** METRO vs. BENCHMARKS Metro NE SE NW SW Western Douglas Sarpy Cass Pott. VS. **TOBACCO USE** Area vs. US **TREND** County HP2030 Omaha Omaha Omaha Omaha **Douglas** County County County NE IA £ * % Smoke Cigarettes 紫 * 14.8 * \$100°C * **100 100** 13.3 13.3 17.0 20.7 20.1 12.0 10.7 15.6 10.1 19.5 13.0 14.7 23.9 6.1 % Someone Smokes at Home * * 11.4 * * * 888 B. (1) \$355 **\$35**1 19.2 16.4 10.0 12.5 5.9 12.6 14.7 7.9 9.2 17.7 15.1 给 93 % Use Vaping Products 11.3 * 1 15.6 12.0 10.0 9.9 8.5 6.7 18.5 16.5 9.6 8.4 10.8 5.8 11.8 % [Smokers] Have Quit Smoking 1+ Days in 49.3 给 给 给 B 255 Past Year 52.8 50.7 52.4 53.1 65.7

Note: In the section above, each subarea is compared against all other areas combined. Throughout these tables, a blank or empty cell indicates that data are not available for this indicator or that sample sizes are too small to provide meaningful results.

£

better similar

worse



COMMUNITY DESCRIPTION

POPULATION CHARACTERISTICS

Total Population

The four-county Metro Area, the focus of this Community Health Needs Assessment, encompasses 2,073 square miles and houses a total population of 895,395 residents, according to latest census estimates.

Total Population (Estimated Population, 2020)

	TOTAL POPULATION	TOTAL LAND AREA (square miles)	POPULATION DENSITY (per square mile)
Douglas County	584,526	326.41	1,791
Sarpy County	190,604	238.10	801
Cass County	26,598	557.34	48
Pottawattamie County	93,667	951.26	98
Metro Area	895,395	2,073.11	421
Nebraska	1,961,504	76,817.44	26
lowa	3,190,369	55,853.11	57
United States	331,449,281	3,533,018.38	94

Sources: • US Census Bureau American Community Survey 5-year estimates.

Population Change 2010-2020

A significant positive or negative shift in total population over time impacts health care providers and the utilization of community resources.

Between the 2010 and 2020 US Censuses, the population of the Metro Area increased by 101,045 persons, or 12.7%.


BENCHMARK ▶ The growth rate is much higher than recorded statewide and nationally.

DISPARITY ► The largest growth is seen in Sarpy and Douglas counties.




Center for Applied Research and Engagement Systems (CARES), University of Missouri Extension. Retrieved June 2024 via SparkMap (sparkmap.org).

Change in Total Population (Percentage Change Between 2010 and 2020)



US Census Bureau Decennial Census (2010-2020).
 Center for Applied Research and Engagement Systems (CARES), University of Missouri Extension. Retrieved June 2024 via SparkMap (sparkmap.org).
 A significant positive or negative shift in total population over time impacts health care providers and the utilization of community resources.

This map shows the areas of greatest increase or decrease in population between 2010 and 2020.





Urban/Rural Population

Urban areas are identified using population density, count, and size thresholds. Urban areas also include territory with a high degree of impervious surface (development). Rural areas are all areas that are not urban.


The Metro Area is predominantly urban, with 80.1% of the population living in areas designated as urban.

BENCHMARK ► Similar to the states' urban populations and higher than the nation's.

DISPARITY Note the disparity in urban/rural classifications between the four counties.

Urban and Rural Population (2020)

■ % Urban ■ % Rural



Sources:

- US Census Bureau Decennial Census.
- Center for Applied Research and Engagement Systems (CARES), University of Missouri Extension. Retrieved June 2024 via SparkMap (sparkmap.org).

This indicator reports the percentage of population living in urban and rural areas. Urban areas are identified using population density, count, and size thresholds.
 Urban areas also include territory with a high degree of impervious surface (development). Rural areas are all areas that are not urban.



Age


It is important to understand the age distribution of the population, as different age groups have unique health needs that should be considered separately from others along the age spectrum.

In the Metro Area, 25.6% of the population are children age 0-17; another 60.4% are age 18 to 64, while 14.0% are age 65 and older.

BENCHMARK > The Metro Area overall skews a bit younger than Nebraska, lowa, and the US.

DISPARITY ► By county, Sarpy has the largest proportion of children under 18 and the smallest proportion of adults age 65+.

Total Population by Age Groups (2020)




US Census Bureau American Community Survey 5-year estimates.

Center for Applied Research and Engagement Systems (CARES), University of Missouri Extension. Retrieved June 2024 via SparkMap (sparkmap.org).

Median Age

Douglas and Sarpy counties are "younger" than the state and the nation in that the median age is lower. (A composite median is not available for the Metro Area as a whole.)


Median Age (2018-2022)





- Sources: US Census Bureau American Community Survey 5-year estimates.
 - Center for Applied Research and Engagement Systems (CARES), University of Missouri Extension. Retrieved June 2024 via SparkMap (sparkmap.org).





Race & Ethnicity


Race


In looking at race independent of ethnicity (Hispanic or Latino origin), 73.4% of Metro Area residents are White and 8.3% are Black.

BENCHMARK ► The area is more diverse than either state but less diverse than the US as a whole.

DISPARITY ▶ Douglas County is the most racially diverse of the four counties.

Total Population by Race Alone (2020)





Center for Applied Research and Engagement Systems (CARES), University of Missouri Extension. Retrieved June 2024 via SparkMap (sparkmap.org).

Race reflects those who identify with a single race category, regardless of Hispanic origin. People who identify their origin as Hispanic, Latino, or Spanish may be of any race.




Ethnicity

A total of 12.2% of Metro Area residents are Hispanic or Latino.

BENCHMARK ► Higher than the Iowa proportion but well below the US proportion.

DISPARITY ▶ The percentage of Hispanic residents is highest in Douglas County.

Hispanic Population (2020)



- Sources:

 US Census Bureau American Community Survey 5-year estimates.
 - Center for Applied Research and Engagement Systems (CARES), University of Missouri Extension. Retrieved June 2024 via SparkMap (sparkmap.org).
 - Origin can be viewed as the heritage, nationality group, lineage, or country of birth of the person or the person's parents or ancestors before their arrival in the
 United States. People who identify their origin as Hispanic, Latino, or Spanish may be of any race.

Linguistic Isolation

A total of 2.5% of the area population age 5 and older live in a home in which <u>no</u> person age 14 or older is proficient in English (speaking only English or speaking English "very well").

BENCHMARK ▶ Higher than the Iowa figure but lower than the US figure.

DISPARITY ► Highest in Douglas County.

Linguistically Isolated Population (2018-2022)



3.1%	1.2%	0.3%	1.5%	2.5%	2.6%	1.9%	3.9%
Douglas	Sarpy	Cass	Pottawattamie	Metro	NE	IA	US


Sources:

US Census Bureau American Community Survey 5-year estimates

OS Centros burlead American Community Survey 3-year estimates.
 Center for Applied Research and Engagement Systems (CARES), University of Missouri Extension. Retrieved June 2024 via SparkMap (sparkmap.org).

This indicator reports the percentage of the population age 5+ who live in a home in which no person age 14+ speaks only English, or in which no person
age 14+ speak a non-English language and speak English "very well."

Note the following map illustrating linguistic isolation throughout the Metro Area.





SOCIAL DETERMINANTS OF HEALTH

ABOUT SOCIAL DETERMINANTS OF HEALTH

Social determinants of health (SDOH) are the conditions in the environments where people are born, live, learn, work, play, worship, and age that affect a wide range of health, functioning, and quality-of-life outcomes and risks.

Social determinants of health (SDOH) have a major impact on people's health, well-being, and quality of life. Examples of SDOH include:

- Safe housing, transportation, and neighborhoods
- Racism, discrimination, and violence
- Education, job opportunities, and income
- Access to nutritious foods and physical activity opportunities
- Polluted air and water
- Language and literacy skills

SDOH also contribute to wide health disparities and inequities. For example, people who don't have access to grocery stores with healthy foods are less likely to have good nutrition. That raises their risk of health conditions like heart disease, diabetes, and obesity — and even lowers life expectancy relative to people who do have access to healthy foods.

Just promoting healthy choices won't eliminate these and other health disparities. Instead, public health organizations and their partners in sectors like education, transportation, and housing need to take action to improve the conditions in people's environments.

- Healthy People 2030 (https://health.gov/healthypeople)

Poverty

The latest census estimate shows 9.7% of the Metro Area total population living below the federal poverty level.

BENCHMARK Lower than the US prevalence but fails to satisfy the Healthy People 2030 objective.

DISPARITY ► Highest in Douglas and Pottawattamie counties.

Among just children (ages 0 to 17), this percentage in the Metro Area is 11.6% (representing an estimated 25,791 children).

BENCHMARK ► Lower than the US prevalence but fails to satisfy the Healthy People 2030 objective.

DISPARITY ► Highest in Douglas and Pottawattamie counties.


Poverty is considered a key driver of health status because it creates barriers to accessing health services, healthy food, and other necessities that contribute to overall health.



Population in Poverty (Populations Living Below the Poverty Level; 2018-2022)

Healthy People 2030 = 8.0% or Lower

Total PopulationChildren




- Sources:

 US Census Bureau American Community Survey 5-year estimates.
 Center for Applied Research and Engagement Systems (CARES), University of Missouri Extension. Retrieved June 2024 via SparkMap (sparkmap.org).
 - US Department of Health and Human Services. Healthy People 2030. August 2020. http://www.healthypeople.gov


 Poverty is considered a key driver of health status. This indicator is relevant because poverty creates barriers to access including health services, healthy food, and other necessities that contribute to poor health status. Notes:

The following maps highlight concentrations of persons living below the federal poverty level.

Population Below the Poverty Level



Children Below the Poverty Level






Education

Among the Metro Area population age 25 and older, an estimated 7.7% (nearly 45,000 people) do not have a high school education.

BENCHMARK ► Lower than the national percentage.


DISPARITY ► Highest in Douglas and Pottawattamie counties.

Population With No High School Diploma (Population Age 25+ Without a High School Diploma or Equivalent, 2018-2022)



US Census Bureau American Community Survey 5-year estimates.

Center for Applied Research and Engagement Systems (CARES), University of Missouri Extension. Retrieved June 2024 via SparkMap (sparkmap.org). This indicator is relevant because educational attainment is linked to positive health outcomes.






Employment

According to data derived from the US Department of Labor, the 2023 unemployment rate in the Metro Area was 2.5%.

BENCHMARK ► Lower than the Iowa and US unemployment rates.

TREND ► Following significant increases in 2020 (attributed to the COVID-19 pandemic), unemployment has dropped below pre-pandemic levels, and lower than found a decade ago.



Sources: • US Department of Labor, Bureau of Labor Statistics.

Center for Applied Research and Engagement Systems (CARES), University of Missouri Extension. Retrieved June 2024 via SparkMap (sparkmap.org).

Notes: • Percent of non-institutionalized population age 16+ who are unemployed (not seasonally adjusted).



Financial Resilience

A total of 24.3% of Metro Area residents would \underline{not} be able to afford an unexpected \$400 expense without going into debt.

BENCHMARK ► Well below the national prevalence.


TREND ► Marks a statistically significant increase since 2021.

DISPARITY In Douglas County, highest among residents east of 72nd Street. By county, highest among Pottawattamie adults. The prevalence decreases with age and household income level, and is reported more often among women, Hispanic residents, White residents, those of Diverse Races, and adults who identify as LGBTQ+.

Respondents were asked: "Suppose that you have an emergency expense that costs \$400. Based on your current financial situation, would you be able to pay for this expense either with cash, by taking money from your checking or savings account, or by putting it on a credit card that you could pay in full at the next statement?"

NOTE: For indicators derived from the population-based survey administered as part of Do Not Have Cash on Hand to Cover a \$400 Emergency Expense

Metro Area




- Sources:
- 2024 PRC Community Health Survey, PRC, Inc. [Item 53]
- 2023 PRC National Health Survey, PRC, Inc.
- Notes: Asked of all respondents.
 - Includes respondents who say they would not be able to pay for a \$400 emergency expense either with cash, by taking money from their checking or savings
 account, or by putting it on a credit card that they could pay in full at the next statement.



this project, text



Do Not Have Cash on Hand to Cover a \$400 Emergency Expense (Metro Area, 2024)



Notes:

- Sources: 2024 PRC Community Health Survey, PRC, Inc. [Item 53]
 - Asked of all respondents.
 - Includes respondents who say they would not be able to pay for a \$400 emergency expense either with cash, by taking money from their checking or savings
 account, or by putting it on a credit card that they could pay in full at the next statement.

INCOME & RACE/ETHNICITY

INCOME ▶ Income categories used to segment survey data in this report are based on administrative poverty thresholds determined by the US Department of Health & Human Services. These guidelines define poverty status by household income level and number of persons in the household (e.g., the 2023 guidelines place the poverty threshold for a family of four at \$30,000 annual household income or lower). In sample segmentation: "very low income" refers to community members living in a household with defined poverty status; "low income" refers to households with incomes just above the poverty level and earning up to twice (100%-199% of) the poverty threshold; and "mid/high income" refers to those households living on incomes which are twice or more (≥200% of) the federal poverty level.

RACE & ETHNICITY ► In analyzing survey results, mutually exclusive race and ethnicity categories are used. All Hispanic respondents are grouped, regardless of identity with any other race group. Data are also detailed for individuals identifying with a race category, without Hispanic origin. "White" reflects those who identify as White alone, without Hispanic origin. "Diverse Races" includes those who identify as American Indian or Alaska Native, Native Hawaiian/Pacific Islander, or as being of multiple races, without Hispanic origin.




Housing

Housing Insecurity

Most surveyed adults rarely, if ever, worry about the cost of housing.

Frequency of Worry or Stress About Paying Rent or Mortgage in the Past Year (Metro Area, 2024)



Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Item 56]
Notes: • Asked of all respondents.

However, nearly one-third (32.5%) report that they were "sometimes," "usually," or "always" worried or stressed about having enough money to pay their rent or mortgage in the past year.


BENCHMARK ► Well below the national figure.


TREND ► Increasing significantly since 2018.

DISPARITY Highest in Douglas County (especially east of 72nd Street). Reported more often among women, young adults, those in low-income households, Hispanic respondents, White respondents, LGBTQ+ respondents, and people who rent.

"Always/Usually/Sometimes" Worried About Paying Rent/Mortgage in the Past Year

Metro Area






2023 PRC National Health Survey, PRC, Inc.

Notes: • Asked of all respondents.



"Always/Usually/Sometimes" Worried About Paying Rent/Mortgage in the Past Year (Metro Area, 2024)



Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Item 56]
Notes: • Asked of all respondents.

Loss of Utilities


A total of 10.3% of Metro Area residents were without electricity, water, or heat at some point in the past year.

TREND Twice the 2018 prevalence (a statistically significant increase).

DISPARITY Highest in Douglas County (especially the Southeast Omaha area). Reported more often among men, adults living in low-income households, LGBTQ+ adults, and those who rent.

Went Without Electricity, Water, or Heating in the Past Year

Metro Area




Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Item 310]
Notes: • Asked of all respondents.



Went Without Electricity, Water, or Heating in the Past Year (Metro Area, 2024)

Among homeowners 7.6% Among renters 14 0%



• 2024 PRC Community Health Survey, PRC, Inc. [Item 310] Asked of all respondents.

Unhealthy or Unsafe Housing

A total of 13.1% of Metro Area residents report living in unhealthy or unsafe housing conditions during the past year.


BENCHMARK ► Lower than the US prevalence.

TREND ▶ Denotes a statistically significant increase since 2018.


DISPARITY Higher in Douglas County (especially east of 72nd Street); due to sample size, the findings for Pottawattamie County are not significant. The Metro Area prevalence decreases with age and household income level and is reported more often among women, Hispanic adults, Black or African American adults, those who identify as LGBTQ+, and renters.

Unhealthy or Unsafe Housing Conditions in the Past Year

Metro Area




- Sources:
- 2024 PRC Community Health Survey, PRC, Inc. [Item 55]
- 2023 PRC National Health Survey, PRC, Inc. Asked of all respondents.
- - Includes respondents who say they experienced ongoing problems in their current home with water leaks, rodents, insects, mold, or other housing conditions that might make living there unhealthy or unsafe.





Unhealthy or Unsafe Housing Conditions in the Past Year (Metro Area, 2024)






- Sources: 2024 PRC Community Health Survey, PRC, Inc. [Item 55]
 - Asked of all respondents.
 - Includes respondents who say they experienced ongoing problems in their current home with water leaks, rodents, insects, mold, or other housing conditions that might make living there unhealthy or unsafe.

Transportation

While the vast majority of survey respondents report owning their own vehicle for transportation purposes, 12.7% rely on other means of transportation.

This includes a total of 6.5% who have someone else who drives them and 6.2% who rely on other modes like public transportation, walking, etc.

Primary Form of Transportation (Metro Area, 2024)



- Sources: 2024 PRC Community Health Survey, PRC, Inc. [Item 311]



Food Access

Low Food Access


US Department of Agriculture data show that 19.3% of the Metro Area population (representing nearly 153,000 residents) have low food access, meaning that they do not live near a supermarket or large grocery store.

BENCHMARK ► Lower than the US percentage.

DISPARITY ► Highest in Sarpy and Pottawattamie counties.

Population With Low Food Access


(Percent of Population That Is Far From a Supermarket or Large Grocery Store, 2019)



Sources: • US Department of Agriculture, Economic Research Service, USDA - Food Access Research Atlas (FARA).

Center for Applied Research and Engagement Systems (CARES), University of Missouri Extension. Retrieved June 2024 via SparkMap (sparkmap.org).

This indicator reports the percentage of the population with low food access. Low food access is defined as living more than ½ mile from the nearest supermarket, supercenter, or large grocery store. This indicator is relevant because it highlights populations and geographies facing food insecurity.





Low food access is

defined as living more

rural areas) from the nearest supermarket, supercenter, or large

grocery store.
RELATED ISSUE
See also Difficulty
Accessing Fresh Produce

of this report.

than 1/2 mile (in urban areas, or 10 miles in

in the Nutrition, Physical

Activity & Weight section

Surveyed adults were asked: "Now I am going to read two statements that people have made about their food situation. Please tell me whether each statement was "often true," "sometimes true," or "never true" for you in the past 12 months:

I worried about whether our food would run out before we got money to buy more.

The food that we bought just did not last, and we did not have money to get more."

Those answering "often" or "sometimes" true for either statement are considered to be food insecure.

Food Insecurity

Over the past year, 25.6% of Metro Area residents "often" or "sometimes" worried about running out of food.


BENCHMARK ► Well below national findings.

TREND ▶ Increasing significantly to the highest percentage recorded to date.

DISPARITY ► Highest in Douglas County (especially the eastern region).

"Often" or "Sometimes" Worry About Food Running Out Before Having Money to Buy More

Metro Area



Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Item 67]

• 2023 PRC National Health Survey, PRC, Inc.


Notes: • Asked of all respondents.

In fact, 20.6% of adults actually ran out of food in the past year before there were funds to buy more.

BENCHMARK ► Well below the US figure.

DISPARITY ► By county, highest in Douglas and Pottawattamie counties. In Douglas County, much higher east of 72nd Street.

Ran Out of Food in the Past Year



Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Item 98]

2023 PRC National Health Survey, PRC, Inc.

Notes: • Asked of all respondents.




Overall, 27.8% of community residents are determined to be "food insecure," having run out of food in the past year and/or been worried about running out of food.

BENCHMARK ► Lower than the national prevalence.

DISPARITY By county, highest in Douglas and Pottawattamie counties. Especially high in Omaha east of 72nd Street. The percentage decreases with age and household income level and is reported more often among women, Hispanic adults, Black or African American adults, and LGBTQ+ adults.

Food Insecure




Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Item 98]

2023 PRC National Health Survey, PRC, Inc.

Notes: • Asked of all respondents.

Food Insecure (Metro Area, 2024)



Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Item 98]

Asked of all respondents.



Discrimination

Unfair Treatment

Over one in four survey respondents (28.2%) reports that in their daily lives, they feel that they are frequently (a few times per month or more often) treated with less courtesy or respect than other people; another 17.8% report frequently being treated as less intelligent than others.

Fewer respondents (10.1%) report frequently receiving **poorer service** at restaurants and stores, and 8.7% report frequently being treated as a **potential danger** by others. A total of 7.4% of survey respondents have been frequently **threatened or harassed**.

TREND ▶ With the exception of being treated as a potential danger, each of these indicators has increased significantly since 2021.


Perceptions of Unfair Treatment in Day-to-Day Life (Metro Area Trends)

2021 2024

Using a short version of the Everyday Discrimination Scale (EDS), respondents were asked about how frequently they encounter treatment they perceive to be unfair.

In your day-to-day life, how often do the following things happen to you:

- You are treated with less courtesy or respect than other people?
- You receive poorer service than other people at restaurants or stores?
- People act as if they think you are not smart?
- People act as if they are afraid of you?



Sources:

- 2024 PRC Community Health Survey, PRC, Inc. [Items 325-329]
- Notes: Asked of all respondents
 - Percentages represent combined "almost daily," "at least weekly," and "a few times a month" responses.

For those who felt they were treated differently, reasons were most often attributed to *race/ethnicity*, *gender*, *age*, *or height/weight* (although nearly one out of four was unsure of the main reason why).


Further looking at these responses according to these respondents' characteristics, notable differences also appeared for income.

DISPARITY Viewed by **household income level**, note the negative correlation between household income and the tested aspects of unfair treatment, with those in <u>very low income</u> households more likely to report each.



Perceptions of Unfair Treatment in Day-to-Day Life (By Household Income Level; Metro Area, 2024)

Very Low Income Low IncomeMid-High Income



- Sources: 2024 PRC Community Health Survey, PRC, Inc. [Items 325-329]
 - Asked of all respondents.
 - Percentages represent combined "almost daily," "at least weekly," and "a few times a month" responses.

DISPARITY Viewed by race/ethnicity, Black or African American adults reported the highest percentage of being threatened or harassed. For all others, responses were notably higher for Hispanic respondents, Black or African American respondents, and respondents of Diverse Races.

Perceptions of Unfair Treatment in Day-to-Day Life (By Race/Ethnicity; Metro Area, 2024)

HispanicWhiteBlackAsianDiverse Races



- Sources: 2024 PRC Community Health Survey, PRC, Inc. [Items 325-329]


 - Percentages represent combined "almost daily," "at least weekly," and "a few times a month" responses.



Community as a Welcoming Place for All Races/Ethnicities

Most Metro Area adults agree that the community is a welcoming place for people of all races and ethnicities, with over 70% giving "agree" or "strongly agree" responses.

Level of Agreement About the Community as a Welcoming Place for People of All Races and Ethnicities (Metro Area, 2024)



• 2024 PRC Community Health Survey, PRC, Inc. [Item 306]


Asked of all respondents.

However, 11.0% of residents do not agree that the community is welcoming to all people.

DISPARITY Disagreement is lowest in Sarpy County. By race/ethnicity, highest among Black or African American adults and those of Diverse Races. Also higher among women, young adults, and those who identify as LGBTQ+.

Disagree That the Community is a Welcoming Place for All Races/Ethnicities

Metro Area






Notes:
• Asked of all respondents

Percentages include "disagree" and "strongly disagree" responses.



Disagree That the Community is a Welcoming Place for All Races/Ethnicities (Metro Area, 2024)




- Asked of all respondents.
- Percentages include "disagree" and "strongly disagree" responses.

Treatment Based on Race/Ethnicity in Health Care Settings

Over one-third (35.0%) of survey respondents feel they were treated "better" than people of other races or ethnicities during their recent health care experiences; most (60.9%) felt they were treated "the same."

Treated in Health Care Settings Over the Past Year in Comparison With People of Other Races/Ethnicities (Metro Area, 2024)



- Sources: 2024 PRC Community Health Survey, PRC, Inc. [Item 313]
 - Asked of all respondents.
 - As compared to the experiences of people of other races or ethnicities.



"In the past 12 months, in general, do you feel your

health care experiences

were "better," "the same," or "worse" than those of other races or ethnicities?"


On the other hand, 4.1% of residents perceive their treatment as being "worse" than people of other races during recent health care experiences.

BENCHMARK ► Lower than the national prevalence.

DISPARITY ► Reported most often in Douglas County (primarily from the county's northeast region). Reported most often among Black or African American respondents.

Recent Health Care Experiences Were Worse Due to Race





Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Item 307]

2023 PRC National Health Survey, PRC, Inc.

Notes: • Asked of all respondents.

When respondents were asked how their "worse" treatment has affected the way they try to get their health care, the largest share (60.0%) indicated that they **have not changed** their health-seeking behaviors as a result, and 6.9% were unsure. However, 6.7% report **avoiding care** or **putting off** their care as long as possible when in need of health care services.



Adverse Childhood Experiences (ACEs)

ABOUT ACEs

Adverse Childhood Experiences (ACEs) are stressful or traumatic events, including abuse and neglect. They are a significant risk factor for substance use disorders and can impact prevention efforts. ACEs include:

- Physical abuse or neglect
- Emotional abuse or neglect
- Sexual abuse
- Intimate partner violence
- Household substance misuse
- Household mental illness
- Parental separation/divorce
- Incarcerated household member

A series of 11 survey questions was used to identify adults' experiences of adverse childhood events prior to the age of 18 years. These 11 questions align with eight ACEs categories, as outlined in the following table.

Adverse Childhood Experiences (ACEs)


CATEGORY	QUESTION			
HOUSEHOLD MENTAL ILLNESS	Before you were 18 years of age, did you live with anyone who was depressed, mentally ill, or suicidal?			
	Before you were 18 years of age, did you live with anyone who was a problem drinker or alcoholic?			
HOUSEHOLD SUBSTANCE USE	Before you were 18 years of age, did you live with anyone who used illegal street drugs or who abused prescription medications?			
INCARCERATED HOUSEHOLD MEMBER	Before you were 18 years of age, did you live with anyone who served time or was sentenced to serve time in a prison, jail, or other correctional facility?			
PARENTAL SEPARATION OR DIVORCE	Before you were 18 years of age, were your parents separated or divorced?			
INTIMATE PARTNER VIOLENCE	Before age 18, how often did your parents or adults in your home slap, hit, kick, punch or beat each other up?			
PHYSICAL ABUSE	Before age 18, how often did a parent or adult in your home hit, beat, kick, or physically hurt you in any way? Do not include spanking.			
EMOTIONAL ABUSE	Before age 18, how often did a parent or adult in your home swear at you, insult you, or put you down?			
	Before you were 18 years of age, how often did an adult or anyone at least 5 years older than you touch you sexually?			
SEXUAL ABUSE	Before you were 18 years of age, how often did an adult or anyone at least 5 years older than you try to make you touch them sexually?			
	Before you were 18 years of age, how often did an adult or anyone at least 5 years older than you force you to have sex?			

- Sources: 2024 PRC Community Health Survey, PRC, Inc.
- Reflects the total sample of respondents

By category, ACEs were most prevalent in the Metro Area for emotional abuse (experienced by 46.9% of respondents during childhood), followed by household substance use (34.1%) and parental separation or divorce (31.2%).



Adverse Childhood Experiences (ACEs) (Metro Area, 2024)



- Sources: 2024 PRC Community Health Survey, PRC, Inc. [Items 344-351]
- Reflects the total sample of respondents
 - . ACEs are stressful or traumatic events, including abuse and neglect. They are a significant risk factor for substance abuse disorders and can impact prevention efforts.

High ACE Scores

The impact of ACEs on future health and well-being are cumulative. PRC looks at these compounding issues by scoring the ACE series — survey respondents receive one "point" for each of the eight ACEs categories containing an affirmative response; a score of four or higher is determined to be a "high" ACE score.


In all, 24.4% of Metro Area residents reported four or more of the adverse childhood experiences tested (a high ACE score).

TREND ▶ Increasing significantly since 2018 (the series was not addressed in 2021).

DISPARITY Highest in Northeast Omaha and Pottawattamie County. The prevalence decreases with age and household income level and is also reported more often among women, Hispanic adults, Black or African American adults, those of Diverse Races, and people who identify as LGBTQ+.

Prevalence of High ACE Scores (Four or More ACEs)

Metro Area






2023 PRC National Health Survey, PRC, Inc.

Asked of all respondents.

Adults who report four or more ACEs is categorized as having a high ACE score.



Prevalence of High ACE Scores (Four or More ACEs) (Metro Area, 2024)



Sources: 2024 PRC Community Health Survey, PRC, Inc. [Item 352]

Asked of all respondents.


Adults who report four or more ACEs is categorized as having a high ACE score.

Relationship of ACEs with Other Health Issues

As a person's ACE score increases, so does their risk for disease, social issues, and emotional problems.

Note the following strong correlations of various health indicators in the Metro Area, comparing those reporting no ACEs with those with low (1-3) and high (4+) ACE risk.

Relationship of ACEs With Other Health Issues (By ACE Risk Classification; Metro Area, 2024)





Asked of all respondents.

Adults with at least one ACE are categorized as having a low score (1 to 3 ACEs) or a high score (4+ ACEs).



Key Informant Input: Social Determinants of Health

The greatest share of key informants taking part in an online survey characterized Social Determinants of Health as a "major problem" in the community.

Perceptions of Social Determinants of Health as a Problem in the Community (Among Key Informants; Metro Area, 2024)



Among those rating this issue as a "major problem," reasons related to the following:

Housing

· Asked of all respondents

Housing is a pretty big issue in the community, especially for low-income families. Education is also a problem in the urban areas of the community. – Community Leader

There is a huge shortage of affordable housing in the metro area, incomes are stagnant, and everything is so expensive. – Social Services Provider

Housing costs (rent, mortgage) are astronomical for average families. The cost of living compared to wages is high. Access to health care, but more so, adequate health coverage or available money for health care. Safe environments for low-income families, especially those of color and refugees. Education is not equal across the city. – Health Provider

There is not enough affordable housing in Omaha. Our most vulnerable populations receive on average \$1,000 a month from social security or disability. That is not enough to cover rent, utilities, and food. Homelessness has increased and resources are available; however, the process of accessing resources is too complicated and overwhelming for the population they are meant to serve. – Health Provider

People living in our community have difficulty accessing affordable housing and suffer from health disparities. – Physician

Lack of housing, transportation and food security are a major problem facing my patient population. – Health Provider

Lack of housing causes increased truancy in school/collage attendance, low or lack of income cause an increase in mental and medical issues, environmental concerns health issues such as lead, poor water quality. Discrimination causes low self-esteem, poor choices in basic needs (Maslow hierarchy). No or little knowledge of medical follow-up. Poor lifestyle choices such as smoking, alcohol, drugs, which lead to medical conditions. – Health Provider

We have a 0.9% homeowner vacancy rate and a 1.4% rental vacancy rate. There are almost literally no places to live. — Community Leader

The impact of housing, poverty, and access to healthy food has geocentric limitations. - Physician

Lack of affordable housing, increase in food insecurity, truancy in the schools. — Social Services Provider

The lack of affordable housing and/or acceptance and availability of housing vouchers is out of control. LGBTQ+
unhoused individuals have NO OPTIONS for housing, unless they are 18-26, because most programs are
always full, religiously affiliated, or focused on emergent intervention for chronically unhoused. There is a need
for primarily transitional and supportive housing, and rapid rehousing rather than interventions in the most
emergent interventions. The state legislature continues to invest in corporate tax incentives without thresholds for
pay, which allows businesses to pay poorly, not provide benefits, and still receive tax incentives while their
employees are forced into state aid program — essentially double-dipping. There is a need for alcohol, tobacco,
and gambling density laws or ordinances in qualified census tract or low-income areas, which are routinely

targeted by commercial industries marketing unhealthy products to historically marginalized communities. -



Community Leader

According to a recent study, we are short nearly 100,000 units of affordable housing in our community. For people living with untreated serious and persistent mental illness, the likelihood of homelessness is significant and there aren't enough Permanent Supportive Housing units available to meet the need. Poverty, lack of safe housing, and untreated mental illness drives the people impacted by these social problems to become high utilizers of emergency room and other expensive (often unreimbursed) medical care. And, of course, due to our ingrained systemic inequities, the impact is especially toxic for our BIPOC communities. – Social Services Provider

We have a housing shortage. We have increased the cost of living without much of an increase in wages. Hate and bigotry every day play a role in people not being able to get even most basic conditions for human survival met. Red Lining is still happening in our community. The bias that people who are experiencing homelessness face is a very real problem, yet most people understand housing instability. — Community Leader

Housing and food are basic needs. If they are unmet, a person cannot achieve their highest level of health and wellbeing. Lack of transportation and education are barriers to gainful employment and livable wages. – Health Provider

Many of the patients I see hospitalized have housing instability, unemployment, and food instability. Housing specifically needs to be addressed, and we need to have more low-income options that are transitions between the shelters to regular housing. – Physician

Income/Poverty

They ultimately drive much of the chronic illness and outcomes for the working poor. Housing is a major issue. – Physician

Growing income gap. Wages not keeping up with cost of living. - Community Leader

More people are having difficulties paying their rent, utilities, and other necessities. More people are needing access to food pantries as they do not have access to food sources. An increase in people are living out of their vehicles or surfing from one couch to another. More people are saying they are unable to work, so more people are trying to apply for disability and not being approved. — Health Provider

Poverty, homelessness, lack of affordable housing, family history, education. - Social Services Provider

Wealth disparity in Omaha affects housing, food, and access to care. - Physician

Income equality causes significant challenges with access to housing, education, legal services, etc. - Physician

Too many people struggle to afford housing, food, health and other insurance, transportation, education, health care, etc. – Physician

Poverty. - Physician

Income disparities. - Social Services Provider

Across our country there are more and more working families who are struggling with finances, housing, food insecurity, etc. We have people living on the streets, people standing on corners panhandling across our community, sometimes with their children. There is a lack of affordable housing, so people are going without food, medication etc. in order to make their rent/mortgage payment. The low paying jobs don't have flexibility so parents lose their jobs when they can't find affordable childcare or when they need to stay home with a sick child. Luxury apartments are being built all over town but limited affordable housing is being created. Our shelters are full. – Social Services Provider

Social determinants are foundational to health. Health is bigger than health care and if you do not have the basics such as living wage, transportation, housing, etc. it is near impossible to achieve or sustain good health. – Business Leader

Poverty levels are higher in Latino communities. - Health Provider

We have pockets of poor people with lack of basic living resources. - Health Provider

There are many people living at the poverty level, many homeless, many with inadequate education, etc. – Health Provider

People have to prioritize basic needs, and when juggling on a tight household income, they aren't able to equally support all. – Social Services Provider

My community has the highest poverty rates, lacks adequate and affordable housing, high lead levels, and is subjected to racism and discrimination. – Community Leader

Access to Care/Services

Because of the lack of mental health services, timely services, housing is an issue, the homeless population, substance use disorders, lack of funding available, not having access to medical insurance, Medicaid eligibility limitations. – Social Services Provider

Hard to get into skilled care or nursing home. Lots of homeless with poor housing options. - Physician



Many of the above problems are direct results and further exacerbated by social determinants of health. Access to care for those without insurance remains difficult. Access to culturally respectful and linguistically appropriate care is limited, especially beyond primary care. Home health, SNF and nursing home care is excessively limited for patients with Medicaid. Medicaid transportation is often not reliable. Limited access to fresh food-access there is often limited to those that speak English and have social security number. Low-income housing is limited and in poor condition. – Physician

It is impossible to go to doctor's appointment if you don't have insurance, or if you have unstable housing, food insecurity, no access to reliable internet services, or if you are working 3 jobs while also going to school and taking care of children or dependent adults in the family. This is particularly impacting non-white people, immigrants, refugees in the community, as well as people in recovery or previously justice involved. – Health Provider

SDOH influence access to care and health equity. Homelessness increases the likelihood of emergency department use. – Advanced Practice Provider

It affects access, follow up, increases recidivism. - Physician

Access to care, health care deserts. - Physician

Racism

Inequality, injustice, poverty, and racism. - Business Leader

Redlining and racism have created wealth inequality and contributed to disinvestment in communities of color and those most impacted by these indicators of health. – Community Leader

Systemic racism and lack of accountability will only continue to perpetuate this issue. – Community Leader Historical and systemic racism. Socioeconomic disparities. Limited access to health care — the State of Nebraska DHHS does not advertise its program and services that would increase access to health care. Environmental injustice. Cultural and language barriers. Food insecurity and food deserts. Educational disparities. – Community Leader

This system doesn't work for all people who live here. Systemic racism and sexism impact how people can afford safe and quality housing, the communities/neighborhoods where they live, and their overall health. – Business Leader

The main social determinant of health is racial inequality, rooted in racial discrimination. Unfortunately, this is what determines if you are healthy or not in this country. Racial inequality impacts a black or brown child's access to quality opportunities, such as quality food, quality housing, quality jobs, quality health care and education. Racial inequity I believe is the contributor and development of social determinants of health. It has led to education barriers, that lead to income barriers, to lead to housing barriers, to living in toxic environments that impacting the health and wellbeing of a child or person. — Social Services Provider

Homelessness

At the core of numerous, if not all, health issues lie the social determinants of health. For individuals experiencing homelessness, the likelihood of having additional severe underlying conditions that compound their current health challenges is alarmingly high. Consider this: residing in an area with limited access to nutritious food while grappling with a heart condition poses a significant dilemma. How can one be expected to uphold their heart health without access to a diet conducive to it? Furthermore, the glaring gap in affordable housing exacerbates these issues. The lack of stable housing not only perpetuates the cycle of homelessness but also amplifies the risk of deteriorating health conditions and complicates access to essential health care services. This confluence of social determinants creates a formidable barrier to achieving and maintaining optimal health for vulnerable populations. – Health Provider

Over 120 homeless camps in the community, lack of affordable and sustainable housing exist in the community. Landlords and management companies are charging extra fees to live in their properties. Limited resources and long waits for housing. HUD guidelines that identify homeless or near homeless with housing programs and financial assistance programs dictate who gets assistance and how long you must be homeless to get the assistance. – Health Provider

There are too many homeless people in this country for as wealthy as it is, but our government spends money on foreign countries or wastes it wherever Congress decides, and yet they don't want to take care of their own. There is a lack of wanting to get and keep people healthy by helping out in all of these areas. Our government, federal, state and local have no intentions of spending what is our money to help people become, healthy, educated and housed. In my opinion, we should be loading money into programs and assistance on the front end instead of continuing to build hospitals and prisons on the back end. But, Corporate America, and big pharma only make money on the sick and dying. – Health Provider

Impact on Quality of Life

So many can't meet basic needs which are directly related to overall health. - Health Provider



Health care does not occur in a silo. Any chronic disease or illness and its management occur in the context of SDOH and the patient's life and community. Douglas County and rural Nebraska present their own unique SDOH for community members. Fear of stigma related to substance overdose leads to under reporting of substance related deaths in rural Nebraska; racial injustices and racial health disparities continue to exist in urban Omaha. – Physician

Everything comes together to impact a person's health. When the basics aren't available and equitable, people struggle. – Social Services Provider

80% of health is dictated by social determinants of health, can't achieve health without stable needs being met. – Health Provider

Incidence/Prevalence

Mental illness and homelessness are on the uprise. – Health Provider Significant inequities exist throughout the region. – Business Leader

Prevention/Screenings

Lack of screening, lack of loop closure mechanism for when a referral is made, checkbox mentality of health systems as opposed to addressing the root problem of financial insecurity or unsafe environment. – Health Provider

Lack of consistency in screening for and securing supports for different SDOH across the community. – Health Provider

Low Education Levels

Without education, low-income people find it difficult to get affordable housing and cannot afford to eat healthier and make better choices. Food high in calories makes people become overweight and for people of color it becomes more difficult, and many times is discriminated. – Social Services Provider

Diagnosis/Treatment

This is a complex issue that most health care provides do not give much thought to. Our federally qualified health centers do a good job in their awareness of SDOHs. – Physician

Transportation

Poor public transit (need more routes in some parts of town, tend to concentrate in areas with better overall transportation resources), need higher minimum wage in NE, poor distribution of health care resources (e.g., no medical specialist and little mental health resources on the north side). – Physician

Follow Up/Support

Our society is currently a mess, with reduction in social supports, funding, and a growing wealth gap. – Business Leader





HEALTH STATUS


OVERALL HEALTH STATUS

Most Metro Area residents rate their overall health favorably (responding "excellent," "very good," or "good").

good," or "good").

The initial inquiry of the PRC Community Health Survey asked: "Would you say that in general your health is excellent, very good, good, fair, or poor?"





Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Item 4]
Notes: • Asked of all respondents.


However, 16.3% of Metro Area adults believe that their overall health is "fair" or "poor."

TREND ► Increasing significantly since 2011.

DISPARITY Highest in Southeast Omaha and Pottawattamie County. The prevalence increases with age, decreases with household income level, and is reported more often among men, and people who identify as LGBTQ+.

Experience "Fair" or "Poor" Overall Health

Metro Area




Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Item 4]

- Behavioral Risk Factor Surveillance System Survey Data. Atlanta, Georgia. United States Department of Health and Human Services, Centers for Disease Control and Prevention (CDC): 2022 Nebraska and Iowa data.
- 2023 PRC National Health Survey, PRC, Inc.

Notes: • Asked of all respondents.



Experience "Fair" or "Poor" Overall Health (Metro Area, 2024)



Sources:

• 2024 PRC Community Health Survey, PRC, Inc. [Item 4]

• Asked of all respondents.



MENTAL HEALTH

ABOUT MENTAL HEALTH & MENTAL DISORDERS

About half of all people in the United States will be diagnosed with a mental disorder at some point in their lifetime. ...Mental disorders affect people of all ages and racial/ethnic groups, but some populations are disproportionately affected. And estimates suggest that only half of all people with mental disorders get the treatment they need.

In addition, mental health and physical health are closely connected. Mental disorders like depression and anxiety can affect people's ability to take part in healthy behaviors. Similarly, physical health problems can make it harder for people to get treatment for mental disorders. Increasing screening for mental disorders can help people get the treatment they need.


- Healthy People 2030 (https://health.gov/healthypeople)

Mental Health Status

Most Metro Area adults rate their overall mental health favorably ("excellent," "very good," or "good").

"Now thinking about your mental health, which includes stress, depression, and problems with emotions, would you say that, in general, your mental health is excellent, very good, good, fair, or poor?"

Self-Reported Mental Health Status (Metro Area, 2024)



Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Item 77]
Notes: • Asked of all respondents.

However, 22.7% believe that their overall mental health is "fair" or "poor."


TREND Increasing significantly from earlier survey administrations.

DISPARITY ► Highest in Southeast Omaha and Pottawattamie County.



Experience "Fair" or "Poor" Mental Health

Metro Area



Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Item 77]
• 2023 PRC National Health Survey, PRC, Inc.

Asked of all respondents.

Depression & Anxiety

Diagnosed Depression

A total of 32.3% of Metro Area adults have been diagnosed by a physician or other health professional as having a depressive disorder (such as depression, major depression, dysthymia, or minor depression).


BENCHMARK ► Well above the Nebraska and Iowa percentages.

TREND ▶ Denotes a statistically significant increase since first asked in 2018.

DISPARITY ► Lowest among Sarpy County respondents.

Have Been Diagnosed With a Depressive Disorder

Metro Area



Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Item 80]

Behavioral Risk Factor Surveillance System Survey Data. Atlanta, Georgia. United States Department of Health and Human Services, Centers for Disease Control and Prevention (CDC): 2022 Nebraska and Iowa data.

2023 PRC National Health Survey, PRC, Inc.

Asked of all respondents.

· Depressive disorders include depression, major depression, dysthymia, or minor depression



Symptoms of Chronic Depression

A total of 41.8% of Metro Area adults have had two or more years in their lives when they felt depressed or sad on most days, although they may have felt okay sometimes (symptoms of chronic depression).


BENCHMARK ► Lower than the national percentage.

TREND ► Increasing significantly since 2011.

DISPARITY Highest in Douglas County (especially the eastern region). Reported more often among women, young adults, those living in low-income households, Hispanic residents, Black or African American residents, and LGBTQ+ adults.

Have Experienced Symptoms of Chronic Depression


Metro Area



- Sources: 2024 PRC Community Health Survey, PRC, Inc. [Item 78]
 - 2023 PRC National Health Survey, PRC, Inc.
- lotes:

 Asked of all respondents.
 - . Chronic depression includes periods of two or more years during which the respondent felt depressed or sad on most days, even if (s)he felt okay sometimes.

Have Experienced Symptoms of Chronic Depression (Metro Area, 2024)



Sources:

- 2024 PRC Community Health Survey, PRC, Inc. [Item 78]
- Asked of all respondents.
 - Chronic depression includes periods of two or more years during which the respondent felt depressed or sad on most days, even if (s)he felt okay sometimes.



The Patient Health Questionnaire-4 (PHQ-4) was developed in order to address anxiety and depression, two of the most prevalent illnesses among the general population and often comorbid in nature.

The PHQ-4 is a four-item questionnaire allowing for ultra-brief and accurate measurement of core symptoms/signs of depression and anxiety. An elevated PHQ-4 score is not diagnostic but is an indicator for further inquiry to establish the presence or absence of a clinical disorder warranting treatment.

Respondents were asked:

During the past two weeks, how often have you been bothered by the following problems:

- Feeling Nervous, Anxious, or On Edge
- Not Being Able to Stop or Control Worrying
- Feeling Down, Depressed, or Hopeless
- Feeling Little Interest or Pleasure in Doing

Responses were scored according to how frequently each was experienced in the previous two weeks (nearly every day, more than half the days, several days, or not at all).

Current Anxiety & Depression


At the time of the survey, 20.0% of Metro Area respondents reported experiencing feelings that signal moderate-to-severe anxiety and/or depression (reflecting a PHQ-4 score of 6 or higher).

TREND ▶ Denotes a statistically significant increase since 2021.

DISPARITY ► Highest in Douglas County (especially Northeast Omaha) and Pottawattamie County. The prevalence decreases with age and household income and is reported more often among women and LGBTQ+ respondents.

Moderate to Severe Anxiety/Depression


Metro Area



- Sources: 2024 PRC Community Health Survey, PRC, Inc. [Item 357]
- lotes:

 Asked of all respondents.
 - Reflects a PHQ-4 score of 6 or higher.

Moderate to Severe Anxiety/Depression (Metro Area, 2024)




Notes:


- 2024 PRC Community Health Survey, PRC, Inc. [Item 357]
- s: Asked of all respondents.
 - Reflects a PHQ-4 score of 6 or higher



Stress

A majority of surveyed adults characterize most days as no more than "moderately" stressful.





Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Item 79]
Notes: • Asked of all respondents.

In contrast, 18.1% of Metro Area adults feel that most days for them are "very" or "extremely" stressful.


BENCHMARK ► Lower than the national percentage.


TREND ► Marks a statistically significant increase from previous survey administrations.

DISPARITY ► Most often reported in Pottawattamie County. Higher among women, young adults, those in low-income households, and those who identify as LGBTQ+.

Perceive Most Days As "Extremely" or "Very" Stressful

Metro Area






2023 PRC National Health Survey, PRC, Inc.

Notes: • Asked of all respondents.



Perceive Most Days as "Extremely" or "Very" Stressful (Metro Area, 2024)



• 2024 PRC Community Health Survey, PRC, Inc. [Item 79] Sources: Asked of all respondents.


Social Support

Three in four Metro Area adults (74.6%) report having someone to turn to "all" or "most" of the time, if they needed or wanted help.

TREND ▶ Decreasing significantly since 2018.

DISPARITY ► Especially low in Douglas County (especially Southeast Omaha), young adults, those in low-income households, Hispanic residents, those of Diverse Races, and LGBTQ+ adults.

Have Someone to Turn to for Help All/Most of the Time




Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Item 320]

Notes:

 Asked of all respondents.



Have Someone to Turn to for Help All/Most of the Time (Metro Area, 2024)



Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Item 320]
Notes: • Asked of all respondents.

Suicide

The Metro Area experiences 14.1 suicides per 100,000 population (2018-2020 annual average age-adjusted rate).


BENCHMARK ▶ Lower than the lowa suicide rate.

TREND The suicide rate has increased over the past decade, echoing state and national trends.

DISPARITY ► Highest in Pottawattamie County and among White residents in the Metro Area.

Suicide: Age-Adjusted Mortality (2018-2020 Annual Average Deaths per 100,000 Population)

Healthy People 2030 = 12.8 or Lower




- Sources: CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics. Data extracted June 2024.
 - US Department of Health and Human Services. Healthy People 2030. August 2020. http://www.healthypeople.gov
 - Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10).
 - Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population.

Refer to "Leading Causes of Death" for an explanation of the use of age-adjusting for these rates.



Suicide: Age-Adjusted Mortality by Race (2018-2020 Annual Average Deaths per 100,000 Population)

Healthy People 2030 = 12.8 or Lower



Sources: • CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics. Data extracted June 2024.

- US Department of Health and Human Services. Healthy People 2030. August 2020. http://www.healthypeople.gov
 Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10).
 Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population.

Suicide: Age-Adjusted Mortality Trends (Annual Average Deaths per 100,000 Population)

Healthy People 2030 = 12.8 or Lower



	2011-2013	2012-2014	2013-2015	2014-2016	2015-2017	2016-2018	2017-2019	2018-2020
Metro Area	10.1	11.0	11.4	12.0	12.0	11.9	13.7	14.1
─ NE	11.5	12.5	12.2	12.7	13.2	13.7	14.7	14.8
——IA	13.7	13.3	13.7	13.8	14.5	15.0	15.7	16.7
US	12.5	12.8	13.1	13.4	13.6	13.9	14.0	13.9

Sources: • CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics. Data extracted June 2024.

US Department of Health and Human Services. Healthy People 2030. August 2020. http://www.healthypeople.gov
 Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10).

• Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population.



Notes:

Notes:

Mental Health Treatment


Mental Health Providers

In 2020, the Metro Area had 214.5 mental health providers (including psychiatrists, psychologists, clinical social workers, and counselors who specialize in mental health care) for every 100,000 population.

BENCHMARK ► Well above the Iowa proportion.

DISPARITY ► Lowest in Sarpy and Cass counties.

Access to Mental Health Providers (Number of Mental Health Providers per 100,000 Population, 2020)



- Sources:
 Centers for Medicare and Medicaid Services, National Plan and Provider Enumeration System (NPPES).
- Center for Applied Research and Engagement Systems (CARES), University of Missouri Extension. Retrieved June 2024 via SparkMap (sparkmap.org).
 - This indicator reports the rate of the county population to the number of mental health providers including psychiatrists, psychologists, clinical social workers, and counselors that specialize in mental health care

Currently Receiving Treatment

A total of 28.6% are currently taking medication or otherwise receiving treatment from a doctor or other health professional for some type of mental health condition or emotional problem.

BENCHMARK ► Higher than the national prevalence.

TREND ► Increasing significantly since 2018.

DISPARITY ► Lowest in western Douglas County.



Note that this indicator only reflects providers

practicing in the Metro Area and residents in the

Metro Area; it does not account for the potential


demand for services from outside the area, nor the potential availability of providers in surrounding

areas.

Currently Receiving Mental Health Treatment

Among respondents ever diagnosed with a depressive disorder, 65.7% are currently receiving treatment.

Metro Area



Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Item 81]

2023 PRC National Health Survey, PRC, Inc.

Notes:

 Asked of all respondents.

"Treatment" can include taking medications for mental health.

Difficulty Accessing Mental Health Services


A total of 11.8% of Metro Area adults report a time in the past year when they needed mental health services but were not able to get them.

TREND ► Increasing significantly since 2018.

DISPARITY Highest in Douglas County (especially the northeast region). Reported more often among women, young adults, those in low-income households, and LGBTQ+ respondents.

Unable to Get Mental Health Services When Needed in the Past Year

Metro Area




Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Item 82]

2023 PRC National Health Survey, PRC, Inc.

Notes: • Asked of all respondents.



Unable to Get Mental Health Services When Needed in the Past Year (Metro Area, 2024)




Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Item 82]
Notes: • Asked of all respondents.

Key Informant Input: Mental Health

Nearly all key informants taking part in an online survey characterized *Mental Health* as a "major problem" in the community.

Perceptions of Mental Health as a Problem in the Community (Metro Area, 2024)



Among those rating this issue as a "major problem," reasons related to the following:

Access to Care/Services

Access to services and also lack of education on self-care and prevention. — Social Services Provider

University based settings tend to support clinicians to perform public-facing activities more than patient-care activities, so the waiting lists are very long when only very few psychiatrists and therapists are doing the work. Most of these institutions only see patients M-F from 8 to 5, making it hard for working adults and those with caregiving responsibilities to see anyone. Private practice groups do not typically accept Medicare, so it is very difficult for Medicare beneficiaries to receive care. Medicare's reimbursement for MH therapists and MFTs who aren't psychologists and social workers are so low many opt out of it. People with ID/DD and older adults are often unaware of or unable to access culturally sensitive and competent services. Transportation and telehealth technology are not always available or accessible. Lack of long-term inpatient hospitalization options. LTC is not willing to take people with SMI, and certainly not registered sex offenders with SMI. — Health Provider

Access, availability, affordability, space for inpatients, and specialty care. - Health Provider

Access to resources and appropriate therapy treatments. No system exists for efficiently and reliably communicating with primary care providers and the public about how to connect patients to therapists in the community. – Physician

Timely access to care. - Health Provider



Access to outpatient care. Lack of inpatient beds for all patients, but especially for pregnant patients. – Physician Lack of resources to respond to rising needs. – Health Provider

Access, understanding their needs and meeting basic needs so they can work on themselves. - Health Provider

The death of therapeutic options for those most in need is a glaring concern, leaving many individuals without essential treatment options. Moreover, access to medication presents a formidable obstacle, particularly for individuals experiencing street homelessness, exacerbating an already dire situation. – Health Provider

Access to care and services, cost, and insurance, or lack thereof. - Health Provider

People who care about their wellbeing and having the ability to access services, especially crisis services. There are often long waits for assessments and therapy/psychiatric appointments for all persons. Compassionate people care for people. Trauma training for everyone is lacking in consistency with information. Not enough housing for people to maintain or access with several mental health diagnoses. – Health Provider

There are not close to enough resources, or accessibility is limited. Transportation to even get to mental health appointments can be a challenge as well. Also, the stigma that is associated with mental health is still very real. – Community Leader

The ability to access services in a timely manner pre-crisis is severely limited, with sometimes months of planning or scheduling required. The constellation of providers accepting insurance, self-pay, managed care, or care networks is cumbersome and challenging to navigate for even an informed person. Access to prescribing mental health professionals is severely limited. – Community Leader

Access to care and counseling, especially for kids. - Health Provider

Access to psychiatric services and to evidence-based psychotherapies. – Physician

Providers accepting new patients. - Social Services Provider

Timely appointments, drug costs and escalation due to response in crisis from law enforcement. – Community Leader

 $Immediate\ access\ to\ mental\ health\ services,\ long\ term\ care,\ crisis\ intervention,\ etc.\ -\ Social\ Services\ Provider$

Depression, anxiety, and lack of resources. - Business Leader

Receiving timely services. - Social Services Provider

Access to care, lack of providers, lack of integration with physical health, lack of assessment of social and non-clinical issues that impact mental health. Poor transitions of care. – Health Provider

People cannot access mental health resources in a timely enough manner to make a difference. Whether from being understaffed, or just lack of concern for mental health resources, it just takes too long for anyone to get help and no doubt, many do not because of it. — Health Provider

Lack of access and the number of people with mental health issues that are homeless. - Community Leader

Access to psychiatry and behavioral health services. - Health Provider

Access to mental health professionals. - Business Leader

Access to therapeutic services, times 1,000. Rising incidences of youth suicide and mental illness. Isolation and coping skills post-pandemic. – Business Leader

I think that access to resources is the biggest challenge. The issues regarding access to resources span from not having facilities and professionals who are able to treat mental health issues to individuals who are unable to afford or get to the facilities. It is also incredibly common for us to treat mental health issues by arresting and incarcerating individuals rather than addressing their needs. Our jail is full of people who need help and is not equipped to give them the help they need. – Community Leader

Finding treatments and appropriate placements if needed. - Social Services Provider

Wait times to get an appointment with a professional, lack of resources to handle larger issues. – Social Services Provider

Getting timely appointments. Good inpatient care for crisis. - Physician

The biggest challenge is that there are not enough mental health resources or professionals in our community. – Business Leader

Access to psychiatrists for medication management. – Social Services Provider

Access to services when needed. Continued engagement with services once started. Motivation/ belief that engaging in services will help them long-term. Ability/ understanding how to connect with resources available to them. So many folks have some form of access but fall away from or never access services for a variety of reasons. – Health Provider

Access to mental health services is limited, particularly in the outpatient setting prior to a crisis situation. – Health Provider

Access to care. - Physician

Access to services. Often individuals have to wait months to get into services for mental health care. – Health Provider

People in acute crisis wait for days for beds due to facilities being overloaded. Those who need outpatient care wait months for appointments. – Health Provider



Access to care. - Health Provider

Access to timely care and reimbursement for services. - Health Provider

Their access to services. There are not enough providers and resources to handle the need for services. – Business Leader

Lack of community resources. - Social Services Provider

We have a lack of resources available to meet our communities' mental health crisis. There aren't enough inpatient beds, and getting in to see a therapist or psychiatrist takes months. – Social Services Provider

Not enough acute inpatient beds, and once in, the stay is far too short. Not enough engagement in the support system in discharge planning, if any planning takes place at all. Too many times they are just discharged to a homeless shelter, with a week of medications. – Health Provider

Mental challenges are becoming more normalized, and the stigma is starting to lower which increases the interest in services. However, there is a lack of workforce in this space to meet the increasing demand. In addition to new patients seeking mental health services, the gap in services is exacerbating those with chronic, pre-existing diagnosis and needs. – Business Leader

Access. - Physician

Accessing mental health services, including behavioral health and psychiatry. - Physician

Many people have difficulty finding an outpatient mental health provider to accept their payer source, or it is taking weeks to months to schedule an appointment for outpatient follow-up. – Health Provider

Limited access to mental health providers due to poor reimbursement for mental health services and burnout in mental health providers as a consequence of the pandemic. – Health Provider

My perception is that there are a lot of folks with brain health issues that are not seeking care, or they are people that do not have access to care due to age, employment and benefits status, or income. – Community Leader

Lack of access to adequate care. - Physician

Access to acute and long-term care, access to transitional cares. - Physician

Limited access. Long wait for psychiatry appointments with many offices not accepting new patients. Inadequate community support resources. – Physician

Lack of access to quality and consistent mental health care. Severe lack of inpatient beds. - Physician

Lack of Providers

There are not enough providers to possibly help all the people who need help with mental health issues in our community. In addition, the systems we do have are not easy to navigate for someone not in crisis. If the person is in crisis, the systems we have now are impossible to navigate without help. There is not enough access to get appointments or assessments for not only therapists, but also psychiatrists and prescribers. – Social Services Provider

Rising numbers of depression and anxiety diagnoses, far too few providers, and we need to address the access children have to social media and iPhones. This is the fundamental problem in our area, from both the number of citizens of all walks of life involved as well as the severity of it for the homeless and poor. – Physician

Not enough trained providers, and the cost. - Physician

The lack of mental health providers and the increasing cost of insurance. – Social Services Provider

Limited number of providers and treatment facilities available to take patients without insurance or Medicaid, and long waiting lists and times for the clinics that do take these patients. – Physician

Especially for children, there are few providers and waiting lists to get in for a diagnosis. – Social Services Provider

Not enough mental health providers, not enough insurance coverage, private and federal, and stigma. – Physician

A growing number of children and adults are experiencing mental health challenges but aren't able to access the help they need. For people without insurance or on Medicaid, there aren't enough providers willing to work for the low wages available at the nonprofit organizations that provide free or sliding scale services (reimbursements for these services are woefully inadequate and haven't kept pace with market forces in the private sector). And there's a shortage of psychiatrists and other prescribers. – Social Services Provider

Lack of enough psychiatrists and counselors, need for housing and other supports. - Physician

Limited access to mental health providers due to poor reimbursement for mental health services and burnout in mental health providers as a consequence of the pandemic. – Health Provider

Inadequate provider numbers, inadequate inpatient beds. – Physician

Shortage of services and providers. Lack of providers and services that accept Medicaid. – Advanced Practice Provider

Lack of adequate number of mental health professionals in the community, stigma associated with mental health issues and lack of connection for residents. – Health Provider



Affordable Care/Services

Accessing and paying for services. Culturally and linguistically concordant services are very limited. Emergency services for youth with mental health crises. – Public Health Representative

Longitudinal counseling services at an affordable cost. - Physician

Resources are difficult to obtain or afford for people in the community. - Health Provider

Access to affordable services. - Social Services Provider

Lack of Diversity in Providers

Lack of diversity in mental health professionals. - Health Provider

Lack of therapists of color and financial barriers to seeking care. Co-occurring mental health and substance use disorders. – Community Leader

Lack of available diverse mental health therapists and available services for long term treatment. – Community Leader

People don't have access to culturally appropriate therapists. Pastors need additional training to support individuals that are seeking mental health services. Taboo of mental health support. Drug and alcohol addiction is more likely due to unresolved trauma and mental health issues left untreated and or addressed. – Social Services Provider

Denial/Stigma

Stigma. Access to care. Cost of Care. What social media has done to kids. - Community Leader

Stigma still remains in the community and health care settings. We have a shortage of psychiatric providers for prescribing psychotropic medications. We have many counselors, but the general public does not know where to look to locate these providers. We also need more bilingual providers for our diverse patient populations, we also have need for more substance use disorder programs. Funding for mental health care is also a concern. The highest levels of care (inpatient, partial hospital, IOP) are a challenge to get covered by insurance, thus those most in crisis will continue to struggle due to inability to afford care. Our greatest need is for those living with severe persistent mental illness. We do not have enough resources for them to obtain consistent care and oversight. They struggle from hospitalization to hospitalization and crisis to crisis. – Physician

Stigma in seeking help. - Business Leader

Multiple Factors

Despair, hopelessness, bigotry, racism, social isolation, and bullying. - Health Provider

Poverty, homelessness, addiction, and family history. - Social Services Provider

Social media, isolation, and lack of support. - Social Services Provider

Awareness/Education

Understanding where to start to look for support. Recognizing when someone may need help themselves and recognizing challenges in others. Cost barriers. – Public Health Representative

They are doing nothing to promote brain health. We can only help those who help themselves. Money is not the solution without compliance and accessibility to programming. – Community Leader

Access for Medicare/Medicaid Patients

The explosion with private practices is causing an exclusion of Medicaid, Medicare and low-income populations from being served. The programs available to serve these populations are non-profits who cannot pay staff enough money to stay long term. More integrated health care needs to happen. Research shows people with a mental health dx. are 3 times more likely to have heart disease and other health complications. You cannot treat the body without treating the mind. Many health systems have moved in this direction, but some are still in the dark ages. — Health Provider

Impact on Quality of Life

I think we continue to see people in a crisis of their mental health, this leads to other medical challenges. For our population I think that it would be beneficial to educate and catch early through primary care, counseling, and tele medicine options so we can meet patients prior to their crisis. With our aging population, undiagnosed or changed mental and behavioral disease to become part of the care plan. – Business Leader

Insurance Issues

Many people have difficulty finding an outpatient mental health provider to accept their payer source, or it is taking weeks to months to schedule an appointment for outpatient follow-up. – Health Provider



Incidence/Prevalence

Children and adults struggling with mental health diagnosis, stress that causes depression and anxiety, drug induces mental illness and medical follow up services. – Health Provider

Workforce Challenges

Lack of stability in workforce due to undiagnosed and misdiagnosed mental health issues. Not enough physicians that are representative of all communities. Not enough holistic options as solutions. – Social Services Provider

Follow Up/Support

There is nowhere for them to follow up, or the wait is extremely long when they are discharged from the hospital. – Health Provider







DEATH, DISEASE & CHRONIC CONDITIONS

LEADING CAUSES OF DEATH

Distribution of Deaths by Cause

Together, cancers and heart disease account for over one-third of all deaths in the Metro Area, with COVID-19 responsible for another 9.5% of deaths in 2020.





Sources: • CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics. Data extracted June 2024.

Notes:
• Lung disease is CLRD, or chronic lower respiratory disease.

Age-Adjusted Death Rates for Selected Causes

AGE-ADJUSTED DEATH RATES

In order to compare mortality in the region with other localities (in this case, Nebraska and Iowa and the United States), it is necessary to look at rates of death — these are figures which represent the number of deaths in relation to the population size (such as deaths per 100,000 population, as is used here).

Furthermore, in order to compare localities without undue bias toward younger or older populations, the common convention is to adjust the data to some common baseline age distribution. Use of these "age-adjusted" rates provides the most valuable means of gauging mortality against benchmark data, as well as Healthy People 2030 objectives.

Note that deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10). Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population.



The following chart outlines 2018-2020 annual average age-adjusted death rates per 100,000 population for selected causes of death in the Metro Area.

For infant mortality data, see Birth Outcomes & Risks in the Births section of this report.

Leading causes of death are discussed in greater detail in subsequent sections of this report.

Age-Adjusted Death Rates for Selected Causes (2018-2020 Deaths per 100,000 Population)

	Metro Area	NE	IA	US	HP2030
Malignant Neoplasms (Cancers)	154.6	148.5	151.3	146.5	122.7
Diseases of the Heart	139.7	144.8	170.3	164.4	127.4*
COVID-19	79.3	84.4	99.0	85.0	-
Fall-Related Deaths (65+)	63.6	67.8	87.4	67.1	63.4
Chronic Lower Respiratory Disease (CLRD)	44.8	45.7	42.3	38.1	_
Alzheimer's Disease	38.5	30.0	30.9	30.9	_
Unintentional Injuries	36.5	40.3	42.9	51.6	43.2
Cerebrovascular Disease (Stroke)	34.8	33.0	32.3	37.6	33.4
Diabetes Mellitus	25.9	25.1	22.3	22.6	_
Alcohol-Induced	15.8	12.0	9.9	11.9	_
Intentional Self-Harm (Suicide)	14.1	14.8	16.7	13.9	12.8
Pneumonia/Influenza	13.7	14.2	13.8	13.4	_
Cirrhosis/Liver Disease	11.7	9.8	9.7	12.5	10.9
Kidney Diseases	10.2	10.3	9.7	12.8	_
Motor Vehicle Deaths	9.5	12.3	10.5	11.4	10.1
Drug-Induced	8.9	7.4	9.4	21.0	_
Homicide	4.4	3.0	3.0	6.1	5.5

- Sources: CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and
 - US Department of Health and Human Services. Healthy People 2030. August 2020. http://www.healthypeople.gov.

 *The Healthy People 2030 Heart Disease target is adjusted to account for all diseases of the heart.

Note:



CARDIOVASCULAR DISEASE

ABOUT HEART DISEASE & STROKE

Heart disease and stroke can result in poor quality of life, disability, and death. Though both diseases are common, they can often be prevented by controlling risk factors like high blood pressure and high cholesterol through treatment.

In addition, making sure people who experience a cardiovascular emergency — like stroke, heart attack, or cardiac arrest — get timely recommended treatment can reduce their risk for long-term disability and death. Teaching people to recognize symptoms is key to helping more people get the treatment they need.

- Healthy People 2030 (https://health.gov/healthypeople)

Age-Adjusted Heart Disease & Stroke Deaths

Heart Disease Deaths


Between 2018 and 2020, there was an annual average age-adjusted heart disease mortality rate of 139.7 deaths per 100,000 population in the Metro Area.

BENCHMARK ► Lower than the Iowa and US mortality rates.

DISPARITY ► Notably higher in Pottawattamie County and among Black or African American residents.

Heart Disease: Age-Adjusted Mortality (2018-2020 Annual Average Deaths per 100,000 Population)

Healthy People 2030 = 127.4 or Lower (Adjusted)



- Sources: CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics. Data extracted June 2024.
 - US Department of Health and Human Services. Healthy People 2030. August 2020. http://www.healthypeople.gov
 - The Healthy People 2030 Heart Disease target is adjusted to account for all diseases of the heart.
 Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10).
 - Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population.




The greatest share of cardiovascular deaths is attributed to heart

disease.

Notes

Heart Disease: Age-Adjusted Mortality by Race (2018-2020 Annual Average Deaths per 100,000 Population)

Healthy People 2030 = 127.4 or Lower (Adjusted)



Notes:

- CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics. Data extracted June 2024.

- US Department of Health and Human Services. Healthy People 2030. August 2020. http://www.healthypeople.gov
 The Healthy People 2030 Heart Disease target is adjusted to account for all diseases of the heart.
 Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10).
- Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population.

Heart Disease: Age-Adjusted Mortality Trends (Annual Average Deaths per 100,000 Population)

Healthy People 2030 = 127.4 or Lower (Adjusted)

	2011-2013	2012-2014	2013-2015	2014-2016	2015-2017	2016-2018	2017-2019	2018-2020	
Metro Area	151.3	150.4	151.2	143.3	141.3	137.1	139.8	139.7	
─ NE	147.2	145.9	148.5	145.9	148.0	145.1	146.6	144.8	
—IA	168.4	165.5	162.3	160.3	163.7	165.1	168.5	170.3	
<u></u> -US	171.3	169.6	168.9	167.5	166.3	164.7	163.4	164.4	

- Sources: CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics. Data extracted June 2024.

 US Department of Health and Human Services. Healthy People 2030. August 2020. http://www.healthypeople.gov

 - The Healthy People 2030 Heart Disease target is adjusted to account for all diseases of the heart.
 - Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10).
 - Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population.




Stroke Deaths

Between 2018 and 2020, there was an annual average age-adjusted stroke mortality rate of 34.8 deaths per 100,000 population in the Metro Area.

DISPARITY ► Affects Metro Area Black residents much more so than White residents or Hispanic residents.

Stroke: Age-Adjusted Mortality (2018-2020 Annual Average Deaths per 100,000 Population)

Healthy People 2030 = 33.4 or Lower



rces: • CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics. Data extracted June 2024.


US Department of Health and Human Services. Healthy People 2030. August 2020. http://www.healthypeople.gov

Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10).

Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population.

Stroke: Age-Adjusted Mortality by Race (2018-2020 Annual Average Deaths per 100,000 Population)

Healthy People 2030 = 33.4 or Lower



 CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics. Data extracted June 2024.

US Department of Health and Human Services. Healthy People 2030. August 2020. http://www.healthypeople.gov

Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10).

Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population.



Notes:

Stroke: Age-Adjusted Mortality Trends (Annual Average Deaths per 100,000 Population)

Healthy People 2030 = 33.4 or Lower



Sources: • CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics. Data extracted June 2024.

• US Department of Health and Human Services. Healthy People 2030. August 2020. http://www.healthypeople.gov

Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10).

Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population.

Prevalence of Heart Disease & Stroke

Prevalence of Heart Disease

Notes:

A total of 8.2% of surveyed adults report that they suffer from or have been diagnosed with heart disease, such as coronary heart disease, angina, or heart attack.


BENCHMARK ► Higher than both state percentages but below the US.

TREND Marks a statistically significant increase.

DISPARITY ► Highest in Pottawattamie County. Strong correlation with age.

Prevalence of Heart Disease

Metro Area





- Behavioral Risk Factor Surveillance System Survey Data. Atlanta, Georgia. United States Department of Health and Human Services, Centers for Disease Control
 and Prevention (CDC): 2022 Nebraska and Iowa data.
- 2023 PRC National Health Survey, PRC, Inc.

Notes: • Asked of all respondents.

Includes diagnoses of heart attack, angina, or coronary heart disease.



Prevalence of Stroke

A total of 3.3% of surveyed adults report that they suffer from or have been diagnosed with cerebrovascular disease (a stroke).


BENCHMARK Higher than the Nebraska prevalence but lower than the national prevalence.

TREND ► Increasing since 2011.

DISPARITY ► Highest in Douglas County (especially Northeast Omaha). Increases with age.

Prevalence of Stroke

Metro Area



Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Item 23]

Behavioral Risk Factor Surveillance System Survey Data. Atlanta, Georgia. United States Department of Health and Human Services, Centers for Disease Control
and Prevention (CDC): 2022 Nebraska and Iowa data.

2023 PRC National Health Survey, PRC, Inc.

Notes:


Asked of all respondents.

Key Informant Input: Heart Disease & Stroke

The greatest share of key informants taking part in an online survey characterized *Heart Disease & Stroke* as a "moderate problem" in the community.

Perceptions of Heart Disease and Stroke as a Problem in the Community (Metro Area, 2024)







Among those rating this issue as a "major problem," reasons related to the following:

Incidence/Prevalence

In my clinical work I am starting to see younger and younger patients hospitalized for heart disease and stroke. – Physician

This is one of the leading causes of morbidity and mortality. - Public Health Representative

Statistics. The CDC, National Institute of Health and American Heart Association. - Social Services Provider

More and more people are having complications from heart disease and stroke at a younger age. – Health Provider

Too prevalent and not enough people seeking care. - Health Provider

Heart disease continues to be a number one cause of death in the United States. - Physician

These are very common in our community and in the hospital. - Physician

It's the number one cause of death in patients 40 and older. – Physician

Death rates or impairments. - Health Provider

High use of Emergency Departments for heart attacks and stroke. Many risk factors made worse by issues with social determinants of health. – Health Provider

Because of the high prevalence of hypertension and diabetes. - Social Services Provider

I hear a lot about the risks of heart disease and stroke in both men and women. - Social Services Provider

Awareness/Education

Lack of early and continuous education regarding heart disease and stroke, unsafe walking trails, lack of culturally sensitive food reduction education programming, lack of affordable fruit and vegetables in North Omaha; food deserts; systemic issues related to social determinants of health, and lack of AA providers, nutritionists in health care systems. Lack of affordable medicines that could treat obesity prevalent among minority populations. Finally the death rates associated with heart disease and stroke reveals that these issues are major problems. – Community Leader

Education about prevention and importance of treatment support for adherence with medication and lifestyle changes once diagnosed. – Community Leader

Not understanding signs of stroke in time, or having someone to assist when it is happening. - Health Provider

Obesity

Heart attack and stroke are closely related to increase in obesity, decreased physical activity and sedentary lifestyle. Access to food and nutrition security play a role in increasing risk. These two vascular conditions remain the majority of health problems and deaths. – Physician

People are fat and they don't eat well. - Social Services Provider

Obesity and its resulting co-morbidities are a leading problem. Also, even with access to health care, there is a lack of trust in the health care system from our community, which COVID has exacerbated. – Health Provider

Lifestyle

Lack of mobility, food with high grams of salt or sugar. Lifestyles very sedentary. – Social Services Provider Poor lifestyle choices. – Physician

Poor diets, lack of exercise and infrequent medical contact. - Community Leader

It is on the rise due to a lack of healthy dieting, exercise, lifestyles. - Health Provider

Aging Population

With the aging population and the continued diet and weight epidemic, it causes an increase in stroke and heart disease. – Business Leader

I believe we have a higher frequency of heart disease and stroke due to a combination of factors, including an aging community, high level of poverty, and lower educational attainment. – Community Leader

Access to Care/Services

Heart disease. Prescription medicine cost is outrageous. - Community Leader

Desperate access to health care, disparate health care experiences, unhealthy lifestyle habits, chronic stress and weathering, diseases of despair. – Health Provider



Access to Affordable Healthy Food

I don't think a person could find/buy healthy food to literally save their life. Corporate America decides what we have access to and its mostly over processed garbage with no health value. The very few foods that might be better for people such as organic and overpriced and unreachable by a great number of our population. There should not be a "Health Food" section in a store. It should all be healthy or at least reasonably. Corporate America and big pharma only make huge profits by keeping people fat and sick. – Health Provider

Racial Disparities

High rates of CAD within our community with notable ethnic and racial disparities, as well as SES barriers. Access to healthy food. Ability to exercise are contributors. Access to goal directed medical therapy and rehab options after an event are also issues. – Physician

Alcohol/Drug Use

Alcohol, drugs, tobacco, unhealthy diet and lack of exercise. - Business Leader

Cultural/Personal Beliefs

Culture and genetics. Lack of intervention and prevention early. - Social Services Provider

Diagnosis/Treatment

Because of the prevalence of untreated and under treated risk factors for heart disease and stroke. – Physician

Vulnerable Populations

The Latino community is highly affected by heart disease and has minimal access to education and prevention. – Health Provider



CANCER

ABOUT CANCER

The cancer death rate has declined in recent decades, but over 600,000 people still die from cancer each year in the United States. Death rates are higher for some cancers and in some racial/ethnic minority groups. These disparities are often linked to social determinants of health, including education, economic status, and access to health care.

Interventions to promote evidence-based cancer screenings — such as screenings for lung, breast, cervical, and colorectal cancer — can help reduce cancer deaths. Other effective prevention strategies include programs that increase HPV vaccine use, prevent tobacco use and promote quitting, and promote healthy eating and physical activity. In addition, effective targeted therapies and personalized treatment are key to helping people with cancer live longer.

Healthy People 2030 (https://health.gov/healthypeople)

Age-Adjusted Cancer Deaths

All Cancer Deaths

Between 2018 and 2020, the Metro Area reported an annual average age-adjusted cancer mortality rate of 154.6 deaths per 100,000 population.


BENCHMARK ▶ Fails to satisfy the Healthy People 2030 objective.

TREND The mortality rate has decreased over time, in keeping with state and national trends.

DISPARITY ► Notably higher among Black residents.

Cancer: Age-Adjusted Mortality (2018-2020 Annual Average Deaths per 100,000 Population)

Healthy People 2030 = 122.7 or Lower



Sources: • CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics. Data extracted June 2024.

US Department of Health and Human Services. Healthy People 2030. August 2020. http://www.healthypeople.gov

Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10).


Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population.



Notes:

Cancer: Age-Adjusted Mortality by Race (2018-2020 Annual Average Deaths per 100,000 Population)

Healthy People 2030 = 122.7 or Lower



Sources: • CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics. Data extracted June 2024.

US Department of Health and Human Services. Healthy People 2030. August 2020. http://www.healthypeople.gov

Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10).

Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population.

Cancer: Age-Adjusted Mortality Trends (Annual Average Deaths per 100,000 Population)

Healthy People 2030 = 122.7 or Lower

	2011-2013	2012-2014	2013-2015	2014-2016	2015-2017	2016-2018	2017-2019	2018-2020	
Metro Area	178.5	174.8	172.4	166.3	162.5	156.9	155.5	154.6	
─ NE	163.4	161.9	159.6	157.0	154.7	152.2	150.2	148.5	
—IA	170.0	167.7	166.2	163.3	160.6	157.7	154.7	151.3	
— US	166.2	162.7	160.1	157.6	155.6	152.5	149.3	146.5	

Sources: • CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics, Data extracted June 2024.

US Department of Health and Human Services. Healthy People 2030. August 2020. http://www.healthypeople.gov

Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10).
 Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population.



Notes:

Notes:

Cancer Deaths by Site

Lung cancer is the leading cause of cancer deaths in the Metro Area.

Other leading sites include prostate cancer, female breast cancer, and colorectal cancer (both sexes).

BENCHMARK

Lung Cancer ► Fails to satisfy the Healthy People 2030 objective.

Female Breast Cancer ▶ Fails to satisfy the Healthy People 2030 objective.

Prostate Cancer ► Higher than the national rate. Fails to satisfy the Healthy People 2030 objective.

Colorectal Cancer ▶ Fails to satisfy the Healthy People 2030 objective.

Age-Adjusted Cancer Death Rates by Site (2018-2020 Annual Average Deaths per 100,000 Population)

	Metro Area	NE	IA	US	HP2030
ALL CANCERS	154.6	148.5	151.3	146.5	122.7
Lung Cancer	34.3	31.8	36.3	33.4	25.1
Prostate Cancer	21.9	18.7	20.2	18.5	16.9
Female Breast Cancer	19.6	20.8	17.9	19.4	15.3
Colorectal Cancer	13.4	14.9	13.9	13.1	8.9

- Sources: CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics. Data extracted June 2024.
- US Department of Health and Human Services. Healthy People 2030. August 2020. http://www.healthypeople.gov
 Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10). Notes:
 - Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population.




Cancer Incidence

"Incidence rate" or "case rate" is the number of newly diagnosed cases in a given population in a given year, regardless of outcome. These rates are also age-adjusted. It is usually expressed as cases per 100,000 population per year.

The highest cancer incidence rates are for female breast cancer and prostate cancer.

DISPARITY Note that Sarpy County reports a significantly lower incidence rate for lung cancer when compared with the other counties. In contrast, the highest incidence rate for colorectal cancer is in Pottawattamie County.

Cancer Incidence Rates by Site (Annual Average Age-Adjusted Incidence per 100,000 Population, 2016-2020)



Sources: • State Cancer Profiles.

Center for Applied Research and Engagement Systems (CARES), University of Missouri Extension. Retrieved June 2024 via SparkMap (sparkmap.org).

This is a spark of the sp

This indicator reports the age adjusted incidence rate (cases per 100,000 population per year) of cancers, adjusted to 2000 US standard population.



Prevalence of Cancer


A total of 8.8% of surveyed Metro Area adults report having ever been diagnosed with cancer.

BENCHMARK ► Below the Nebraska and Iowa percentages.

DISPARITY ► Statistically high in Pottawattamie County. The prevalence increases with age and is reported more often among White respondents.

Prevalence of Cancer

Metro Area




Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Item 24]

- Behavioral Risk Factor Surveillance System Survey Data. Atlanta, Georgia. United States Department of Health and Human Services, Centers for Disease Control and Prevention (CDC): 2022 Nebraska and lowa data.
- 2023 PRC National Health Survey, PRC, Inc.

Notes: • Reflects all respondents.

Prevalence of Cancer (Metro Area, 2024)



Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Item 24]

Notes: • Reflects all respondents



Cancer Screenings

The American Cancer Society recommends that both men and women get a cancer-related checkup during a regular doctor's checkup. It should include examination for cancers of the thyroid, testicles, ovaries, lymph nodes, oral cavity, and skin, as well as health counseling about tobacco, sun exposure, diet and nutrition, risk factors, sexual practices, and environmental and occupational exposures. Screening levels in the community were measured in the PRC Community Health Survey relative to the following cancer sites:

FEMALE BREAST CANCER

The US Preventive Services Task Force (USPSTF) recommends biennial screening mammography for women age 50 to 74 years.

CERVICAL CANCER

The US Preventive Services Task Force (USPSTF) recommends screening for cervical cancer every 3 years with cervical cytology alone in women age 21 to 29 years. For women age 30 to 65 years, the USPSTF recommends screening every 3 years with cervical cytology alone, every 5 years with high-risk human papillomavirus (hrHPV) testing alone, or every 5 years with hrHPV testing in combination with cytology (cotesting). The USPSTF recommends against screening for cervical cancer in women who have had a hysterectomy with removal of the cervix and do not have a history of a high-grade precancerous lesion (i.e., cervical intraepithelial neoplasia [CIN] grade 2 or 3) or cervical cancer.

COLORECTAL CANCER

The US Preventive Services Task Force (USPSTF) recommends screening for colorectal cancer starting at age 50 years and continuing until age 75 years.

 US Preventive Services Task Force, Agency for Health care Research and Quality, US Department of Health & Human Services

Note that other organizations (e.g., American Cancer Society, American Academy of Family Physicians, American College of Physicians, National Cancer Institute) may have slightly different screening guidelines.

Among women age 50 to 74, 82.2% have had a mammogram within the past 2 years.

BENCHMARK ► Higher than the Nebraska and US percentages.

DISPARITY ► Reported <u>most</u> often in the Western Douglas County area (not shown).

Among Metro Area women age 21 to 65, 73.5% have had appropriate cervical cancer screening.

BENCHMARK ► Fails to satisfy the Healthy People 2030 objective.

TREND ▶ Decreasing significantly since 2011.

DISPARITY ► Lowest in Southeast Omaha and Cass County (not shown).

"Appropriate cervical cancer screening" includes Pap smear testing (cervical cytology) every 3 years in women age 21 to 29 and Pap smear testing and/or HPV testing every 5 years in women age 30 to 65.




"Appropriate colorectal cancer screening" includes a fecal occult blood test within the past year and/or lower endoscopy (sigmoidoscopy or colonoscopy) within the past 10 years.


Among all adults age 50 to 75, 78.9% have had appropriate colorectal cancer screening.

BENCHMARK ► Well above the state and national percentages, and satisfying the Healthy People 2030 objective.

TREND ► Increasing significantly since 2011.



- Sources: 2024 PRC Community Health Survey, PRC, Inc. [Items 101-103]
 - Behavioral Risk Factor Surveillance System Survey Data. Atlanta, Georgia. United States Department of Health and Human Services, Centers for Disease Control
 and Prevention (CDC): 2022 Nebraska and Iowa data.
 - 2023 PRC National Health Survey, PRC, Inc.
 - US Department of Health and Human Services. Healthy People 2030. August 2020. http://www.healthypeople.gov
- Notes: Each indicator is shown among the gender and/or age group specified.



2018

2021

2024

2011

2015

2018

2021

2024

Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Items 101-103]

2021

2024

US Department of Health and Human Services. Healthy People 2030. August 2020. http://www.healthypeople.gov

2011

2015

Notes: • Each indicator is shown among the gender and/or age group specified.



2011

2015

2018

Key Informant Input: Cancer

The greatest share of key informants taking part in an online survey characterized *Cancer* as a "moderate problem" in the community.

Perceptions of Cancer as a Problem in the Community (Metro Area, 2024)



Among those rating this issue as a "major problem," reasons related to the following:

Incidence/Prevalence

So many people have it. - Social Services Provider

Excessively high rates of breast cancer, prostate, colon, and pancreatic cancer. - Community Leader

Cancer is running rampant in all communities. I believe that there are cures out there, but that they are not accessible to all individuals. – Social Services Provider

Many community members have been diagnosed with different types of cancer and eventually some died, while others are battling it with no resources at all. – Business Leader

More and more individuals are diagnosed with cancer and at younger ages. - Community Leader

The numbers are increasing for all age groups, and I have personally experienced loss to cancer in all age groups. – Social Services Provider

It seemed for a decent period of time that treatments were more successful and there were fewer cases of cancer. It seems to have come back with a vengeance in recent times. I fear people can't get in to see a care provider and time is of the essence, especially with cancer. I have lost several friends and a young niece in the last year alone and know or am aware of several people battling for their lives currently. I think that pharmaceuticals' profit comes before everything and there's no money for curing or saving people. — Health Provider

It is rising in incidence and becoming more prevalent. More people are diagnosed at an earlier stage, which means more people are living with or from the consequences of cancer. Additional resources to this survivor population would be beneficial. — Physician

Increasing prevalence rates of many cancers. - Health Provider

Prevention/Screenings

Patients have limited access to convenient and timely cancer screenings. – Physician

Limited access to health care affects screenings and diagnostic testing for some patients, resulting in later detection. – Advanced Practice Provider

There's limited use of preventive screenings and health education campaigns to reduce cancer risk factors (aside from tobacco use). With many behaviors like smoking, limited physical activity, and poor nutrition, being prevalent in the community, I think more needs to be done focused on prevention of cancer. I think more policy work in this realm would be helpful. – Public Health Representative

Environmental Contributors

Many young people are developing cancer at an early age. The health of the air, soil and water are in question with major power and energy plants in the north Omaha area. America should quicky adapt regulations from Australia and Europe around contaminate allowed in our environment and our food. — Social Services Provider

Rising rates in rural areas, potentially as a result of agricultural pollution. Rural areas are less likely to have access to preventative or screening resources. – Business Leader



Obesity

The epidemic of obesity, nitrate content of our water supply due to agricultural inputs, poor diet high in processed sugars and oils, continued tobacco use and alcohol consumption well above any recommended limits. – Physician

Access to Affordable Healthy Food

Food choices are unhealthy and unaffordable for healthy foods to maintain. - Health Provider

Access to Care for Uninsured/Underinsured

Many people go untreated for a long period of time due to not having health insurance and or a trusted care physician. – Social Services Provider

Racial Disparities

There are disparities due to ethnicities and access to care. – Health Provider



RESPIRATORY DISEASE

ABOUT RESPIRATORY DISEASE

Respiratory diseases affect millions of people in the United States. ... More than 25 million people in the United States have asthma. Strategies to reduce environmental triggers and make sure people get the right medications can help prevent hospital visits for asthma. In addition, more than 16 million people in the United States have COPD (chronic obstructive pulmonary disease), which is a major cause of death. Strategies to prevent the disease — like reducing air pollution and helping people quit smoking — are key to reducing deaths from COPD.

- Healthy People 2030 (https://health.gov/healthypeople)

Note that this section also includes data relative to COVID-19 (coronavirus disease).


Age-Adjusted Respiratory Disease Deaths

Lung Disease Deaths

Between 2018 and 2020, the Metro Area reported an annual average age-adjusted lung disease mortality rate of 44.8 deaths per 100,000 population.

DISPARITY ► Lowest in Sarpy County. Notably higher among Metro Area Black residents.

CLRD: Age-Adjusted Mortality (2018-2020 Annual Average Deaths per 100,000 Population)



Notes:

Sources: • CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics, Data extracted June 2024.

CLRD is chronic lower respiratory disease

- Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10).
- Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population




Note: Here, lung disease

reflects chronic lower respiratory disease

(CLRD) deaths and includes conditions such as emphysema, chronic bronchitis, and asthma.

CLRD: Age-Adjusted Mortality by Race (2018-2020 Annual Average Deaths per 100,000 Population)



CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and. Informatics. Data extracted June 2024.

Notes:

- CLRD is chronic lower respiratory disease.
 Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10).
- Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population.

CLRD: Age-Adjusted Mortality Trends (Annual Average Deaths per 100,000 Population)

	2011-2013	2012-2014	2013-2015	2014-2016	2015-2017	2016-2018	2017-2019	2018-2020	
Metro Area	50.4	50.4	53.6	52.6	52.4	49.4	48.7	44.8	
─ NE	49.0	49.1	50.2	50.6	51.2	49.7	48.8	45.7	
—IA	47.4	47.4	48.2	48.5	48.1	46.3	44.7	42.3	
US	42.0	41.7	41.8	41.3	41.0	40.4	39.6	38.1	

Sources: • CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics. Data extracted June 2024.

CLRD is chronic lower respiratory disease

- Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10).
- Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population.




Pneumonia/Influenza Deaths

Between 2018 and 2020, the Metro Area reported an annual average age-adjusted pneumonia/influenza mortality rate of 13.7 deaths per 100,000 population.

DISPARITY ▶ Particularly high in Pottawattamie County.


Pneumonia/Influenza: Age-Adjusted Mortality (2018-2020 Annual Average Deaths per 100,000 Population)



Sources:

- CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics. Data extracted June 2024.
- Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10).
- Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population.

Pneumonia/Influenza: Age-Adjusted Mortality by Race (2018-2020 Annual Average Deaths per 100,000 Population)



CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics. Data extracted June 2024.

Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10). Notes:

Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population



Pneumonia/Influenza: Age-Adjusted Mortality Trends (Annual Average Deaths per 100,000 Population)



	2011-2013	2012-2014	2013-2015	2014-2016	2015-2017	2016-2018	2017-2019	2018-2020
Metro Area	14.7	15.8	17.0	16.3	15.8	15.0	14.8	13.7
—−NE	13.8	14.1	15.5	15.4	15.8	15.5	15.6	14.2
——IA	16.4	15.7	15.2	13.2	13.0	13.5	14.0	13.8
US	15.3	15.2	15.4	14.6	14.3	14.2	13.8	13.4

 CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics Data extracted June 2024

Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10).

Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population.

Prevalence of Respiratory Disease

Asthma

Notes

A total of 12.5% of Metro Area adults have asthma.


BENCHMARK ► Well above the Nebraska and Iowa percentages but lower than found across the US.

TREND ► Marks a gradual but significant increase from earlier survey administrations.

DISPARITY Highest in Northeast Omaha. The prevalence decreases with age and household income level and is reported more often among women, Black or African American residents, and LGBTQ+ residents.

Prevalence of Asthma

Metro Area



Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Item 26]


- Behavioral Risk Factor Surveillance System Survey Data. Atlanta, Georgia. United States Department of Health and Human Services, Centers for Disease Control and Prevention (CDC): 2022 Nebraska and Iowa data.
 2023 PRC National Health Survey, PRC, Inc.

Asked of all respondents.

Survey respondents were asked to indicate whether they suffer from or have been diagnosed with various respiratory conditions, including asthma and COPD.



Prevalence of Asthma (Metro Area, 2024)



Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Item 26] Notes: Asked of all respondents.

Chronic Obstructive Pulmonary Disease (COPD)

A total of 5.9% of Metro Area adults suffer from chronic obstructive pulmonary disease (COPD).


BENCHMARK ► Well below the US figure.

TREND ▶ Decreasing significantly since 2011 (and especially since 2018).

DISPARITY Considerably higher in Pottawattamie County when compared with the other counties.

Prevalence of Chronic Obstructive Pulmonary Disease (COPD)

Metro Area



2024 PRC Community Health Survey, PRC, Inc. [Item 21]

- Behavioral Risk Factor Surveillance System Survey Data. Atlanta, Georgia. United States Department of Health and Human Services, Centers for Disease Control and Prevention (CDC): 2022 Nebraska and Iowa data.
 2023 PRC National Health Survey, PRC, Inc.

- Assect or all responsers.
 Includes those having ever suffered from or been diagnosed with COPD or chronic obstructive pulmonary disease, including bronchitis or emphysema.
 In prior data, the term "chronic lung disease" was used, which also included bronchitis or emphysema.



Note: COPD includes


lung diseases such as emphysema and chronic

bronchitis.

Key Informant Input: Respiratory Disease

The greatest share of key informants taking part in an online survey characterized *Respiratory Disease* as a "moderate problem" in the community (followed closely by "minor problem" ratings).

Perceptions of Respiratory Diseases as a Problem in the Community (Metro Area, 2024)



Among those rating this issue as a "major problem," reasons related to the following:

Incidence/Prevalence

Asthma and lung disease. - Health Provider

COPD and sleep apnea are prevalent. - Physician

High incidences of COPD. - Community Leader

Respiratory issues, such as asthma. - Health Provider

Tobacco Use

Smoking and noncompliance. - Physician

Large number of smokers or ex-smokers in the community. Community resistance to getting vaccinated. – Social Services Provider

Tobacco use is high in the community. Not all people received vaccines and or boosters. – Social Services Provider

Awareness/Education

Lack of awareness and options. – Business Leader

Environmental Contributors

Significant pollution in Omaha and Nebraska from pesticides and agricultural practices, as well as high rates of smoking. – Physician



INJURY & VIOLENCE

ABOUT INJURY & VIOLENCE

INJURY ► In the United States, unintentional injuries are the leading cause of death in children, adolescents, and adults younger than 45 years. ...Many unintentional injuries are caused by motor vehicle crashes and falls, and many intentional injuries involve gun violence and physical assaults. Interventions to prevent different types of injuries are key to keeping people safe in their homes, workplaces, and communities.

Drug overdoses are now the leading cause of injury deaths in the United States, and most overdoses involve opioids. Interventions to change health care providers' prescribing behaviors, distribute naloxone to reverse overdoses, and provide medications for addiction treatment for people with opioid use disorder can help reduce overdose deaths involving opioids.

VIOLENCE ► Almost 20,000 people die from homicide every year in the United States, and many more people are injured by violence. ...Many people in the United States experience physical assaults, sexual violence, and gun-related injuries. Adolescents are especially at risk for experiencing violence. Interventions to reduce violence are needed to keep people safe in their homes, schools, workplaces, and communities.

Children who experience violence are at risk for long-term physical, behavioral, and mental health problems. Strategies to protect children from violence can help improve their health and well-being later in life.

- Healthy People 2030 (https://health.gov/healthypeople)

Unintentional Injury

Age-Adjusted Unintentional Injury Deaths

Between 2018 and 2020, there was an annual average age-adjusted unintentional injury mortality rate of 36.5 deaths per 100,000 population in the Metro Area.


BENCHMARK ► Lower than the Iowa and US mortality rates. Satisfies the Healthy People 2030 objective.

DISPARITY ► Highest in Cass County.



Unintentional Injuries: Age-Adjusted Mortality (2018-2020 Annual Average Deaths per 100,000 Population)

Healthy People 2030 = 43.2 or Lower




Sources: • CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics. Data extracted June 2024.

- US Department of Health and Human Services. Healthy People 2030. August 2020. http://www.healthypeople.gov Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10). Notes:
 - Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population

Unintentional Injuries: Age-Adjusted Mortality by Race (2018-2020 Annual Average Deaths per 100,000 Population)

Healthy People 2030 = 43.2 or Lower



Sources: • CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics. Data extracted June 2024.

US Department of Health and Human Services. Healthy People 2030. August 2020. http://www.healthypeople.gov

Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10).

• Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population.



Unintentional Injuries: Age-Adjusted Mortality Trends

(Annual Average Deaths per 100,000 Population)

Healthy People 2030 = 43.2 or Lower



Sources: • CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics. Data extracted June 2024.

US Department of Health and Human Services. Healthy People 2030. August 2020. http://www.healthypeople.gov


Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10).

Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population.

Leading Causes of Unintentional Injury Deaths

Poisoning (including unintentional drug overdose), falls, and motor vehicle crashes accounted for most unintentional injury deaths in the Metro Area between 2018 and 2020.

Leading Causes of Unintentional Injury Deaths (Metro Area, 2018-2020)



CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics. Data extracted June 2024.

RELATED ISSUE
For more information
about unintentional drugrelated deaths, see also
Substance Use in the
Modifiable Health Risks
section of this report.



Falls

ABOUT FALLS

Falls are the leading cause of fatal and nonfatal injuries for persons age 65 and older Even when those injuries are minor, they can seriously affect older adults' quality of life by inducing a fear of falling, which can lead to self-imposed activity restrictions, social isolation, and depression.


Modifiable fall risk factors include muscle weakness, gait and balance problems, poor vision, use of psychoactive medications, and home hazards. Falls among older adults can be reduced through evidence-based fall-prevention programs that address these modifiable risk factors. Most effective interventions focus on exercise, alone or as part of a multifaceted approach that includes medication management, vision correction, and home modifications.

- Division of Unintentional Injury Prevention, National Center for Injury Prevention and Control, CDC

Among surveyed Metro Area adults age 45 and older, most have not fallen in the past year.

RELATED ISSUE
For fall-related mortality
data, see Age-Adjusted
Death Rates for Selected
Causes in the Leading
Causes of Death section
of this report.

Number of Falls in Past 12 Months (Adults Age 45 and Older; Metro Area, 2024)



Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Item 331]


Notes: • Asked of all respondents age 45+



However, 35.3% have experienced a fall at least once in the past year.

TREND ► Increasing significantly from 2018 findings.

Fell One or More Times in the Past Year (Adults Age 45 and Older)



Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Items 331-332] Asked of those respondents age 45 and older

Intentional Injury (Violence)

Age-Adjusted Homicide Deaths

The Metro Area reports 4.4 homicides per 100,000 population (2018-2020 annual average ageadjusted rate).


BENCHMARK ► Worse than both state homicide rates but below the US rate. Satisfies the Healthy People 2030 objective.

TREND The rate has decreased over the past decade, mirroring the Nebraska trend.

DISPARITY > The homicide rate is nine times higher among Black residents than White residents.

Homicide: Age-Adjusted Mortality (2018-2020 Annual Average Deaths per 100,000 Population)

Healthy People 2030 = 5.5 or Lower



Notes:


- Sources: CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics. Data extracted June 2024.
 - US Department of Health and Human Services. Healthy People 2030. August 2020. http://www.healthypeople.gov
 - Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10)
 - Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population.

RELATED ISSUE See also Mental Health (Suicide) in the General Health Status section of this report.



Homicide: Age-Adjusted Mortality by Race (2018-2020 Annual Average Deaths per 100,000 Population)

Healthy People 2030 = 5.5 or Lower




Sources: • CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics. Data extracted June 2024.
US Department of Health and Human Services. Healthy People 2030. August 2020. http://www.healthypeople.gov
Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10).

Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population.

Homicide: Age-Adjusted Mortality Trends (Annual Average Deaths per 100,000 Population)

Healthy People 2030 = 5.5 or Lower



	2011-2013	2012-2014	2013-2015	2014-2016	2015-2017	2016-2018	2017-2019	2018-2020
Metro Area	6.2	6.1	6.6	5.6	5.3	4.2	4.0	4.4
──NE	3.8	3.7	3.8	3.6	3.3	2.6	2.6	3.0
—IA	2.0	2.2	2.4	2.6	2.9	3.0	2.9	3.0
US	5.4	5.3	5.3	5.2	5.3	5.7	6.0	6.1

Sources: • CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics. Data extracted June 2024.

 US Department of Health and Human Services. Healthy People 2030. August 2020. http://www.healthypeople.gov
 Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10). Notes:

Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population.



Violent Crime


Violent Crime Rates

The Metro Area reported 369.3 violent crimes per 100,000 population (2015-2017).

BENCHMARK ► Well above both state violent crime rates.

DISPARITY ► Highest in Douglas County; particularly low in Sarpy and Cass counties.

Violent Crime (Rate per 100,000 Population, 2015-2017)



Sources:

- Federal Bureau of Investigation, FBI Uniform Crime Reports (UCR).
 Center for Applied Research and Engagement Systems (CARES), University of Missouri Extension. Retrieved June 2024 via SparkMap (sparkmap.org).
 This indicator reports the rate of violent crime offenses reported by the sheriff's office or county police department per 100,000 residents. Violent crime includes
 - homicide, forcible rape, robbery, and aggravated assault.

 Participation by law enforcement agencies in the UCR program is voluntary. Sub-state data do not necessarily represent an exhaustive list of crimes due to gaps in
 - reporting. Also, some institutions of higher education have their own police departments, which handle offenses occurring within campus grounds; these offer are not included in the violent crime statistics but can be obtained from the Uniform Crime Reports Universities and Colleges data tables.

Violent crime is composed of four offenses (FBI Index offenses): murder and non-negligent manslaughter; forcible rape; robbery; and aggravated assault.

Note that the quality of crime data can vary widely from location to location, depending on the consistency and completeness of reporting among various jurisdictions.



Respondents were read:
"By an intimate partner, I
mean any current or
former spouse, boyfriend,
or girlfriend. Someone
you were dating, or
romantically or sexually
intimate with would also
be considered an intimate
partner."

Intimate Partner Violence


A total of 19.8% of Metro Area adults acknowledge that they have ever been hit, slapped, pushed, kicked, or otherwise hurt by an intimate partner.

TREND ▶ Denotes a statistically significant increase since 2011.

DISPARITY ► Highest in Pottawattamie County and eastern Omaha.

Have Ever Been Hit, Slapped, Pushed, Kicked, or Hurt in Any Way by an Intimate Partner

Metro Area



Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Item 33]

2023 PRC National Health Survey, PRC, Inc.


es: • Asked of all respondents.

Perceived Neighborhood Safety

While most Metro Area adults consider their own neighborhoods to be "extremely safe" or "quite safe," 18.9% consider them only "slightly safe" or "not at all safe."

DISPARITY ► Least favorable in Douglas County (especially east of 72nd Street). Less favorable perceptions are also expressed among women, young adults, those in low-income households, Hispanic residents, Black or African American residents, those of Diverse Races, and LGBTQ+ residents.

Perceived Safety of Own Neighborhood (Metro Area, 2024)









Perceive Own Neighborhood as "Slightly" or "Not At All" Safe

Metro Area



Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Item 319]
Notes: • Asked of all respondents.

Perceive Own Neighborhood as "Slightly" or "Not At All" Safe (Metro Area, 2024)



Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Item 319]

otes:


• Asked of all respondents.



Key Informant Input: Injury & Violence

The largest share of key informants taking part in an online survey characterized *Injury* & *Violence* as a "moderate problem" in the community.

Perceptions of Injury and Violence as a Problem in the Community (Metro Area, 2024)



Among those rating this issue as a "major problem," reasons related to the following:

Incidence/Prevalence

We treat many patients who are victims of violence at the hospital, and violence has lifelong psychological and social impacts on victims. – Physician

The community is divisive and violent and on the increase. - Social Services Provider

Observed feedback from community groups, like Omaha 360. - Social Services Provider

Our organization focuses on sexual violence and domestic assault. Our recent report of domestic violence in Douglas County indicates an increase. – Business Leader

Crime. - Physician

Assault rates are extremely high. People in the community are prone to resorting to violence for even minor inconveniences. This violence is not only amongst intimate partners, but to those providing community services. – Health Provider

Significant community violence issues. Some areas of the city are distant from trauma centers. Only one level-one trauma center. – Physician

There are always reports on the news of injury or violence happening in our community, shootings, stabbings, drunk driving accidents, robberies, child abuse, human and sex trafficking, etc. – Social Services Provider

High prevalence of people experiencing violence. – Physician

Increased violence with not enough programs or pathways out of circumstances or to address systemic causes. – Business Leader

Turn on the news, it is all over the place. - Health Provider

Based on data from several community partners. - Health Provider

Gun Violence

The United States has an epidemic of gun violence. - Physician

We have a higher rate of shootings in North and South Omaha. Families are traumatized and looking or mental health support services. Our unhoused population is increasing looking for services. With not enough services or assistance available increases use of drugs and violence. – Social Services Provider

High incidences of shootings and assaults. - Community Leader

Frequent news reports of guns and other violence in Omaha. - Health Provider

Domestic/Family Violence

Domestic violence and instability in the home cause family displacement. Gun violence, loss of loved ones and if survived, disability, needing caregiver services, family strained relationships. Hit-and-run accidents causes loss of income, in a family, disability, loss of life and not being able to follow up due to the circumstances involved. – Health Provider

Domestic violence, unsafe roads, poor infrastructure, lack of safety regulations and safeguards. – Business Leader



Impact on Quality of Life

A large portion of the patients I serve list some form of past abuse in their chart, such as physical, emotional, or sexual abuse. We know the impact these adverse events have when experienced or witnessed as children, and as adults on long-term health. – Physician

Poverty, trauma, and mental health issues. - Health Provider

Traffic Fatalities

Traffic fatalities (particularly those for vulnerable roadway users such as cyclists/pedestrians/motorcyclists) continue to be a persistent problem. High-speeds, impairment from drugs and alcohol, distracted driving, and infrastructure that prioritizes high speeds all contribute to these issues and require cross-sectoral efforts to make progress. – Community Leader

Parental Influence

Poor parenting skills, single young women, drug abuse, alcohol intake, and domestic violence. – Social Services Provider

Social Media

News and social media. – Social Services Provider



DIABETES

ABOUT DIABETES

More than 30 million people in the United States have diabetes, and it's the seventh leading cause of death. ... Some racial/ethnic minorities are more likely to have diabetes. And many people with diabetes don't know they have it.

Poorly controlled or untreated diabetes can lead to leg or foot amputations, vision loss, and kidney damage. But interventions to help people manage diabetes can help reduce the risk of complications. In addition, strategies to help people who don't have diabetes eat healthier, get physical activity, and lose weight can help prevent new cases.


Healthy People 2030 (https://health.gov/healthypeople)

Age-Adjusted Diabetes Deaths

Between 2018 and 2020, there was an annual average age-adjusted diabetes mortality rate of 25.9 deaths per 100,000 population in the Metro Area.

DISPARITY Highest in Pottawattamie County. Much higher among Black residents.

Diabetes: Age-Adjusted Mortality (2018-2020 Annual Average Deaths per 100,000 Population)




Sources: • CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics. Data extracted June 2024.

Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10).

Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population.




Diabetes: Age-Adjusted Mortality by Race (2018-2020 Annual Average Deaths per 100,000 Population)



- Sources: CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics. Data extracted June 2024.
 - Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10).
 - Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population.

Diabetes: Age-Adjusted Mortality Trends (Annual Average Deaths per 100,000 Population)



23.5

21.3

21.9

21.3

21.6

21.5

22.3

•IA

-US

18.8

21.3

21.2 CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and

24.4

 Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10).
 Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population. Notes:

23.8

21.3

20.7

21.2



Prevalence of Diabetes

A total of 11.2% of Metro Area adults report having been diagnosed with diabetes.


TREND While the prevalence of diabetes has remained fairly stable, the prevalence of borderline or pre-diabetes has increased significantly over time (not shown below).

DISPARITY Diabetes prevalence is highest in Northeast Omaha. Diabetes increases with age and decreases with household income level, and is reported more often among men and Black or African American respondents.

Prevalence of Diabetes

Another 12.1% of adults have been diagnosed with "pre-diabetes" or "borderline" diabetes.

Metro Area




Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Item 106]

- Behavioral Risk Factor Surveillance System Survey Data. Atlanta, Georgia. United States Department of Health and Human Services, Centers for Disease Control and Prevention (CDC): 2022 Nebraska and Iowa data.
- 2023 PRC National Health Survey, PRC, Inc.

Asked of all respondents.

Prevalence of Diabetes (Metro Area, 2024)



Sources: Notes:

- 2024 PRC Community Health Survey, PRC, Inc. [Item 106]
- - Asked of all respondents.

 Excludes gestational diabetes (occurring only during pregnancy).



Age-Adjusted Kidney Disease Deaths

ABOUT KIDNEY DISEASE & DIABETES

Chronic kidney disease (CKD) is common in people with diabetes. Approximately one in three adults with diabetes has CKD. Both type 1 and type 2 diabetes can cause kidney disease. CKD often develops slowly and with few symptoms. Many people don't realize they have CKD until it's advanced and they need dialysis (a treatment that filters the blood) or a kidney transplant to survive.


Centers for Disease Control and Prevention (CDC) https://www.cdc.gov/diabetes/managing/diabetes-kidney-disease.html

Between 2018 and 2020, there was an annual average age-adjusted kidney disease mortality rate of 10.2 deaths per 100,000 population in the Metro Area.

BENCHMARK ► Lower than the national mortality rate.

DISPARITY Lowest in Sarpy County. By race, the mortality rate is much higher among Black residents than White or Hispanic residents.


Kidney Disease: Age-Adjusted Mortality (2018-2020 Annual Average Deaths per 100,000 Population)



- CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics. Data extracted June 2024.
- Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10).
 Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population. Notes:




Kidney Disease: Age-Adjusted Mortality by Race (2018-2020 Annual Average Deaths per 100,000 Population)



- Sources: CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics. Data extracted June 2024.
 - Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10).
 Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population.

Kidney Disease: Age-Adjusted Mortality Trends (Annual Average Deaths per 100,000 Population)



	2011-2013	2012-2014	2013-2015	2014-2016	2015-2017	2016-2018	2017-2019	2018-2020
Metro Area	11.6	11.4	12.2	11.1	11.1	10.2	10.8	10.2
─ NE	9.8	10.4	10.9	10.7	10.1	9.5	10.1	10.3
—IA	8.2	8.1	8.1	8.0	8.4	8.7	9.3	9.7
US	13.2	13.2	13.3	13.3	13.2	13.0	12.9	12.8

 CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Sources: Informatics. Data extracted June 2024. Notes:


Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10).
 Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population.



Key Informant Input: Diabetes

Over half of key informants taking part in an online survey characterized *Diabetes* as a "major problem" in the community.

Perceptions of Diabetes as a Problem in the Community (Metro Area, 2024)



Among those rating this issue as a "major problem," reasons related to the following:

Affordable Medications/Supplies

Lack of access to first line medications, such as GLP1 agonists and SGLT2 inhibitors, due to cost, poor insurance coverage, or just not available in pharmacies. Overabundance of unhealthy foods in grocery stores and fast food. – Physician

Being able to afford diabetic medications. Some newer diabetic medications are not in stock even when insurance will cover the cost, making it difficult for diabetics to get access to their medications. – Health Provider

Affordable medications and access to healthy affordable foods. Medications for diabetes are all separate and have to be ordered individually. Some pharmacies do not carry all diabetic supplies. Rural communities have barriers to accessing their medications, especially if the person is disabled. – Health Provider

Cost of life saving medications. - Social Services Provider

Access to medications, including insulin and type-2 drugs. - Community Leader

Access to affordable medications, and prevention of the disease. - Community Leader

Medication costs both with and without insurance. Significant supply chain issues affecting availability of medications (specifically GLP-1s although some insulin as well). Access to culturally and linguistically appropriate education. Access to healthy foods. Limitation of fresh fruits, vegetables (non-carb heavy) options at food pantries. Access to safe spaces to exercise. – Physician

Costs of medications and testing supplies. Limited access to newer agents for testing, such as continuous glucose monitoring and newer medications, due to lack of formulary coverage by insurers. – Advanced Practice Provider

Access to medications, like GLP-1 and SGLT2, and medication classes. – Physician

Ability to afford CGMs, pumps, and expensive medications and insulin. - Physician

Inability to afford the medication (or access the medications that work best for diabetes and weight management like GLP-1 agonists that were being obtained by people wanting to lose weight). Food insecurity and additional costs of healthier food choices. Lack of access to safe and affordable physical activity options. Lack of diabetes prevention programs in addition to culturally competent diabetes education classes that are conveniently located within the communities most impacted. Lack of podiatrists in north and south Omaha. – Community Leader

Cost of supplies and adherence to care. - Health Provider

From what I have heard, access to medications needed to treat diabetes are being used for other things like weight loss. – Business Leader

Access to Affordable Healthy Food

Accessing healthy foods. - Community Leader

Due to lack of income, no access to fresh food items and healthier items. - Social Services Provider

Access to healthy, less starchy food. - Community Leader

Overall accessibility to healthy food options. - Community Leader

Access to healthy foods. Access to affordable medication, health literacy and prevention. - Health Provider



Access to fresh healthy foods, eating healthy is more expensive and more work to prepare, we need to educate our community on simple choices they can make to improve their eating habits. I also feel that access to affordable health care is a challenge for our "working" population. Medical coverage as well as medication is expensive as are diabetic supplies. – Social Services Provider

Access to healthy nutrition, exercise, and the cost of diabetes medications. A healthy lifestyle is not the default for many people. – Physician

Awareness/Education

Education and access to healthy foods and exercise. Lack of intervention early. – Social Services Provider Early intervention, education, and compliance. – Physician

Health literacy and education on the diagnosis and lifestyle changes, finances for medications. – Social Services Provider

I would say the largest gap is awareness. We know how well we are managing the patients that come to us but do not have a full grasp of all the individuals that need support that aren't engaged with a health provider. So, awareness is an opportunity, financial support to help manage diabetes and resources to support healthy living and lifestyle options. – Business Leader

Lack of quality education regarding diabetes management. Ability for lower income individuals to afford needed supplies for testing their blood sugar. – Physician

Nutrition

Nutrition and foot care. - Community Leader

The fact that foods that cause diabetes are allowed on the shelf in the first place. The lack of education of proper nutrition for our children and adults. Access to healthy food options in North Omaha food deserts. – Social Services Provider

A culture in which unhealthy foods are subsidized and marketed to people starting at a young age. Some people do not have access to safe outdoor areas or gyms for exercise, but many Medicaid plans now cover gym access so that is helpful. the cost of newer Diabetes medicines means that people with lower income or wealth may not have the same access to medicine as those who are wealthy. – Physician

Being able to follow directions on making good food choices, being able to afford healthy choices, and following medical guidelines. – Health Provider

Food security, access to treatment, access to the ability to be active and safe, access to medications due to cost. – Physician

Obesity

Obesity rates, older population, access to affordable drug therapies. - Business Leader

Increasing number of overweight people who are developing diabetes at a very young age. – Physician Obesity. – Health Provider

Diabetes is secondary to obesity. Healthy food is expensive and people living in poverty must opt for less expensive food that's not good for you. People, in general, are living a sedentary lifestyle as well. – Social Services Provider

I think many people are not diagnosed, but likely have Type 2, especially with the weight epidemic. Scheduling with a care provider can take far too long and I think people give up and just chance it. I don't think people are educated as to where they can get help/insulin and especially if they don't have insurance. I believe there is a federal cap on insulin for \$25 or \$35 now, but for some people that means not eating that week if they have to buy it, and they likely won't. — Health Provider

Prevention/Screenings

Screening, diagnosis, and management of diabetes, particularly culturally and linguistically appropriate care. – Public Health Representative

Testing kits, appropriate diet, and preventative measures. – Business Leader

Tertiary prevention, which is too late. Difficulty accessing and affording medications. Little understanding or evaluation of the environment's impact on diabetes, such as endocrine disrupters and pollution. – Health Provider

I think it is those that are pre-diabetic. We need more assessment and data monitoring to prevent diabetes and screen those individuals that do not know they are pre-diabetic. – Public Health Representative

Access to Care/Services

Lack of insurance, expense of medications. - Physician

Lack of adequate access to care, education, and treatment. Financial limitations on food choices and optimal medication choices. Issues with weight management for all of the previously noted reasons. – Physician



Weight loss programs, availability of weight loss medication GLP-1, and access to endocrinologists. – Physician Lack of access to the latest technology and treatments. – Health Provider

Diagnosis/Treatment

Lack of integrated services to combine classic medication-based approach with community-based nutrition, activity, and peer support. Medication availability and pricing. Education for those at risk of prediabetes. Access to technology that may help to assist with long term control and prevent diabetes in the first place. – Health Provider

Mental health services are not being offered in conjunction with medical treatment for those on dialysis. Diabetic and pre-diabetic patients should have access to trained and paid community health workers located at support groups outside of medical institutions within their communities. Recent data has revealed that African American diabetes patients on dialysis nationally are not put on kidney transplants lists at the level of white patients and given the racial and ethnic health disparities among diabetes deaths among African Americans in Douglas County, I see this as possibly the biggest challenge. — Community Leader

Disease Management

Diabetes is not managed well. - Health Provider

Adherence to medication and access to providers of diabetes. Lack of physical activity and obesity. - Physician

Vulnerable Populations

The Latino community is highly affected by diabetes and prevention education. - Health Provider

For our homeless population it is medication access and making sure that they do not lose their medication. For those that are lower income, it is affording their medications. – Health Provider

Affordable Care/Services

Insurance coverage for continuous glucose monitoring. Access to healthy, minimally processed foods. Safe places to exercise. – Physician

Cost. - Health Provider

Comorbidities

Mental health, addiction, and poverty. - Social Services Provider

Inequality

High maternal and infant mortality, both driven in part by discrimination and access issues. Inability to access mental health services. High rate of syphilis and other STIs. – Physician

Lack of Diversity in Providers

Access to professionals of color. – Health Provider

Lifestyle

T2D and unhealthy lifestyles lead to this diagnosis. – Social Services Provider

Parental Influence

For adults in the community, it is a choice, for children the food by parents who do not pay attention to leading factors as it relates to a restrictive diet or one that includes balance; many have access but may be income challenged or not know what to purchase/prepare. – Business Leader



DISABLING CONDITIONS

Activity Limitations

ABOUT DISABILITY & HEALTH

Studies have found that people with disabilities are less likely to get preventive health care services they need to stay healthy. Strategies to make health care more affordable for people with disabilities are key to improving their health.

In addition, people with disabilities may have trouble finding a job, going to school, or getting around outside their homes. And they may experience daily stress related to these challenges. Efforts to make homes, schools, workplaces, and public places easier to access can help improve quality of life and overall well-being for people with disabilities.


Healthy People 2030 (https://health.gov/healthypeople)

A total of 29.6% of Metro Area adults are limited in some way in some activities due to a physical, mental, or emotional problem.

TREND ▶ Denotes a statistically significant increase since 2011.

DISPARITY Highest in Pottawattamie County and Northeast Omaha. The percentage increases with age, decreases with household income levels, and is reported more often among White adults, Black or African American adults, those of Diverse Races, and those who identify as LGBTQ+.

Limited in Activities in Some Way Due to a Physical, Mental or Emotional Problem




Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Items 83-84]

2023 PRC National Health Survey, PRC, Inc.

Notes: • Asked of all respondents.



Limited in Activities in Some Way Due to a Physical, Mental or Emotional Problem (Metro Area, 2024)



Sources:

• 2024 PRC Community Health Survey, PRC, Inc. [Item 83]

• Asked of all respondents.



Alzheimer's Disease

ABOUT DEMENTIA

Alzheimer's disease is the most common cause of dementia. Nearly 6 million people in the United States have Alzheimer's, and that number will increase as the population ages.

Dementia refers to a group of symptoms that cause problems with memory, thinking, and behavior. People with dementia are more likely to be hospitalized, and dementia is linked to high health care costs.

While there's no cure for Alzheimer's disease, early diagnosis and supportive care can improve quality of life. And efforts to make sure adults with symptoms of cognitive decline — including memory loss — are diagnosed early can help improve health outcomes in people with dementia. Interventions to address caregiving needs can also help improve health and well-being in people with dementia.

Healthy People 2030 (https://health.gov/healthypeople)

Age-Adjusted Alzheimer's Disease Deaths


Between 2018 and 2020, there was an annual average age-adjusted Alzheimer's disease mortality rate of 38.5 deaths per 100,000 population in the Metro Area.

BENCHMARK ► Much worse than state and national mortality rates.

TREND ► Increasing over the past decade.

DISPARITY ► Lowest in Cass County. Higher among Metro Area Black residents than White residents.

Alzheimer's Disease: Age-Adjusted Mortality (2018-2020 Annual Average Deaths per 100,000 Population)




Sources: • CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics. Data extracted June 2024.

Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10).

Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population



Alzheimer's Disease: Age-Adjusted Mortality by Race (2018-2020 Annual Average Deaths per 100,000 Population)



Notes:

- Sources: CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics. Data extracted June 2024.
 - Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10).
 Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population.

Alzheimer's Disease: Age-Adjusted Mortality Trends (Annual Average Deaths per 100,000 Population)



	2011-2013	2012-2014	2013-2015	2014-2016	2015-2017	2016-2018	2017-2019	2018-2020
Metro Area	28.1	28.5	31.1	32.4	34.6	34.3	36.0	38.5
─ NE	24.7	23.3	23.4	24.3	26.5	27.4	28.7	30.0
——IA	30.3	29.4	29.2	30.3	32.2	32.8	32.1	30.9
—US	23.1	24.7	27.4	29.7	30.2	30.6	30.4	30.9

Sources:

- CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics. Data extracted June 2024.
- Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10).
 Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population. Notes:



Key Informant Input: Disabling Conditions

Key informants taking part in an online survey most often characterized *Disabling Conditions* as a "moderate problem" in the community.

Perceptions of Disability & Chronic Pain as a Problem in the Community (Metro Area, 2024)



Among those rating this issue as a "major problem," reasons related to the following:

Aging Population

Increasing dementia incidence with aging of the population. Many people living with chronic pain and have limited treatment options. – Health Provider

Wider incidences of these conditions are observed with older adults in our community, especially those vulnerable elderly who live alone and need assistance to remain in their homes because they cannot afford an assisted living facility. Also, higher numbers of these conditions among a growing population of elderly in the homeless shelters. – Health Provider

As the population ages, including myself, health changes drastically. All of the above-listed problems along with no longer having a decent balance mean problems and injuries for this demographic. I believe many people, as they age, don't have anyone or anyone who cares to look out for them and help them with their health issues. There are many for-profit care givers for home health care and monitoring, but not many can afford such a service and fall through the cracks. – Health Provider

Baby boomers are aging, and dementia is becoming more predominant. – Health Provider

Aging population and health issues that come with age, such as cognitive, mental, complex chronic conditions. – Social Services Provider

Aging population with increasing number of comorbid chronic health conditions that lead to all of these concerns. This is higher in those with multiple disparities in SDOH. – Physician

Access to Care/Services

Our neighborhoods, gathering places, and hospital systems are not physically or digitally accessible. Although programs exist, it is not easy for people with physical, sensory, intellectual, and cognitive disabilities and needs to access services. – Health Provider

In my clinical work, I see chronic pain and dementia as severe problems that the health care community and the general public do not have adequate resources or education to manage. These are both complex disease processes that have no simple solutions. The needs of individuals suffering from chronic pain and dementia require multiple resources and medical providers working in coordination. The current medical system is fragmented and there simply isn't the time or personnel to do a thorough job. Not to mention the support that dementia caregivers need. And the opioid crisis's contribution to poor outcomes and unrealistic expectations for quick relief. – Physician

No post-acute facilities to take care of a dementia patient. - Social Services Provider

Not enough beds available at high quality skilled nursing and long-term care facilities. - Physician

Incidence/Prevalence

Again, it just seems as if disabling conditions, cancer, diabetes, etc. are becoming more and more prevalent than historically. I am also seeing a difference in the availability of resources based on socioeconomic status and a shortage of health care professionals who specialize in these areas. – Social Services Provider



There is a high proportion of patients in my health care system with disabling conditions, including decreased mobility, dementia, and vision and hearing problems. – Physician

Injuries, illnesses, and sanitary conditions. - Business Leader

I hear about people dealing with dementia more and more every single day, both in my professional as well as my personal life. Doctors don't always know how to address the symptoms patients and their families are presenting with. Memory care facilities typically have a waitlist and so many don't take Medicaid. There is a lack of resources available for those trying to care for their loved ones at home. – Social Services Provider

Impact on Quality of Life

These challenges can drive the onset of other health issues or exacerbate existing health issues significantly. – Business Leader

Many of the people we see on a daily basis have some sort of medical condition that prevents them from leading a normal life. Many times, people are not able to get their level of care met if they have housing instability nor are actively homeless. This is a situation where one small push can send someone over the edge to then needing higher levels of care and get stuck in a cycle of needing care but then not being to get that care, which spirals out into all aspects of life. — Community Leader

Any of these disability conditions severely impact the ability to manage complex chronic health conditions like hypertension and diabetes from a physical standpoint as well as a mental effort standpoint. Disability conditions take a big toll on mental health, and as mental health declines other chronic conditions suffer. Moreover, these disabling conditions all lead to diminished or complete absence of physical activity. Physical activity by itself is incredibly important in the improvement of chronic conditions both physical and mental. — Physician

Diagnosis/Treatment

Diets and medications that have side effects which may not be discussed during patient visits. I also believe there are more young people with auditory issues because of the earpieces and headphones that allow significant levels of high decibel sounds directly into the ear. – Business Leader

There are many disabling conditions not seen by the eye that often go undiagnosed. When they involve in-depth diagnoses, it is hard to get appointments or to get clear results and next steps. – Social Services Provider

Conditions are taken care of when they start, and it becomes more difficult to care for. - Social Services Provider

Affordable Care/Services

Lack of affordable personal care assistance options, lack of support family, friends for those in the community, and the ability to afford hearing and vision care DME. – Health Provider

Comorbidities

Autism, emotional and mental health, physical conditions, such as diabetes, arthritis, orthopedic, and traumatic brain injury. – Social Services Provider

Follow Up/Support

The lack of, support for, and insurance coverage for these services, which is especially problematic for low-income persons. – Physician

Lifestyle

It is secondary to obesity and a sedentary lifestyle. – Social Services Provider

Autoimmune Diseases

Autoimmune diseases, such as thyroid disease. – Community Leader

Transportation

Poor transportation systems in town. - Physician





BIRTHS

PRENATAL CARE

ABOUT INFANT HEALTH

Keeping infants healthy starts with making sure women get high-quality care during pregnancy and improving women's health in general. After birth, strategies that focus on increasing breastfeeding rates and promoting vaccinations and developmental screenings are key to improving infants' health. Interventions that encourage safe sleep practices and correct use of car seats can also help keep infants safe.

The infant mortality rate in the United States is higher than in other high-income countries, and there are major disparities by race/ethnicity. Addressing social determinants of health is critical for reducing these disparities.

Healthy People 2030 (https://health.gov/healthypeople)

Early and continuous prenatal care is the best assurance of infant health.

Between 2017 and 2019, 5.1% of all Metro Area births did <u>not</u> receive prenatal care in the first six months of pregnancy.

BENCHMARK ► Higher than the Iowa percentage but lower than the US.

TREND ► Increasing over the past decade.

DISPARITY ► Highest in Douglas County.

Lack of Prenatal Care in the First Six Months of Pregnancy (Percentage of Live Births, 2017-2019)



Sources: • Centers for Disease Control and Prevention, National Vital Statistics System. Accessed via CDC WONDER. Centers for Disease Control and Prevention, Wide-Ranging Online Data for Epidemiologic Research.

Center for Applied Research and Engagement Systems (CARES), University of Missouri Extension. Retrieved June 2024 via SparkMap (sparkmap.org).

This indicates reported the percentage of women who do not able to propose of the percentage of t

This indicator reports the percentage of women who do not obtain prenatal care before their seventh month of pregnancy (if at all).



Note:

Lack of Prenatal Care in the First Six Months of Pregnancy (Percentage of Live Births)



	2008-2010	2011-2013	2014-2016	2017-2019
Metro Omaha	4.0%	4.0%	5.0%	5.1%
—−NE	4.3%	4.6%	5.2%	4.9%
—IA	4.1%	3.8%	4.0%	4.3%
—US	4.3%	5.0%	5.7%	6.1%

Sources:

Centers for Disease Control and Prevention, National Vital Statistics System. Accessed via CDC WONDER. Centers for Disease Control and Prevention, Wide-Ranging Online Data for Epidemiologic Research.

Center for Applied Research and Engagement Systems (CARES), University of Missouri Extension. Retrieved June 2024 via SparkMap (sparkmap.org).

This indicator reports the percentage of women who do not obtain prenatal care before their seventh month of pregnancy (if at all).



BIRTH OUTCOMES & RISKS

Low-Weight Births

A total of 8.2% of 2016-2022 Metro Area births were low-weight.


BENCHMARK ► Higher than the Iowa percentage.

DISPARITY ► Highest among Douglas County births.

Low-Weight Births (Percent of Live Births, 2016-2022)

Low birthweight babies, those who weigh less than 2,500 grams (5 pounds, 8 ounces) at birth, are much more prone to illness and neonatal death than are babies of normal birthweight.

Largely a result of receiving poor or inadequate prenatal care, many low-weight births and the consequent health problems are preventable.



Sources: • University of Wisconsin Population Health Institute, County Health Rankings.

Center for Applied Research and Engagement Systems (CARES), University of Missouri Extension. Retrieved June 2024 via SparkMap (sparkmap.org).

This indicator reports the percentage of total births that are low birth weight (Under 2500g).



Infant Mortality

Between 2018 and 2020, there was an annual average of 5.8 infant deaths per 1,000 live births.


BENCHMARK ► Higher than the Iowa infant mortality rate.

DISPARITY ► Highest in Pottawattamie County. Much higher among Black births than White or Hispanic births in the Metro Area.

Infant Mortality Rate

(Annual Average Infant Deaths per 1,000 Live Births, 2018-2020)

Healthy People 2030 = 5.0 or Lower




Sources:

- CDC WONDER Online Query System. Centers for Disease Control and Prevention, National Center for Health Statistics, Division of Vital Statistics.
 Data extracted June 2024.
- US Department of Health and Human Services. Healthy People 2030. https://health.gov/healthypeople

Notes: • Infant deaths include deaths of children under 1 year old.

Infant Mortality Rate by Race/Ethnicity (Annual Average Infant Deaths per 1,000 Live Births, 2018-2020)

Healthy People 2030 = 5.0 or Lower



Sources:

- CDC WONDER Online Query System. Centers for Disease Control and Prevention, National Center for Health Statistics, Division of Vital Statistics.
 Data extracted June 2024.
- US Department of Health and Human Services. Healthy People 2030. https://health.gov/healthypeople

Notes: Infant deaths include deaths of children under 1 year old

Race categories reflect individuals without Hispanic origin.



Infant mortality rates

reflect deaths of children less than one year old per 1,000 live births.

Infant Mortality Trends (Annual Average Infant Deaths per 1,000 Live Births) Healthy People 2030 = 5.0 or Lower



	2011-2013	2012-2014	2013-2015	2014-2016	2015-2017	2016-2018	2017-2019	2018-2020
Metro Area	5.2	5.6	6.1	6.2	6.3	6.3	5.8	5.8
─ NE	5.2	5.1	5.5	5.8	5.8	5.8	5.4	5.4
—IA	4.8	4.9	4.5	5.1	5.2	5.4	5.1	4.8
US	6.0	5.9	5.9	5.9	5.8	5.7	5.6	5.5

Sources: • CDC WONDER Online Query System. Centers for Disease Control and Prevention, National Center for Health Statistics, Division of Vital Statistics. Data extracted June 2024.

Centers for Disease Control and Prevention, National Center for Health Statistics.

US Department of Health and Human Services. Healthy People 2030. https://health.gov/healthypeople
Rates are three-year averages of deaths of children under 1 year old per 1,000 live births. Notes:



FAMILY PLANNING

ABOUT FAMILY PLANNING

Nearly half of pregnancies in the United States are unintended, and unintended pregnancy is linked to many negative outcomes for both women and infants. ... Unintended pregnancy is linked to outcomes like preterm birth and postpartum depression. Interventions to increase use of birth control are critical for preventing unintended pregnancies. Birth control and family planning services can also help increase the length of time between pregnancies, which can improve health for women and their infants.

Adolescents are at especially high risk for unintended pregnancy. Although teen pregnancy and birth rates have gone down in recent years, close to 200,000 babies are born to teen mothers every year in the United States. Linking adolescents to youth-friendly health care services can help prevent pregnancy and sexually transmitted infections in this age group.

Healthy People 2030 (https://health.gov/healthypeople)

Births to Adolescent Mothers


Between 2016 and 2022, there were 16.5 births to adolescents age 15 to 19 per 1,000 women age 15 to 19 in the Metro Area.

DISPARITY ► Highest in Pottawattamie and Douglas counties. Much higher among Black or Hispanic females.

Teen Birth Rate

(Births to Adolescents Age 15-19 per 1,000 Females Age 15-19, 2016-2022)

Healthy People 2030 = 31.4 or Lower



Sources: • Centers for Disease Control and Prevention, National Vital Statistics System.

Center for Applied Research and Engagement Systems (CARES), University of Missouri Extension. Retrieved June 2024 via SparkMap (sparkmap.org).


• This indicator reports the rate of total births to women under the age of 15-19 per 1,000 female population age 15-19.



Teen Birth Rate

(Births to Adolescents Age 15-19 per 1,000 Females Age 15-19, 2016-2022)

Healthy People 2030 = 31.4 or Lower



Sources:

- Centers for Disease Control and Prevention, National Vital Statistics System.
- Center for Applied Research and Engagement Systems (CARES), University of Missouri Extension. Retrieved June 2024 via SparkMap (sparkmap.org).

Notes: • This indicator reports the rate of total births to women under the age of 15-19 per 1,000 female population age 15-19.

Race categories reflect individuals without Hispanic origin.

Key Informant Input: Infant Health & Family Planning

Over half of key informants taking part in an online survey largely characterized *Infant Health* & *Family Planning* as a "moderate problem" in the community.

Perceptions of Infant Health and Family Planning as a Problem in the Community (Metro Area, 2024)



Sources: • PRC Online Key Informant Survey, PRC, Inc.
Notes: • Asked of all respondents.

Among those rating this issue as a "major problem," reasons related to the following:

Racial Disparities

Infant mortality rate shows us that Black and Brown children are dying at birth or shortly after at a higher rate. Proper health and wellness are needed prior to conception and with toxins so high in our food and water, it is causing our babies to die before they are full term, die stillborn, or are born with autism and or major birth defects. – Social Services Provider

The infant mortality rates among Black babies are almost three times the rate of White babies. 70% of all Black births in Douglas County are to single mothers. – Community Leader

Disparities in infant mortality rates for Black families. Lack of equitable infant and maternal health outcomes. Lack of doula coverage and accessibility. Racism and providers not being held accountable for patient experience and poor outcomes. Historical trauma within communities of color contributes to the generations of mistrust. Siloed efforts to address this issue across the state of Nebraska. – Community Leader



Large African American population where high-risk births are associated. – Health Provider

Awareness/Education

There is confusion and lack of information/resources regarding abortions. Unwanted pregnancy occurs from assaults, and having a choice is a challenge. We see babies with serious health issues due to mothers not obtaining prenatal care. Medicaid is complicated for low-income and vulnerable populations to navigate regarding family planning and infant health issues. — Health Provider

Lack of education to get healthy foods. - Social Services Provider

Parents that are not aware of the need to follow up with pediatricians. Lack of understanding of resources available to parents in the community. – Health Provider

Government/Policy

Access to family planning is limited. Increased legislation on women's rights. Limited funding for reliable methods (IUDS, Nexplanon, etc.). Continued racial and ethnic disparities in infant outcomes. Limited access to culturally and linguistically specific resources and providers. – Physician

Because women do not have a right to decide what is best for their body. There is a specific hospital in Omaha that does not have adequate staff to instruct new mothers with what to expect during or after the birth of a child. – Social Services Provider

The political environment which impacts the information and resources available, such as comprehensive sexual health information and contraception. – Business Leader

Infant Mortality

Infant mortality rates. - Health Provider

Nebraska has a lower or worse than average maternal and infant mortality. We need to improve access and prenatal care to change this horrible statistic. – Physician

High infant mortality and low birth weights. High rates of unintended pregnancies. - Community Leader

Unplanned Pregnancy

Early age pregnancy, unplanned, access to sex health care, such as pregnancy preventive strategies, routine testing, routine supplies, etc. Social supportive environments and lack of empowerment. – Physician

Due to the number of unplanned births, the number of teen births, and infant mortality rates in our community. – Social Services Provider

Income/Poverty

Poverty, homelessness, infertility, addiction, and mental health challenges. Social media. – Social Services Provider

Poverty and lack of financial means. - Business Leader

Diagnosis/Treatment

Not prioritized enough during pregnancy. Lack of short-term and long-term family planning. – Social Services Provider

Lack of Providers

Insufficient providers. – Physician





MODIFIABLE HEALTH RISKS

NUTRITION

ABOUT NUTRITION & HEALTHY EATING

Many people in the United States don't eat a healthy diet. ...People who eat too many unhealthy foods — like foods high in saturated fat and added sugars — are at increased risk for obesity, heart disease, type 2 diabetes, and other health problems. Strategies and interventions to help people choose healthy foods can help reduce their risk of chronic diseases and improve their overall health.


Some people don't have the information they need to choose healthy foods. Other people don't have access to healthy foods or can't afford to buy enough food. Public health interventions that focus on helping everyone get healthy foods are key to reducing food insecurity and hunger and improving health.

Healthy People 2030 (https://health.gov/healthypeople)

Difficulty Accessing Fresh Produce

Most Metro Area adults report little or no difficulty buying fresh produce at a price they can afford.

Level of Difficulty Finding Fresh Produce at an Affordable Price (Metro Area, 2024)



Sources:

• 2024 PRC Community Health Survey, PRC, Inc. [Item 66]

• Asked of all respondents.

Respondents were asked, "How difficult is it for you to buy fresh produce like fruits and vegetables at a price you can afford? Would you say very difficult, somewhat difficult, not too difficult, or not at all difficult?"

RELATED ISSUE
See also Food Access in
the Social Determinants
of Health section of this
report.




However, 23.7% of Metro Area adults find it "very" or "somewhat" difficult to access affordable fresh fruits and vegetables.

BENCHMARK ► Lower than the national response.

DISPARITY ► Highest in eastern Omaha and in Pottawattamie County. Reported more often among women, young adults, those in low-income households, Hispanic respondents, Black or African American respondents, those of Diverse Races, and LGBTQ+ adults.

Find It "Very" or "Somewhat" Difficult to Buy Affordable Fresh Produce

Metro Area




Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Item 66]

2023 PRC National Health Survey, PRC, Inc.

Notes:

Asked of all respondents.

Find It "Very" or "Somewhat" Difficult to Buy Affordable Fresh Produce (Metro Area, 2024)



Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Item 66]

Asked of all respondents.




Sugar-Sweetened Beverages

A total of 26.8% of Metro Area adults report drinking an average of at least one sugarsweetened beverage per day in the past week.

DISPARITY ► Highest in eastern Omaha and Pottawattamie County. Reported more often among men, young adults, those living in low-income households, Hispanic respondents, Black or African American respondents, and LGBTQ+ respondents.

Had Seven or More Sugar-Sweetened Beverages in the Past Week


Metro Area



Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Item 312] Notes:

 Asked of all respondents

Had Seven or More Sugar-Sweetened Beverages in the Past Week (Metro Area, 2024)



Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Item 312]

Asked of all respondents.



PHYSICAL ACTIVITY

ABOUT PHYSICAL ACTIVITY

Physical activity can help prevent disease, disability, injury, and premature death. The Physical Activity Guidelines for Americans lays out how much physical activity children, adolescents, and adults need to get health benefits. Although most people don't get the recommended amount of physical activity, it can be especially hard for older adults and people with chronic diseases or disabilities.

Strategies that make it safer and easier to get active — like providing access to community facilities and programs — can help people get more physical activity. Strategies to promote physical activity at home, at school, and at childcare centers can also increase activity in children and adolescents.

- Healthy People 2030 (https://health.gov/healthypeople)

Leisure-Time Physical Activity

Just over one in four Metro Area adults (26.5%) reports no leisure-time physical activity in the past month.

BENCHMARK ► Higher than the Nebraska percentage but lower than the US figure. Satisfies the Healthy People 2030 objective.


TREND ▶ Despite a drop since 2021, denotes an unfavorable, statistically significant increase since 2011.

DISPARITY Least favorable in Southeast Omaha and in Pottawattamie County.

No Leisure-Time Physical Activity in the Past Month

Healthy People 2030 = 21.8% or Lower

Metro Area



Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Item 69]

Behavioral Risk Factor Surveillance System Survey Data. Atlanta, Georgia. United States Department of Health and Human Services, Centers for Disease Control
and Prevention (CDC): 2022 Nebraska and Iowa data.

2023 PRC National Health Survey, PRC, Inc.

US Department of Health and Human Services. Healthy People 2030. August 2020. http://www.healthypeople.gov

Notes:

Asked of all respondents.



Leisure-time physical

activity includes any



Activity Levels

Adults

ADULTS: RECOMMENDED LEVELS OF PHYSICAL ACTIVITY

For adults, "meeting physical activity recommendations" includes adequate levels of both aerobic and strengthening activities:

- Aerobic activity is one of the following: at least 150 minutes per week of light to moderate activity (such as walking), 75 minutes per week of vigorous activity (such as jogging), or an equivalent combination of both.
- Strengthening activity is at least two sessions per week of exercise designed to strengthen muscles (such as push-ups, sit-ups, or activities using resistance bands or
- 2013 Physical Activity Guidelines for Americans, US Department of Health and Human Services. www.cdc.gov/physicalactivity

A total of 26.4% of Metro Area adults regularly participate in adequate levels of both aerobic and strengthening activities (meeting physical activity recommendations).

BENCHMARK ► Higher than the state percentages but lower than the US. Fails to satisfy the Healthy People 2030 objective.


TREND ► Increasing significantly since 2018.

DISPARITY ► Lowest in Northeast Omaha and Pottawattamie County. Reported less often among women, older adults (age 65+), and those living in low-income households.

Meets Physical Activity Recommendations

Healthy People 2030 = 29.7% or Higher

Metro Area




2024 PRC Community Health Survey, PRC, Inc. [Item 110]
Behavioral Risk Factor Surveillance System Survey Data. Atlanta, Georgia. United States Department of Health and Human Services, Centers for Disease Control and Prevention (CDC): 2022 Nebraska and lowa data.
2023 PRC National Health Survey, PRC, Inc.
US Department of Health and Human Services. Healthy People 2030. August 2020. http://www.healthypeople.gov
Asked of all respondents.
Meeting both guidelines is defined as the number of persons age 18+ who report light or moderate aerobic activity for at least 150 minutes per week or who report vigorous physical activity? 5 minutes per week or an equivalent combination of moderate and vigorous-intensity activity and report doing physical activities specifically designed to strengthen muscles at least twice per week.



Meets Physical Activity Recommendations

(Metro Area, 2024)

Healthy People 2030 = 29.7% or Higher



Sources:

- 2024 PRC Community Health Survey, PRC, Inc. [Item 110] US Department of Health and Human Services. Healthy People 2030. August 2020. http://www.healthypeople.gov
- - Meeting both guidelines is defined as the number of persons age 18+ who report light or moderate aerobic activity for at least 150 minutes per week or who report vigorous physical activity 75 minutes per week or an equivalent combination of moderate and vigorous-intensity activity and report doing physical activities specifically designed to strengthen muscles at least twice per week.

Built Environment


Recreation & Fitness Facilities

In 2021, there were 17.5 recreation/fitness facilities for every 100,000 population in the Metro Area.

BENCHMARK ► Higher than the state and national ratios.

DISPARITY ► Highest in Douglas County.

Population With Recreation & Fitness Facility Access (Number of Recreation & Fitness Facilities per 100,000 Population, 2021)



Sources:
• US Census Bureau, County Business Patterns. Additional data analysis by CARES.

Center for Applied Research and Engagement Systems (CARES), University of Missouri Extension. Retrieved June 2024 via SparkMap (sparkmap.org).

- Recreation and fitness facilities are defined by North American Industry Classification System (NAICS) Code 713940, which include Establishments engaged in operating facilities which offer "exercise and other active physical fitness conditioning or recreational sports activities." Examples include athletic clubs, gymnasiums, dance centers, tennis clubs, and swimming pools. This indicator is relevant because access to recreation and fitness facilities encourages physical activity and other healthy behaviors.
- Counts of establishments <3 are suppressed.

Here, recreation/fitness facilities include establishments engaged in operating facilities which offer "exercise and other active physical fitness conditioning or recreational sports activities.

Examples include athletic clubs, gymnasiums, dance centers, tennis clubs, and swimming pools



Neighborhood Barriers

Survey respondents were next asked about the presence of five neighborhood factors that potentially prevent people from exercising, including lack (or poor condition) of sidewalks; heavy traffic; lack (or poor condition) of trails; crime; and lack of streetlights or nonworking streetlights.

Overall, a lack of sidewalks/poor sidewalks received the largest share of responses among Metro Area adults (mentioned by 20.6%), followed by heavy traffic and lack of (or poor condition) trails.

TREND ▶ With the exception of sidewalks, <u>each of these</u> barriers has worsened since 2015.

DISPARITY ► Residents of Sarpy County were <u>least</u> likely to mention these potential barriers to outdoor physical activity. Adults in eastern Omaha were far <u>more</u> likely to report these barriers.

Presence of Neighborhood Barriers That Prevent Physical Activity (Metro Area)


2015 2018 2021 2024




Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Items 314-318]
• Asked of all respondents.

Presence of Neighborhood Barriers That Prevent Physical Activity (By County; Metro Area, 2024)

Douglas Co. Sarpy Co. Cass Co. Pott. Co. Metro Area







Presence of Neighborhood Barriers That Prevent Physical Activity

(By Douglas County Subareas; Metro Area, 2024)

NE Omaha SE Omaha NW Omaha SW Omaha Western Douglas Metro Area



Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Items 314-318] Asked of all respondents.



WEIGHT STATUS

ABOUT OVERWEIGHT & OBESITY

Obesity is linked to many serious health problems, including type 2 diabetes, heart disease, stroke, and some types of cancer. Some racial/ethnic groups are more likely to have obesity, which increases their risk of chronic diseases.

Culturally appropriate programs and policies that help people eat nutritious foods within their calorie needs can reduce overweight and obesity. Public health interventions that make it easier for people to be more physically active can also help them maintain a healthy weight.

- Healthy People 2030 (https://health.gov/healthypeople)

Body Mass Index (BMI), which describes relative weight for height, is significantly correlated with total body fat content. The BMI should be used to assess overweight and obesity and to monitor changes in body weight. In addition, measurements of body weight alone can be used to determine efficacy of weight loss therapy. BMI is calculated as weight (kg)/height squared (m²). To estimate BMI using pounds and inches, use: [weight (pounds)/height squared (inches²)] x 703.

In this report, overweight is defined as a BMI of 25.0 to 29.9 kg/m² and obesity as a BMI ≥30 kg/m². The rationale behind these definitions is based on epidemiological data that show increases in mortality with BMIs above 25 kg/m². The increase in mortality, however, tends to be modest until a BMI of 30 kg/m² is reached. For persons with a BMI ≥30 kg/m², mortality rates from all causes, and especially from cardiovascular disease, are generally increased by 50 to 100 percent above that of persons with BMIs in the range of 20 to 25 kg/m².

 Clinical Guidelines on the Identification, Evaluation, and Treatment of Overweight and Obesity in Adults: The Evidence Report. National Institutes of Health. National Heart, Lung, and Blood Institute in Cooperation With The National Institute of Diabetes and Digestive and Kidney Diseases. September 1908

Adult Weight Status

CLASSIFICATION OF OVERWEIGHT AND OBESITY BY BMI	BMI (kg/m²)
Underweight	<18.5
Healthy Weight	18.5 – 24.9
Overweight	25.0 – 29.9
Obese	≥30.0

Source: Clinical Guidelines on the Identification, Evaluation, and Treatment of Overweight and Obesity in Adults: The Evidence Report. National Institutes of Health. National Heart, Lung, and Blood Institute in Cooperation With The National Institute of Diabetes and Digestive and Kidney Diseases. September 1998.



Here, "overweight" includes those respondents with a BMI value ≥25.

Overweight Status


Most Metro Area adults (72.2%) are overweight.

BENCHMARK ► Higher than the Nebraska and US figures.

TREND ► Increasing significantly since 2011.

DISPARITY ► Highest in Sarpy and Pottawattamie counties.

Prevalence of Total Overweight (Overweight and Obese)



Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Items 112, 313]
• Behavioral Risk Factor Surveillance System Survey Data. Atlanta, Georgia. United States Department of Health and Human Services, Centers for Disease Control and Prevention (CDC): 2022 Nebraska and Iowa data.

2023 PRC National Health Survey, PRC, Inc.

Based on reported heights and weights, asked of all respondents.
 The definition of overweight is having a body mass index (BMI), a ratio of weight to height (kilograms divided by meters squared), greater than or equal to 25.0, regardless of gender. The definition for obesity is a BMI greater than or equal to 30.0.

"Obese" (also included in overweight prevalence discussed previously) includes respondents with a BMI value ≥30.

The overweight prevalence above includes 38.4% of Metro Area adults who are obese.

BENCHMARK ► Worse than the Nebraska and US percentages. Fails to satisfy the Healthy People 2030 objective.

TREND ► Increasing significantly since 2011.


DISPARITY ► Considerably lower in Douglas County when compared to the other three counties. Higher among women, adults age 40 to 64, those in low-income households, and LGBTQ+ adults. Asian respondents are least likely to be obese in the Metro Area.



Prevalence of Obesity

Healthy People 2030 = 36.0% or Lower

Metro Area




2024 PRC Community Health Survey, PRC, Inc. [Item 112]
Behavioral Risk Factor Surveillance System Survey Data. Atlanta, Georgia. United States Department of Health and Human Services, Centers for Disease Control and Prevention Behavioral Hisk Factor Surveillance System Survey Data: Autama, Georga. Onited States Department of Factor Surveilland Iowa data.

2023 PRC National Health Survey, PRC, Inc.
US Department of Health and Human Services. Healthy People 2030. August 2020. http://www.healthypeople.gov
Based on reported heights and weights, asked of all respondents.

The definition of obesity is having a body mass index (BMI), a ratio of weight to height (kilograms divided by meters squared), greater than or equal to 30.0, regardless of gender.

Prevalence of Obesity (Metro Area, 2024)

Healthy People 2030 = 36.0% or Lower



Sources:

• 2024 PRC Community Health Survey, PRC, Inc. [Item 112]

• US Department of Health and Human Services. Healthy People 2030. August 2020. http://www.healthypeople.gov

Based on reported heights and weights, asked of all respondents.

• The definition of obesity is having a body mass index (BMI), a ratio of weight to height (kilograms divided by meters squared), greater than or equal to 30.0, regardless of gender.



The correlation between overweight and various health issues cannot be disputed.

Relationship of Overweight With Other Health Issues


Overweight and obese adults are more likely to report a number of adverse health conditions, as outlined in the following chart.

Relationship of Overweight With Other Health Issues (Metro Area, 2024)

Among Healthy Weight

Among Overweight/Not Obese

Among Obese



- Sources: 2024 PRC Community Health Survey, PRC, Inc. [Item 128]
- . Based on reported heights and weights, asked of all respondents

Key Informant Input: Nutrition, Physical Activity & Weight

Over half of key informants taking part in an online survey characterized Nutrition, Physical Activity & Weight as a "major problem" in the community.

Perceptions of Nutrition, Physical Activity, and Weight as a Problem in the Community (Metro Area, 2024)



Moderate Problem

Minor Problem

No Problem At All



Among those rating this issue as a "major problem," reasons related to the following:

Obesity

Increasing obesity rates and resulting medical consequences. - Health Provider

Overweight and obesity continues to affect a large portion of the population. - Health Provider

Look at the statistics regarding obesity in our country. - Health Provider

Obesity epidemic and high cost of effective new drugs. - Physician

Obesity and obesity-related illnesses. - Physician

Obesity rates, lack of safe walking across the city. - Health Provider



As long as Corporate America and big pharma profit from our population being fat and sick, there will never be the changes necessary in this country to actually help people find/live a healthy life. There's no such thing as nutrition with the products most can afford and have to buy/eat. Private, for-profit gyms and work out centers are overpriced and typically those places are stacked with so many people, most people will not want to continue to fight a crowd, including me, to try to get some exercise. And the for-profit places are good at locking people in for monthly membership that just about takes an act of Congress to get out of. They are banking on the idea that most people will continue paying the monthly membership, instead of "coming in" which they make you do to cancel, even if you can cancel. It's been a scam for many years, and I've been caught in it several times myself. — Health Provider

There is significant obesity and related diseases such as diabetes, hypertension, cancer that could be prevented with a better diet. The connection between how we feel and what we eat seems to be absent. – Health Provider

Lifestyle

People's weight is getting higher and higher, and their activity levels are decreasing. Many people do not know how to eat healthily and/or choose not to. People are becoming more sedentary. – Health Provider

Our community is largely overweight due to the normalization of obesity in our culture, lack of physical activity, and food insecurity for those with lower income. – Physician

A growing number of children and adults are presenting with obesity secondary to a sedentary lifestyle and poor nutrition. Gym memberships can be expensive and/or people are unsure how to use the equipment or don't know how to get started or what sort of exercise/nutrition program would be best suited for them. – Social Services Provider

Inactivity, fast foods, and obesity. - Physician

I personally don't feel we as a community are as active as we should or could be. Social media and video games keep us more stationary. Life for everyone is so busy and most families are on the go which means less nutritious meals and snacks. Joining a local gym is not an option for everyone due to cost, work, etc. – Social Services Provider

Our diet has been getting worse over the years. Fast food is cheaper and faster than food cooked at home. With the increasing use of social media, our young people are not getting as much physical activity. – Health Provider A culture of unhealthy food and inactivity. – Physician

People are exposed to fat intake foods, sugars, and sodium. Lack of exercise. Not enough fresh food. – Social Services Provider

Food insecurity, cultural influences, socioeconomic disparities, barriers to physical activity, cultural perceptions of body image, health care disparities, stress, and mental health. – Community Leader

Inactivity and poor access to healthy foods. - Health Provider

Access to Affordable Healthy Food

Needs to be attacked from a family and household level. Need easier access to nutritious food in schools and in neighborhoods. Need incentives in communities where this isn't prioritized. – Social Services Provider

Lack of healthy foods, cost of healthy foods. - Physician

Lack of access to healthy foods is affected due to costs and limited supply of healthy food options and ingredients. Processed foods are too abundant and cheap. – Advanced Practice Provider

Poor access to healthy foods. - Physician

Access to good and affordable nutrition and a lack of knowledge. - Physician

Not enough access to healthy foods. - Health Provider

Convenience and time are key contributors to decisions or ability to manage a healthy weight, whether it's the increasing price of food, or the time to prepare healthy meals yourself, it makes it very challenging for many to achieve. – Business Leader

Access to healthier foods and the price of healthier foods. It costs so much more to purchase healthier foods than non-healthy foods. – Business Leader

Awareness/Education

Limited education and promotion of physical activity. Not enough space for low-cost options. Limited multimodal active transit opportunities, such as bike lanes, walking paths, etc. Healthy food is perceived to be expensive. Overweight and obesity is normalized. – Public Health Representative

Education. - Social Services Provider

I think a lot of our issues tie back to educational attainment, age, and poverty. - Community Leader

Education, lack of integrated plans for medication and activity or nutrition. Expense of therapies and behavioral focused weight programs. Lack of a community focus on the issue. – Health Provider

Lack of understanding of physical activity and nutrition. Lack of resources, lack of integrated planning to build vibrant neighborhoods that promote physical activity, instead of needing to be so car centric in daily life. – Business Leader



Education, access to care, motivation, and financial limitations. - Physician

Education and access to healthy eating options for healthy living. - Community Leader

Income/Poverty

Lack of financial stability to make healthier lifestyle choices that support healthy food options and regular physical activity. – Community Leader

Poverty, unhealthy diet, lack of choices. - Business Leader

Lack of funds to buy the proper nutritious food, sedentary lifestyles, and too much screen time. – Social Services

Nutrition

Overabundant access to processed and unhealthy foods. There is not enough access to healthy, nutritious foods, like fruits and vegetables. School lunch programs serve lots of processed foods that are not healthy for our youth. – Social Services Provider

There are too many fast-food restaurants that are open 24 hours and have the ability to deliver food to their house. Physical activity is encouraged but not taken advantage of by all people in the community. – Health Provider

Access to Services

Lack of facilities and programs. - Community Leader

Built Environment

Places to safely exercise, as I specifically have patients who are afraid to walk in their neighborhood. Ability to afford healthy food. – Physician

Comorbidities

Diabetes, poverty, and homelessness. - Social Services Provider

Denial/Stigma

Stigma around weight. Omaha is not a walkable city, leading to less outdoor exercise. – Physician



SUBSTANCE USE

ABOUT DRUG & ALCOHOL USE

Substance use disorders can involve illicit drugs, prescription drugs, or alcohol. Opioid use disorders have become especially problematic in recent years. Substance use disorders are linked to many health problems, and overdoses can lead to emergency department visits and deaths.

Effective treatments for substance use disorders are available, but very few people get the treatment they need. Strategies to prevent substance use — especially in adolescents — and help people get treatment can reduce drug and alcohol misuse, related health problems, and deaths.

Healthy People 2030 (https://health.gov/healthypeople)

Alcohol Use

Age-Adjusted Alcohol-Induced Deaths


Between 2018 and 2020, the Metro Area reported an annual average age-adjusted mortality rate of 15.8 alcohol-induced deaths per 100,000 population.

BENCHMARK ► Well above the state and US rates.

TREND ► Increasing considerably over the past decade.

DISPARITY Highest in Douglas County and among Black individuals in the Metro Area.

Alcohol-Induced Deaths: Age-Adjusted Mortality (2018-2020 Annual Average Deaths per 100,000 Population)



Sources: • CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics. Data extracted June 2024.

US Department of Health and Human Services. Healthy People 2030. August 2020. http://www.healthypeople.gov


Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10).

• Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population.



Notes


Alcohol-Induced Deaths: Age-Adjusted Mortality by Race (2018-2020 Annual Average Deaths per 100,000 Population)



Sources: • CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics. Data extracted June 2024.

- Notes:
- US Department of Health and Human Services. Healthy People 2030. August 2020. http://www.healthypeople.gov
 Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10).
 Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population.

Alcohol-Induced Deaths: Age-Adjusted Mortality Trends (Annual Average Deaths per 100,000 Population)



	2011-2013	2012-2014	2013-2015	2014-2016	2015-2017	2016-2018	2017-2019	2018-2020
Metro Area	8.5	9.6	9.3	10.5	12.2	13.6	14.1	15.8
─ NE	7.9	8.5	8.2	8.4	9.0	10.1	10.8	12.0
——IA	7.8	8.4	8.8	9.1	9.2	9.2	9.2	9.9
<u></u> -US	9.9	10.3	10.6	10.8	10.8	10.9	11.1	11.9

Sources: • CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics. Data extracted June 2024.

US Department of Health and Human Services. Healthy People 2030. August 2020. http://www.healthypeople.gov
 Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10).

• Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population.



Notes:

Excessive Drinking

Excessive drinking includes heavy and/or binge drinkers:

- HEAVY DRINKING ➤ men reporting 2+ alcoholic drinks per day or women reporting 1+ alcoholic drink per day in the month preceding the interview.
- BINGE DRINKING ▶ men reporting 5+ alcoholic drinks or women reporting 4+ alcoholic drinks on any single occasion during the past month.

A total of 22.7% of area adults engage in excessive drinking (heavy and/or binge drinking).


BENCHMARK ► Higher than the Nebraska prevalence but well below the US.

TREND ► Decreasing significantly since 2018.

DISPARITY Highest in Douglas County. Reported more often among men, young adults, and those living in higher-income households. Note the low proportion of Asian respondents who engage in excessive drinking.

Excessive Drinking

Metro Area



Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Item 116]
• 2023 PRC National Health Survey, PRC, Inc.

 Behaivoral Risk Factor Surveillance System Survey Data. Atlanta, Georgia. United States Department of Health and Human Services, Centers for Disease Control and Prevention (CDC): 2022 Nebraska and Iowa data.


lotes:

 Asked of all respondents.

• Excessive drinking reflects the number of persons aged 18 years and over who drank more than two drinks per day on average (for men) or more than one drink per day on average (for women) OR who drank 5 or more drinks during a single occasion (for men) or 4 or more drinks during a single occasion (for women) during the past 30 days.



Excessive Drinking (Metro Area, 2024)



- Sources: 2024 PRC Community Health Survey, PRC, Inc. [Item 116]
 - Asked of all respondents.
 - Excessive drinking reflects the number of persons aged 18 years and over who drank more than two drinks per day on average (for men) or more than one drink per day on average (for women) OR who drank 5 or more drinks during a single occasion (for men) or 4 or more drinks during a single occasion (for women) during the past 30 days.

Drug Use

Age-Adjusted Unintentional Drug-Induced Deaths


Between 2018 and 2020, there was an annual average age-adjusted mortality rate of 8.9 unintentional drug-induced deaths per 100,000 population in the Metro Area.

BENCHMARK ► Higher than the Nebraska mortality rate but well below that of the US.

TREND The mortality rate has increased over the past decade, echoing the state trends (although well below the sharply increasing US trend).

DISPARITY Lower in Sarpy County. Slightly higher among Black residents than White residents.

Unintentional Drug-Related Deaths: Age-Adjusted Mortality (2018-2020 Annual Average Deaths per 100,000 Population)






Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10).

Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population



Unintentional Drug-Related Deaths: Age-Adjusted Mortality by Race

(2018-2020 Annual Average Deaths per 100,000 Population)




Notes:

- CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics. Data extracted June 2024.
 Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10).
- - Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population.

Unintentional Drug-Related Deaths: Age-Adjusted Mortality Trends

(Annual Average Deaths per 100,000 Population)



	2011-2013	2012-2014	2013-2015	2014-2016	2015-2017	2016-2018	2017-2019	2018-2020	
Metro A	rea 7.3	6.8	6.4	7.2	7.7	7.7	7.8	8.9	
NE	5.6	5.6	5.5	5.5	5.9	6.0	6.5	7.4	
—IA	6.4	6.8	7.4	7.8	8.5	8.3	8.6	9.4	
US	11.0	12.1	13.0	14.9	16.7	18.1	18.8	21.0	

Sources: • CDC WONDER Online Query System. Centers for Disease Control and Prevention, Epidemiology Program Office, Division of Public Health Surveillance and Informatics. Data extracted June 2024.

Deaths are coded using the Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10).
 Rates are per 100,000 population, age-adjusted to the 2000 US Standard Population.



Notes:

Opioids are a class of drugs used to treat pain. Examples presented to respondents include morphine, codeine, hydrocodone, oxycodone, methadone, and fentanyl. Common brand name opioids include Vicodin, Dilaudid, Percocet, OxyContin, and Demerol.

Use of Prescription Opioids


A total of 13.4% of Metro Area adults report using a prescription opioid drug in the past year.

TREND ► Decreasing significantly since 2018.

DISPARITY ► Highest in Northeast Omaha and Pottawattamie County. Reported more often among adults age 40 and older, those in low-income households, White adults, and Black or African American adults.

Used a Prescription Opioid in the Past Year

Metro Area




Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Item 41]

2023 PRC National Health Survey, PRC, Inc.

Notes:

 Asked of all respondents.

Used a Prescription Opioid in the Past Year (Metro Area, 2024)



Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Item 41]

2023 PRC National Health Survey, PRC, Inc.

Notes: • Asked of all respondents.



Alcohol & Drug Treatment


A total of 7.2% of Metro Area adults report that they have sought professional help for an alcohol or drug problem at some point in their lives.

TREND ▶ The prevalence has increased significantly since 2011.

DISPARITY ► Highest in Douglas County (especially Southeast Omaha).

Have Ever Sought Professional Help for an Alcohol/Drug-Related Problem

Metro Area




2023 PRC National Health Survey, PRC, Inc. Asked of all respondents.

Key Informant Input: Substance Use

The greatest share of key informants taking part in an online survey characterized Substance Use as a "major problem" in the community (followed closely by "moderate problem" responses).

Perceptions of Substance Use as a Problem in the Community (Metro Area, 2024)





Asked of all respondents.



Among those rating this issue as a "major problem," reasons related to the following:

Access to Care/Services

Not enough places to go for advice or counselling. Without family support, this issue becomes very difficult to deal with. – Social Services Provider

Very limited number of clinics and providers for managing serious substance use disorders. – Health Provider Residential treatment programs. – Physician

No facilities are available. - Business Leader

Receiving services in a timely manner. The cost of a substance abuse evaluation, shortage of staff and licensed professionals. – Social Services Provider

Lack of access and providers dedicated to this. Education and stigma. - Health Provider

There is a lack of substance abuse treatment options in the area. Many patients are committed to acute psychiatric care in the hospital setting which doesn't get the patient to the care they need, leading to long waits for treatment and preventing others from getting the services needed in the acute psychiatric unit. – Health Provider

It is difficult to find immediate openings for substance use treatment. There are often waiting lists and delays when attempting to go into a treatment facility. Often times the person relapses or chooses not to go if they have to long of a wait to get into a program. – Health Provider

Not enough 24/7 thirty-day detox, and then recovery programs that accept Nebraska Medicaid. – Health Provider Lack of insurance. – Health Provider

Accessing substance abuse treatment centers poses a formidable challenge, with individuals often encountering significant hurdles in securing admission and locating available facilities. As a result, many individuals find themselves discharged back into the same environments where their substance abuse issues originated, perpetuating a cycle of dependency and relapse. The scarcity of treatment options exacerbates this predicament, compounding the difficulties faced by those seeking help. Consequently, the barrier posed by insufficient treatment options continues to escalate, leaving many individuals grappling with addiction without the vital support they urgently require. – Health Provider

Affordable evaluations, open beds. No inpatient facilities in the community. The closest inpatient facility that accepts Iowa Medicaid is 45-60 minutes away. Transportation and accessibility are an issue too. – Health Provider

In addition to the financial hurdles, I think that one of the biggest barriers is getting an appointment or getting to the appointment. – Community Leader

Lack of reimbursement. Leads to a lack of resources and levels of care that are needed. Lack or compensation and pay wages for these disciplines to do this work. – Health Provider

Limited resources and limited capacity. - Community Leader

Access to care and treatment, capacity to get out of environments with high substance use. – Physician Sober living houses and the challenges of those going through drug court not being able to find employment that will allow for the meetings, UA testing, and programming requests. – Social Services Provider

I think the need is exceeding the availability. - Physician

Denial/Stigma

Stigma and insurance. - Physician

Stigma. – Business Leader

Stigma for accessing care. - Health Provider

Stigmas and fear to admit they need help. - Community Leader

Consumer buy-in that they have a problem and need help to combat it. Transportation to services. Accountability to the substance user. They can't see a path forward. – Community Leader

Stigma and the financial capability to afford treatment options. - Community Leader

Lack of Providers

Once again, there is a lack of substance abuse providers. We have a workforce shortage of therapists to address substance misuse in our communities. Many state funded programs cannot afford to pay people to be fully staffed, which reduces accesses and increases wait time. — Social Services Provider

LACK OF EVIDENCE BASED PROVIDERS, stigma, STIGMA!!!!!! People refuse to provide evidence-based treatment with medications. Literally, doctors will refuse to prescribe medication to patients that is needed for their disease. It is completely abhorrent that this is happening, and it is discriminatory to patients. No doctor would refuse to prescribe a medication for their blood pressure- but they refuse to prescribe it for another chronic health condition, their substance use disorder. – Physician



Shortage of trained clinicians and inadequate reimbursement rates. Lack of long-term residential treatment options in our community available for free, or on a sliding-fee scale. – Social Services Provider

Need more substance abuse providers - very difficult to recruit and retain therapists. We need more options and better community understanding of how to access care and where. Should be noted that Providers do exist in the community, there seems to be some access, however, individuals struggle with motivation to follow-through with evaluation and treatment. And substance abuse still seems to be an accepted norm in society. – Health Provider

Affordable Care/Services

Cost and availability. - Social Services Provider

Affordable facilities to get treatment, both intensive outpatient and inpatient treatment. Lack of insurance coverage for substance use treatment. – Social Services Provider

Unknown to me, but cost and specific programs are barriers. - Physician

Awareness/Education

Lack of knowledge of available resources. - Physician

Lack of information about them and too few resources. Unable to afford care. Culture that promotes and glorifies drug use and selling drugs. – Physician

Not adequate access to education and prevention. - Health Provider

Income/Poverty

Poverty, family history and mental health. – Social Services Provider

Stigma

Willingness of those in need and support. - Business Leader

The desire to seek out treatment. - Health Provider

Comorbidities

Mental health, trauma, depression, anxiety, and social isolation. – Health Provider

Cultural Mistrust

Cultural mistrust, lack of culturally competent care, financial barriers, limited availability of services, language barriers and legal concerns. – Community Leader

Diagnosis/Treatment

Good urgent withdrawal treatment. Good urgent long term treatment options. – Physician

Easy Access

Represented by large amounts of availability. Seeing a houselessness increase over the last few years. – Social Services Provider

Hopelessness

People have no hope, so they don't care about using drugs. Difficult getting people in. – Health Provider



Most Problematic Substances

Key informants (who rated this as a "major problem") clearly identified **alcohol** as causing the most problems in the community, followed distantly by **methamphetamine/other amphetamines**, **heroin or other opioids**, **marijuana**, **prescription medications**, and **over-the-counter medications**.

SUBSTANCES VIEWED AS MOST PROBLEMATIC IN THE COMMUNITY

(Among Key Informants Rating Substance Use as a "Major Problem")

ALCOHOL	73.5%
METHAMPHETAMINE OR OTHER AMPHETAMINES	12.2%
HEROIN OR OTHER OPIOIDS	6.1%
MARIJUANA	4.1%
PRESCRIPTION MEDICATIONS	2.0%
OVER-THE-COUNTER MEDICATIONS	2.0%



TOBACCO USE

ABOUT TOBACCO USE

Most deaths and diseases from tobacco use in the United States are caused by cigarettes. Smoking harms nearly every organ in the body and increases the risk of heart disease, stroke, lung diseases, and many types of cancer. Although smoking is widespread, it's more common in certain groups, including men, American Indians/Alaska Natives, people with behavioral health conditions, LGBT people, and people with lower incomes and education levels.


Several evidence-based strategies can help prevent and reduce tobacco use and exposure to secondhand smoke. These include smoke-free policies, price increases, and health education campaigns that target large audiences. Methods like counseling and medication can also help people stop using tobacco.


- Healthy People 2030 (https://health.gov/healthypeople)

Cigarette Smoking

Prevalence of Cigarette Smoking

A total of 14.8% of Metro Area adults currently smoke cigarettes, either regularly (every day) or occasionally (on some days).







Note the following findings related to cigarette smoking prevalence in the Metro Area.

BENCHMARK Higher than the Nebraska findings but lower than the US. Fails to satisfy the Healthy People 2030 objective.


TREND ▶ Though increasing since 2018, the percentage remains below 2015 findings.

DISPARITY > Highest in eastern Omaha and Pottawattamie County. Reported more often among men, adults under age 65, those in low-income households, Black or African American residents, those of Diverse Races, and LGBTQ+ adults.

Current Smokers

Healthy People 2030 = 6.1% or Lower

Metro Area



Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Item 34]
• Behavioral Risk Factor Surveillance System Survey Data. Atlanta, Georgia. United States Department of Health and Human Services, Centers for Disease Control and Prevention (CDC): 2022 Nebraska and Iowa data.


2023 PRC National Health Survey, PRC, Inc.

US Department of Health and Human Services. Healthy People 2030. August 2020. http://www.healthypeople.gov

Asked of all respondents.
 Includes regular and occasional smokers (those who smoke cigarettes every day or on some days).

Current Smoke Cigarettes (Metro Area, 2024)

Healthy People 2030 = 6.1% or Lower





US Department of Health and Human Services. Healthy People 2030. August 2020. http://www.healthypeople.gov

Notes Asked of all respondents.

Includes regular and occasion smokers (every day and some days)



Environmental Tobacco Smoke

Among all surveyed households in the Metro Area, 11.4% report that someone has smoked cigarettes, cigars, or pipes anywhere in their home an average of four or more times per week over the past month.


BENCHMARK ► Lower than the national figure.

TREND ▶ Decreasing significantly since 2011.

DISPARITY Highest in Douglas County (especially east of 72nd Street) and Pottawattamie County.

Member of Household Smokes at Home

Metro Area



Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Items 35, 114]

2023 PRC National Health Survey, PRC, Inc.

Asked of all respondents.

• "Smokes at home" refers to someone smoking cigarettes, cigars, or a pipe in the home an average of four or more times per week in the past month.

Smoking Cessation


Just less than one-half of adults who regularly smoke cigarettes (49.3%) went without smoking for one day or longer in the past year because they were trying to quit smoking.

BENCHMARK ► Fails to satisfy the Healthy People 2030 objective.

Have Stopped Smoking for One Day or Longer in the Past Year (Everyday Smokers)

Healthy People 2030 = 65.7% or Higher

Metro Area





2023 PRC National Health Survey, PRC, Inc.


- Behavioral Risk Factor Surveillance System Survey Data. Atlanta, Georgia. United States Department of Health and Human Services, Centers for Disease Control and Prevention (CDC): 2018 Nebraska and Iowa data.
- US Department of Health and Human Services. Healthy People 2030. August 2020. http://www.healthypeople.gov

Asked of respondents who smoke cigarettes every day.



Use of Vaping Products

Most Metro Area adults do not use electronic vaping products.



Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Item 36]
Notes: • Asked of all respondents.

However, 11.3% currently use electronic vaping products either regularly (every day) or occasionally (on some days).


BENCHMARK ▶ Higher than the state percentages but below that found nationally.

TREND ► Increasing significantly from 2011 findings.

DISPARITY ► Much higher in eastern Omaha. The prevalence decreases with age and household income level and is reported more often among adults of Diverse Races and LGBTQ+ respondents.

Currently Use Vaping Products (Every Day or on Some Days)

Metro Area





Behavioral Risk Factor Surveillance System Survey Data. Atlanta, Georgia. United States Department of Health and Human Services, Centers for Disease Control
and Prevention (CDC): 2022 Nebraska and Iowa data.


2023 PRC National Health Survey, PRC, Inc.

Notes: • Asked of all respondents.

• Includes regular and occasional users (those who smoke e-cigarettes every day or on some days).



Currently Use Vaping Products (Metro Area, 2024)



- Sources: 2024 PRC Community Health Survey, PRC, Inc. [Item 36]
 - Asked of all respondents.
 - Includes regular and occasional users (those who smoke e-cigarettes every day or on some days).

Key Informant Input: Tobacco Use

The greatest share of key informants taking part in an online survey characterized Tobacco Use as a "moderate problem" in the community.

Perceptions of Tobacco Use as a Problem in the Community (Metro Area, 2024)



 PRC Online Key Informant Survey, PRC, Inc. Asked of all respondents.

Among those rating this issue as a "major problem," reasons related to the following:

Incidence/Prevalence

So many people still smoke, even with the negative health effects. - Social Services Provider Continued significant tobacco use in the community with medical problems secondary to this. - Physician High rates of smoking. - Health Provider

When I moved to this community seven years ago, I was struck by how many people I saw smoking. I think this is tied to age, educational attainment, and poverty. - Community Leader

Tobacco and vaping are a big problem in our community because there are many people who use tobacco or nicotine and cannot quit using it. We continue to see vape and tobacco shops open in our community, which increases access to the products (including access to youth), but when the stores keep opening, you know that use is a problem, simply because the market for the tobacco / nicotine products are there. If there were not a market, they would not open up new shops to claim some of the market share. - Social Services Provider



Far too many people smoke or use to smoke, making themselves vulnerable to future diseases. – Social Services Provider

Easy Access

Easily accessible for minors to obtain with new tobacco products on the market. – Health Provider

Easy access to tobacco stores. - Health Provider

Easy to access. - Social Services Provider

Co-Occurrences

Often concomitant with other substance use. - Physician

It leads to heart problems, stroke, weight gain or loss, dental diseases, unemployment, and violence and mental illness. – Business Leader

E-Cigarettes

Vaping among young people. - Social Services Provider

I think tobacco use may be on the decline, but the increase in vaping is concerning. - Health Provider

Income/Poverty

People say they can't pay their bills or afford necessities such as food, and yet they smoke a pack or more of cigarettes a day. – Health Provider

Tobacco is primarily used by the poor community. - Health Provider

Addiction

It's an expensive addictive drug that is the first item of necessity. Often monetary resources first allocation. – Community Leader

Impact on Caregivers/Families

Tobacco use is a problem because the addiction affects other people that reside with the smoker, causing long term effects. – Health Provider



SEXUAL HEALTH

ABOUT HIV & SEXUALLY TRANSMITTED INFECTIONS

Although many sexually transmitted infections (STIs) are preventable, there are more than 20 million estimated new cases in the United States each year — and rates are increasing. In addition, more than 1.2 million people in the United States are living with HIV (human immunodeficiency virus).

Adolescents, young adults, and men who have sex with men are at higher risk of getting STIs. And people who have an STI may be at higher risk of getting HIV. Promoting behaviors like condom use can help prevent STIs.

Strategies to increase screening and testing for STIs can assess people's risk of getting an STI and help people with STIs get treatment, improving their health and making it less likely that STIs will spread to others. Getting treated for an STI other than HIV can help prevent complications from the STI but doesn't prevent HIV from spreading.

Healthy People 2030 (https://health.gov/healthypeople)


HIV

In 2021, the Metro Area reported a prevalence of 76.3 HIV cases per 100,000 population.

BENCHMARK ► Well below the state rates; exceedingly lower than the national rate.

DISPARITY ► Locally highest in Douglas and (especially) Pottawattamie counties. The prevalence rate is much higher among Black residents than White or Hispanic residents.

HIV Prevalence (Prevalence Rate of HIV per 100,000 Population, 2021)




Sources: • Centers for Disease Control and Prevention, National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention.

Center for Applied Research and Engagement Systems (CARES), University of Missouri Extension. Retrieved June 2024 via SparkMap (sparkmap.org).



HIV Prevalence by Race/Ethnicity (Rate per 100,000 Population, 2021)



- Centers for Disease Control and Prevention, National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention.
- Center for Applied Research and Engagement Systems (CARES), University of Missouri Extension. Retrieved June 2024 via SparkMap (sparkmap.org).

Race categories reflect individuals without Hispanic origin.

Sexually Transmitted Infections (STIs)

Chlamydia & Gonorrhea

In 2021, the chlamydia incidence rate in the Metro Area was 572.1 cases per 100,000 population.

BENCHMARK ► Worse than the Nebraska rate.


DISPARITY ► Highest in Douglas County.

The Metro Area gonorrhea incidence rate in 2021 was 238.8 cases per 100,000 population.

BENCHMARK ► Well above both state incidence rates.

DISPARITY ► Highest in Douglas County.

Chlamydia & Gonorrhea Incidence (Incidence Rate per 100,000 Population, 2021)







- Centers for Disease Control and Prevention, National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention.
 - Center for Applied Research and Engagement Systems (CARES), University of Missouri Extension. Retrieved June 2024 via SparkMap (sparkmap.org)



Key Informant Input: Sexual Health

A plurality of key informants taking part in an online survey characterized Sexual Health as a "moderate problem" in the community.





Among those rating this issue as a "major problem," reasons related to the following:

Incidence/Prevalence

· Asked of all respondents

Data from the Health Department shows STI rates are far above national and state averages. – Business Leader High rates of STDs. – Community Leader

Large number of STDs in collage age and senior citizens in the community. - Social Services Provider

I believe we have a higher rate of STDs than the state rate in Iowa. Drug use and being impaired contributes to STDs because of sexual assault and the ability to make good decisions while under the influence of all substances. – Social Services Provider

Rates are still high. There is shame associated with practicing healthy sexual behaviors and or getting screened. – Social Services Provider

We have been above the national average for years. - Community Leader

Reports provided by local Emergency Room staff and the local health department. - Social Services Provider

Elevated rates of CVD, syphilis, and other STIs in the community. - Health Provider

Very high rates of chlamydia. - Physician

Omaha is known for their high rate of STDs and being higher than the national average. – Social Services Provider

Increasing HIV, syphilis, high STI rates, ectopic pregnancies. - Physician

My understanding is Douglas County has a large population of STIs. - Health Provider

Rates of STIs are increasing. - Physician

High rates chlamydia, GC, and syphilis. - Physician

High rate of STIs, including GC, chlamydia, and syphilis. - Physician

Douglas County has one of the highest STD rates, per capita, in the nation. - Physician

High rates of STIs, in particular syphilis. Lack of basic comprehensive sex education in the public school system. – Community Leader

Rising number of STD cases in Omaha. Lack of access to Planned Parenthood or other testing facilities. – Social Services Provider

Awareness/Education

Lack of awareness, hygiene, and low morals. - Business Leader

Statistics and lack of information, poverty, mental health, and addiction. - Social Services Provider

Omaha has been horrible for years. Maybe the lack of sexual health education. - Community Leader

Education, lack of access, stigma of access and treatment, lack of centralized anonymous screening and treatment, lack of on-demand screening and treatment possibilities. – Health Provider

STD prevention is not talked about enough with young people. Not talking about it does not keep it from happening, but rather without any safety precautions. – Health Provider



Access to Care

Politicization of sexual & reproductive health and continued changes in laws targeted by the State Legislature have a chilling effect on those seeking care or contemplating seeking care. Gender-affirming care is restricted to minors under age 19. It is often weeks before community clinics have available slots of HIV/STI testing, and this can radically impact the ability to access post-exposure prophylactic treatment scheduling. Certain STIs are increasing in prevalence (syphilis) or becoming treatment resistant (gonorrhea), primarily due to the widespread use of pre-exposure prophylactic (which should be encouraged and more accessible) but sometimes standard STI panels will omit one or more STIs. A lack of inclusive and updated medical forms at primary care physicians (PCP) don't ask sexual practices/orientation, leaving physicians unable to adequately treat or recommend preventative strategies or testing regimes. Lack of DoxyPEP usage. – Community Leader

Income/Poverty

Compounding social determinants of health that affect access and prevention, such as poverty. – Health Provider


Funding

Lack of ongoing funding to continuously educate through rigorous campaigns targeting those most at risk. – Community Leader

Culture

Kids are pushed to grow up. Society rushes kids to become more independent too soon. – Social Services Provider





ACCESS TO HEALTH CARE


HEALTH INSURANCE COVERAGE

Type of Health Care Coverage

A total of 64.1% of Metro Area adults age 18 to 64 report having health care coverage through private insurance. Another 20.9% report coverage through a government-sponsored program (e.g., Medicaid, Medicare, military benefits).

Survey respondents were asked a series of questions to determine their health care insurance coverage, if any, from either private or government-sponsored sources.

Health Care Insurance Coverage (Adults Age 18-64; Metro Area, 2024)



Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Item 117]

Reflects respondents age 18 to 64.

Lack of Health Insurance Coverage

Among adults age 18 to 64, 7.8% report having no insurance coverage for health care expenses.

BENCHMARK ▶ Lower than the Nebraska response.

TREND ▶ The prevalence has decreased significantly since 2011.

DISPARITY Highest in Douglas County (especially Southeast Omaha). Lack of coverage is reported more often among men, young adults, those in low-income households, Hispanic respondents, Black or African American respondents, and those who identify as LGBTQ+.

Here, lack of health insurance coverage reflects respondents age 18 to 64 (thus, excluding the Medicare population) who have no type of insurance coverage for health care services neither private insurance nor governmentsponsored plans (e.g., Medicaid).




Lack of Health Care Insurance Coverage

(Adults Age 18-64)

Healthy People 2030 = 7.6% or Lower

Metro Area



Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Item 117]


- Behavioral Risk Factor Surveillance System Survey Data. Atlanta, Georgia. United States Department of Health and Human Services, Centers for Disease Control and Prevention (CDC): 2022 Nebraska and Iowa data.

 2023 PRC National Health Survey, PRC, Inc.
- US Department of Health and Human Services. Healthy People 2030. August 2020. http://www.healthypeople.gov

Notes: • Asked of all respondents under the age of 65.

Lack of Health Care Insurance Coverage (Adults Age 18-64; Metro Area, 2024)

Healthy People 2030 = 7.6% or Lower



Sources:

2024 PRC Community Health Survey, PRC, Inc. [Item 117]
 US Department of Health and Human Services. Healthy People 2030. August 2020. http://www.healthypeople.gov

• Asked of all respondents under the age of 65.



DIFFICULTIES ACCESSING HEALTH CARE

ABOUT HEALTH CARE ACCESS

Many people in the United States don't get the health care services they need. ...People without insurance are less likely to have a primary care provider, and they may not be able to afford the health care services and medications they need. Strategies to increase insurance coverage rates are critical for making sure more people get important health care services, like preventive care and treatment for chronic illnesses.

Sometimes people don't get recommended health care services, like cancer screenings, because they don't have a primary care provider. Other times, it's because they live too far away from health care providers who offer them. Interventions to increase access to health care professionals and improve communication — in person or remotely — can help more people get the care they need.

Healthy People 2030 (https://health.gov/healthypeople)

Difficulties Accessing Services

A total of 42.2% of Metro Area adults report some type of difficulty or delay in obtaining health care services in the past year.


BENCHMARK ► Well below the national prevalence.

TREND Increasing significantly from early survey administrations.

DISPARITY ► The prevalence decreases with age and household income level and is reported more often among women, Hispanic adults, and LGBTQ+ respondents.

Experienced Difficulties or Delays of Some Kind in Receiving Needed Health Care in the Past Year

Metro Area



- Sources: 2024 PRC Community Health Survey, PRC, Inc. [Item 119]
 - 2023 PRC National Health Survey, PRC, Inc.


Notes: • Asked of all respondents.

Percentage represents the proportion of respondents experiencing one or more barriers to accessing health care in the past 12 months

This indicator reflects the percentage of the total population experiencing problems accessing health care in the past year, regardless of whether they needed or sought care. It is based on reports of the barriers outlined in the following section.



Experienced Difficulties or Delays of Some Kind in Receiving Needed Health Care in the Past Year (Metro Area, 2024)



Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Item 119] Notes:

Asked of all respondents

Percentage represents the proportion of respondents experiencing one or more barriers to accessing health care in the past 12 months.

Barriers to Health Care Access

Of the tested barriers, appointment availability impacted the greatest share of Metro Area adults.

BENCHMARK Each of the barriers illustrated fared better than its national counterpart.


TREND ▶ Each of the barriers below has increased significantly from 2011 findings.

DISPARITY Residents of Douglas County (especially east of 72nd Street) are more likely than others to experience many of these barriers to medical care (not shown).

Barriers to Access Have Prevented Medical Care in the Past Year (Metro Area)

■2011 ■2015 ■2018 ■2021 ■2024 ■US

In addition, 17.0% of adults have skipped doses or stretched a needed prescription in the past year in order to save costs.





Notes Asked of all respondents.

access prevented them from seeing a physician or obtaining a needed prescription in the past year. Again, these percentages reflect the total population, regardless of whether medical care was needed or sought.

To better understand health care access barriers, survey

participants were asked whether any of seven


types of barriers to



Key Informant Input: Access to Health Care Services

Key informants taking part in an online survey most often characterized *Access to Health Care Services* as a "moderate problem" in the community.

Perceptions of Access to Health Care Services as a Problem in the Community (Metro Area, 2024)



Among those rating this issue as a "major problem," reasons related to the following:

Access to Care/Services

Asked of all respondents.

Options for insurance, which limit health care professionals, and the distance to travel for some services. – Health Provider

Insurance, transportation, and treatment of clients. - Community Leader

Wait times for specialty providers. - Health Provider

Services for older adults. - Social Services Provider

Primarily behavioral health care and dental care here is Council Bluffs. - Health Provider

Availability during hours people can access transportation. Access to therapeutic interventions. Insurance. Geographic access and health care deserts. – Physician

Having providers who take new patients, having the availability to get in, being able to afford it, more access on bus lines. – Health Provider

For individual's dependent on federally qualified health centers for their health care, securing an appointment on the lowa side can often feel like an insurmountable task. – Health Provider

Affordable Care/Services

Cost. Many individuals cannot afford to see a health care provider, and then pay for prescriptions. Affordable health care options should be available in every part of the community. – Physician

Higher premium, very complicated rules, and guidelines, out of pocket co-pays, illiteracy, and difficult portal due to technology. – Business Leader

There are not enough services for low-income patients. - Health Provider

Cost, accessibility, especially mental health, getting an appointment takes six to nine months, and many times it is not covered. – Health Provider

Affordable health care services, access to primary care services, timely access to specialty care services. People are concerned about making ends meet with the various health related social needs that sometimes accessing preventative care services or routine care checks are not attainable. – Community Leader

Lack of Providers

Large driver of access is the low number of primary care physicians resulting in use of emergency or urgent services rather than reliable preventative care. SDOH including transportation access is a significant barrier. For hourly wage earners and many other salaried workers, they have no access to leave to allow for attending health related appointments. Another barrier is underinsured with large deductibles and choosing to avoid or delay care due to large out of pocket cost despite insurance. – Physician



There are not enough care providers and many care providers, including doctors and dentists and even chiropractors are working a 4-day week. That means having to schedule out months and wait months to see a provider. Personally, I have care providers that I appreciate very much, but even with that relationship of many years, if I have an issue, I have to go to an urgent care clinic or wait 1-3 months to see my own providers. I go every 6 months for a dental checkup and teeth cleaning. Twice last year, the office was closed on the day I was scheduled. Once for snow, and it didn't snow and the other, the dentist, for whatever reason, needed the day off. At face value, that would be tolerable, but you go to the back of the line for rescheduling and the 6-month checkup ends up being 9 months. A lot can happen in 1-3 months waiting to see a care provider and I fear with many people, that IuII in time can mean the difference between life or death. — Health Provider

Not enough providers, especially in certain specialty areas, such as psychiatry and neurology. – Physician There are not enough providers, and the rural areas are significantly lacking resources. – Social Services Provider

Income/Poverty

Poverty, homelessness, access to affordable housing, and an increase in mental health and physical disability conditions. – Social Services Provider

Finances, even with insurance, the costs of health care influence the decisions of the majority of patients. Compounding the problem are the many patients with no insurance or are underinsured. – Advanced Practice Provider

Transportation

Transportation access to connect rural people to resources, resulting in delayed or neglected treatment. – Business Leader

Transportation with or without insurance is very difficult to access and very inaccessible at all hours of the day and night. Insurance is a barrier, not a resource anymore. It creates too many obstacles. – Health Provider

Awareness/Education

Information can be hard to find. Trust between community and health professionals is broken or non-existent. Not all patients are treated equally. Many health providers cannot relate or do not try to relate to patients. Access to some care is difficult because of poor public transportation. – Social Services Provider

Lifestyle

Obesity is driven by lack of physical activity, nutrition, food access and the built environment. Traffic fatalities and serious injuries, particularly for vulnerable roadway users (pedestrians/cyclists/motorcyclists). Mental health and substance abuse disorders impacting suicide and traffic fatalities and serious injuries. – Community Leader

Insurance Issues

The main access in accessing health care services is lack of insurance. The state government helped to mitigate this through expanding Medicaid. As of 2021 9.8% of Nebraskans under age 65 did not have insurance which is a higher percentage than Iowa, Minnesota, or North Dakota. Health systems such as Nebraska Medicine could make the process for getting financial assistance simpler which would allow more uninsured people to access care. – Physician

Overuse of Emergency Room

Too many patients using the emergency departments as their primary care providers, and not getting the follow through that is needed because that is not the role of the emergency department providers. – Health Provider

Value-Based Care Model

A move towards value-based care would incentivize health systems to improve quality, efficiency and outcomes measurements, while lowering the cost of care by addressing the social determinants of care and behavioral health. – Physician



PRIMARY CARE SERVICES

ABOUT PREVENTIVE CARE

Getting preventive care reduces the risk for diseases, disabilities, and death — yet millions of people in the United States don't get recommended preventive health care services.

Children need regular well-child and dental visits to track their development and find health problems early, when they're usually easier to treat. Services like screenings, dental check-ups, and vaccinations are key to keeping people of all ages healthy. But for a variety of reasons, many people don't get the preventive care they need. Barriers include cost, not having a primary care provider, living too far from providers, and lack of awareness about recommended preventive services.

Teaching people about the importance of preventive care is key to making sure more people get recommended services. Law and policy changes can also help more people access these critical services.


- Healthy People 2030 (https://health.gov/healthypeople)

Access to Primary Care

There are currently 1,022 primary care physicians in the Metro Area, translating to a rate of 114.1 primary care physicians per 100,000 population.

DISPARITY ► The proportion is lowest in Cass County.

Access to Primary Care (Number of Primary Care Physicians per 100,000 Population, 2024)



Sources: • Centers for Medicare and Medicaid Services, National Plan and Provider Enumeration System (NPPES).

Center for Applied Research and Engagement Systems (CARES), University of Missouri Extension. Retrieved June 2024 via SparkMap (sparkmap.org).
 Notes:
 Doctors classified as "nomany care physicians" by the AMA include general family medicine MDs and DOs general practice MDs and DOs general intensions.

Doctors classified as "primary care physicians" by the AMA include general family medicine MDs and DOs, general practice MDs and DOs, general internal
medicine MDs, and general pediatrics MDs. Physicians age 75 and over and physicians practicing sub-specialties within the listed specialties are excluded.

Note that this indicator takes into account *only* primary care physicians. It does <u>not</u> reflect primary care access available through advanced practice providers, such as physician assistants or nurse practitioners.



Specific Source of Ongoing Care

A total of 76.6% of Metro Area adults were determined to have a specific source of ongoing medical care.


BENCHMARK ► Higher than the US prevalence but fails to satisfy the Healthy People 2030 objective.

TREND ► Increasing significantly since 2018.

DISPARITY Lowest in Northeast Omaha.

Have a Specific Source of Ongoing Medical Care

Healthy People 2030 = 84.0% or Higher



- Sources: 2024 PRC Community Health Survey, PRC, Inc. [Item 118]
 - 2023 PRC National Health Survey, PRC, Inc.
 - US Department of Health and Human Services. Healthy People 2030. August 2020. http://www.healthypeople.gov

Notes:

Asked of all respondents.

Utilization of Primary Care Services

Adults

A total of 71.5% of Metro Area adults visited a physician for a routine checkup in the past year.

BENCHMARK ► Well below the state percentages but higher than the US percentage.

TREND ► Marks a statistically significant increase since 2011.

DISPARITY Lowest in Southeast Omaha. Reported less often among men, young adults, those in low-income households, Hispanic adults, and LGBTQ+ residents.



Having a specific source

of ongoing care includes having a doctor's office,

public health clinic, community health center,

urgent care or walk-in clinic, military/VA facility, or some other kind of

place to go if one is sick or needs advice about his or her health. This resource is crucial to the

centered medical homes"

A hospital emergency room is not considered a

specific source of


ongoing care in this

concept of "patient-

(PCMH).

instance.

Have Visited a Physician for a Checkup in the Past Year




Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Item 16]

- Behavioral Risk Factor Surveillance System Survey Data. Atlanta, Georgia. United States Department of Health and Human Services, Centers for Disease Control
 and Prevention (CDC): 2022 Nebraska and Iowa data.
- 2023 PRC National Health Survey, PRC, Inc.

Asked of all respondents.

Have Visited a Physician for a Checkup in the Past Year (Metro Area, 2024)



Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Item 16]
Notes: • Asked of all respondents.



EMERGENCY ROOM UTILIZATION

A total of 11.8% of Metro Area adults have gone to a hospital emergency room more than once in the past year about their own health.


BENCHMARK ► Lower than the national figure.

TREND ▶ The prevalence has more than doubled since 2011.

DISPARITY Highest in Northeast Omaha. Reported more often among young adults, those in lowincome households, Black or African American respondents, and LGBTQ+ adults.

Have Used a Hospital Emergency Room More Than Once in the Past Year


Metro Area



Sources: • 2024 PRC Community Health Survey, PRC, Inc. [Item 19] • 2023 PRC National Health Survey, PRC, Inc.

Asked of all respondents.

Have Used a Hospital Emergency Room More Than Once in the Past Year (Metro Area, 2024)





Asked of all respondents.



ORAL HEALTH

ABOUT ORAL HEALTH

Tooth decay is the most common chronic disease in children and adults in the United States. ...Regular preventive dental care can catch problems early, when they're usually easier to treat. But many people don't get the care they need, often because they can't afford it. Untreated oral health problems can cause pain and disability and are linked to other diseases.

Strategies to help people access dental services can help prevent problems like tooth decay, gum disease, and tooth loss. Individual-level interventions like topical fluorides and community-level interventions like community water fluoridation can also help improve oral health. In addition, teaching people how to take care of their teeth and gums can help prevent oral health problems.

Healthy People 2030 (https://health.gov/healthypeople)

Dental Care

Adults

Two in three (67.4%) Metro Area adults have visited a dentist or dental clinic (for any reason) in the past year.


BENCHMARK ► Well above the national prevalence.

TREND ► Except for the low finding in 2021, decreasing significantly from earlier survey administrations.

DISPARITY Lowest in eastern Omaha and Pottawattamie County. Reported less often among young adults, those in low-income households, Hispanic adults, Black or African American adults, those of Diverse Races, and those who identify as LGBTQ+.

Have Visited a Dentist or Dental Clinic Within the Past Year

Healthy People 2030 = 45.0% or Higher





- Behavioral Risk Factor Surveillance System Survey Data. Atlanta, Georgia. United States Department of Health and Human Services, Centers for Disease Control
 and Prevention (CDC): 2022 Nebraska and Iowa data.
- 2023 PRC National Health Survey, PRC, Inc.
- US Department of Health and Human Services. Healthy People 2030. August 2020. http://www.healthypeople.gov


Notes:

 Asked of all respondents.



Have Visited a Dentist or Dental Clinic Within the Past Year

(Metro Area, 2024) Healthy People 2030 = 45.0% or Higher




Sources:

2024 PRC Community Health Survey, PRC, Inc. [Item 17]
 US Department of Health and Human Services. Healthy People 2030. August 2020. http://www.healthypeople.gov

Asked of all respondents.

Key Informant Input: Oral Health

Key informants taking part in an online survey most often characterized Oral Health as a "moderate problem" in the community.





 PRC Online Key Informant Survey, PRC, Inc. Asked of all respondents.

Among those rating this issue as a "major problem," reasons related to the following:

Access for Medicare/Medicaid Patients

For families with Medicaid coverage, there are virtually no dentists in Iowa accepting new patients. I had a parent today tell me they are on a waiting list for a dental appointment, but whenever the appointment gets close, the office calls and cancels. - Social Services Provider

No providers accepting patients with Medicaid. - Social Services Provider

It is very difficult to find dentists willing to accept Medicaid as an insurance. There are few providers and the wait time to be seen in the office is often months or more. It is very difficult to find emergency dental care. - Health

The majority of dentists in the community no longer accept lowa Medicaid. - Health Provider

Lack of services available for Medicaid or uninsured clients, especially youth. - Health Provider

There are not enough dentists to take Medicaid clients. The wait lists are horrible. Research shows poor dental care results in chronic health conditions. - Health Provider



Access to Care/Services

Dental care access creates long-term health complications. - Physician

Most of my patients lack teeth or lack healthy teeth and lack access to quality and timely dental care and oral surgical care. – Physician

Lack of dental health facilities. - Community Leader

It is very difficult for my patients to find a dentist or dental home, and some have severe tooth decay, affecting their health. – Physician

Lack of access. - Health Provider

Lack of health care access. - Social Services Provider

Affordable Care/Services

Because no one can afford it, so if you don't have insurance or cash, dental hygiene is the first to go. – Health Provider

It is not available to low resource children and families. - Social Services Provider

It is not affordable. - Social Services Provider

Very expensive, lack of insurance, not a lot of options. - Business Leader

Too costly to get dental care. - Social Services Provider

Access to Care for Uninsured/Underinsured

Very limited access to dental care for those without insurance, or with financial barriers. Many go without treatment. – Physician

Lack of affordable dental insurance and services. - Advanced Practice Provider


Environmental Contributors

No fluorinated water in rural communities. Access to dental care and or affordable providers. Delayed dental care for children. – Business Leader

Impact on Quality of Life

Poverty, mental health, and addiction. - Social Services Provider






LOCAL RESOURCES

HEALTH CARE RESOURCES & FACILITIES

Federally Qualified Health Centers (FQHCs)

The following map details Federally Qualified Health Centers (FQHCs) within the Metro Area as of December 2023.





Resources Available to Address Significant Health Needs

The following represent potential measures and resources (such as programs, organizations, and facilities in the community) identified by key informants as available to address the significant health needs identified in this report. This list only reflects input from participants in the Online Key Informant Survey and should not be considered to be exhaustive nor an all-inclusive list of available resources.

Access to Health Care Services

All Care Community Health Center

Bluffs Taxi

Caring for Our Communities

Casino Cab

Charles Drew Health Center

CHI Health Center

CHI Health Immanuel

CHI Health Mercy

City of Omaha Vision Zero Plan

Community Alliance

Community Health Centers

Community Health Workers

Connections Area Agency on Aging

Council Bluffs Transit

Creighton University

Doctor's Offices

Douglas County

Douglas County Detox Center

Douglas County Health Department

Faith Based Nursing Resources

Federal Programs

Federally Qualified Health Centers

Firefly Mobile Wellness Unit

Health Care Facilities

Health Fairs

Health Systems

Heartland Family Services

Healing Gift Free Clinic (HGFC)

Hospitals

Insurance Company

Jennie Edmundson Hospital

Lutheran Family Services

MAPA Safe Streets and Roads

Marketplace Insurance

Methodist Community Health Clinic (MCHC)

Medicare/Medicaid

Mental Health Respite Programs

Methodist

Metro Area Transit

Mobile Care Units

Nebraska Medicine

Nonprofits

North Omaha Area Health

One World Community Health Center

Secret Heart

Southwest Iowa Transit System

University of Nebraska Medical Center

VA

Cancer

A Time to Heal

American Cancer Association

American Cancer Society

American Chemical Society (ACS)

American Lung Association

Buffett Cancer Center

Cancer Shop

Cancer Support

Charles Drew Health Center

CHI Health Center

Douglas County Health Department

Estabrook Cancer Center

Every Woman Matters

Great Plains Colon Cancer Task Force

Health Care Facilities

Hope Lodge

Hospitals

Jennie Edmundson Hospital

Leukemia and Lymphoma Society

Marketplace Insurance

Mayo Clinic

Methodist

Midwest GI

Mobile Diabetes Center

My Sister's Keeper

NC II Certification

Nebraska Cancer Specialists

Nebraska Coalition to End Childhood Cancer

Nebraska Medicine

One World Community Health Center

Project Pink'd

Smoking Cessation Programs

Susan G. Komen Foundation

University of Nebraska Medical Center



Wings of Hope Cancer Support Center Women's Center for Advancement

Diabetes

American Diabetes Association

American Heart Association

Americans with Disabilities Act (ADA)

Black Men United Food Pantry

Caring for Our Communities

Charles Drew Health Center

CHI Health Center

Children's Hospital and Medical Center

Clair Methodist Garden and Food Distribution

Clinical Pharmacist

Community Health Centers

Community Health Clinics

Community Health Services

Community Health Workers

Connections AEA Services

Creighton University

Dear Diabetes

Diabetes and Education Self Help Center

Diabetes Center

Diabetes Communities of Practice

Diabetes Education

Diabetes Prevention Program

Doctor's Offices

Douglas County Community Health Workers

Douglas County Help Center

Employers

Faith Based Supports

Federally Qualified Health Centers

Fitness Centers/Gyms

Food Banks/Food Pantries

HALT Diabetes Nebraska

Health Department

Health Fairs

Health Systems

Heart Ministry

Healing Gift Free Clinic (HGFC)

Home Health Agencies

Hospitals

HyVee

I Be Black Girl

Insurance Company

Juvenile Diabetes Resource Foundation

Medicare/Medicaid

Medication Assistance Programs

Methodist

Methodist College

Methodist Community Health Clinic (MCHC)

Mobile Diabetes Center

Nebraska Medicine

Nebraska Total Care

Nebraska Urban Indian Health Coalition

No More Empty Pots

North Omaha Area Health

One World Community Health Center

Pharmacv

Ponca Tribe of Nebraska Wellness Center

Pottawattamie County General Assistance

Supplemental Nutrition Assistance Program

Together Inc

U Card Benefits

University of Nebraska Medical Center

YMCA/YWCA

Disabling Conditions

Alzheimer's Association

Assisted Living

CHI Health Center

Circle Theatre Omaha

Community Alliance

Community Health Centers

Creighton University

Department of Health and Human Services

Doctor's Offices

Eastern Nebraska Office on Aging

Federally Qualified Health Centers

Health Department

Health Systems

Home Instead

Hospitals

League of Human Dignity

Meals on Wheels

Memory Care Units

Methodist

MOBY Paratransit

Nebraska Medicine

Nebraska Urban Indian Health Coalition

Nursing Homes

Ponca Tribe of Nebraska Wellness Center

Region Six

Silver Sneakers

Social Security

University of Nebraska Medical Center

VA

Heart Disease & Stroke

American Heart Association

Black Family Health and Wellness

Charles Drew Health Center



CHI Health Center

Community Health Centers

County Services

Doctor's Offices

Federally Qualified Health Centers

Fitness Centers/Gyms

Health Fairs

Health Systems

Heart Ministry

Hospitals

Managed Care Services

Methodist

Methodist Community Health Center

Nebraska Medicine

Nebraska Stroke Association

No More Empty Pots

North Omaha Area Health

Offer in One Language Only

One World Community Health Center

Reading Materials

Restaurants

Supplemental Nutrition Assistance Program

University of Nebraska Medical Center

YMCA/YWCA

Infant Health & Family Planning

Charles Drew Health Center

Children's Hospital and Medical Center

CHIP Program

Community Health Centers

Contraceptive Access Program

Department of Health and Human Services

Diaper Banks

Doctor's Offices

Douglas County Health Department

Douglas County STD Clinic

Doula Passage Program

Early Development Network

Emergency Pregnancy Services

Essential Pregnancy Services

Federally Qualified Health Centers

Girls Inc

Health Start

Help Me Grow Program

Hospitals

I Be Black Girl

Medicare/Medicaid

Methodist

Milk Works

Mothers Love

Nebraska Children's Home

Nebraska Medicine

Nonprofits

NPI-Q Questionnaire

Omaha Black Doula Association

One World Community Health Center

Planned Parenthood

Women, Infants and Children

Injury & Violence

Advocacy

Boys and Girls Club

Charles Drew Health Center

Churches

Community Health Clinics

Completely Kids

Department of Health and Human Services

Doctor's Offices

Douglas County Victim Assistance Program

EMCOMPASS Omaha

Empowerment Network

Girls Inc

Heartland Family Services

Hospitals

Law Enforcement

Nonprofits

Northstart

Omaha 360

Omaha Police Department

Project Harmony

Victim Assistance/Advocacy Program

Women's Center for Advancement

Women's Fund

YMCA/YWCA

YouTurn

Mental Health

988

AEA Services

All Care Community Health Center

Anywhere Care

Behavioral Health Connection

Behavioral Health Education Center of

Nebraska (BHCEN)

Behavioral Health Peer Supports

Best Care EAP

Boys Town

Bridge to Mental Health

Calvary Church

Capstone Behavioral Health

Catholic Charities

Center for Healing and Hope

Center for Holistic Development



Charles Drew Health Center

CHI Health Center

CHI Health Immanuel

CHI Health Mercy

CHI Heritage

CHI McDermott

Child and Family Resource Network

Community Alliance

Community Based Organizations

Community Health Centers

Community Health Clinics

Concord Mediation

Counseling Clinics

Department of Health and Human Services

Doctor's Offices

Douglas County

Douglas County Community Mental Health

Center

Douglas County Health Center

Douglas County Health Department

Douglas County Hospital

Durham Outpatient Center Clinic

Employee Assistance Program

Employers

Family Connections

Family Services

Federally Qualified Health Centers

Food Banks/Food Pantries

Fresh Hope

Full Circle Recovery

Group Homes

Harbor Pointe

Health Care Facilities

Health Department

Health Systems

Heartland Family Services

Healing Gift Free Clinic (HGFC)

Homeless Shelters

Hope Valentine and Melissa Jansen

Hospitals

Jennie Edmundson Hospital

Lasting Hope Recovery Center

Lutheran Family Services

Managed Care Services

Mental Health Court

Meridian

Methadone Clinic

Methodist Community Health Clinic (MCHC)

Mobile Crisis

Munroe Meyer

National Alliance on Mental Illness

Nebraska Association of Behavioral Health

Organizations

Nebraska Health Systems

Nebraska Medicine

Nebraska Mental Health & Aging Coalition

Nonprofits

Omaha ForUS LGBTQ Center

Omaha Police Department

One World Community Health Center

PES Emergency Room Services

Private Mental Health Therapists

Project Harmony

Psychiatric Immediate Need Clinic

Public Entities Directing/Developing Programs

Region Six

Regional Services

Remedy Health

Rural Health Clinic

Safe Harbor Peer Support

Salvation Army

School System

Southwest Iowa Mental Health and Disabilities

Service Region

Southwest Iowa Region

Spence Counseling

Stephen's Center

SWIA Region

SWIAM

Teen Mental Health Alliance

The Kim Foundation

University of Nebraska Medical Center

VA

What Makes Us Campaign

Nutrition, Physical Activity & Weight

75 North Highlander Programming

Anytime Fitness

Big Garden

Charles Drew Health Center

CHI Health Center

Children's Hospital and Medical Center

College of Lifestyle Medicine

Community Gardens

Community Health Clinics

Community Health Workers

Creighton University

Doctor's Offices

Douglas County Health Department

Farmer's Markets

Fitness Centers/Gyms

Food Bank for the Heartland

Food Banks/Food Pantries

Government Food Assistance Programs

Health Care Facilities



Health Department

Heart Ministry

Heartland B-Cycle

Heartland Family Services

Homeless Shelters

Hospitals

HyVee

Kroc Center

Libraries

Metro

Nebraska Medicine

Neighbor Good Pantry

No More Empty Pots

Nonprofits

Nutrition Services

Omaha Health Clinic

One World Community Health Center

Parks and Recreation

Planet Fitness

Podcasts

Salvation Army

School System

Senior Farmer's Market Nutrition Program

Summer Food Program

Supplemental Nutrition Assistance Program

Together Inc

University of Nebraska Medical Center

Weight Watchers

Whispering Roots

Women, Infants and Children

YMCA/YWCA

Oral Health

All Care Community Health Center

All Smiles

Charles Drew Health Center

Creighton University

Dentist's Offices

Department of Health and Human Services

Doctor's Offices

Give Kids a Smile Day

Heartland Ministry Dental Clinic

Iowa Mission of Mercy

I-Smile

Managed Care Services

Nebraska Medicine

Nonprofits

One World Community Health Center

Primer Dental Free Dental Care Day

Public Health

School System

Summit Dental

Universities

Respiratory Diseases

Charles Drew Health Center

CHI Health Center

Health Systems

Methodist Community Health Clinic (MCHC)

Nebraska Medicine

One World Community Health Center

Public Health

University of Nebraska Medical Center

Sexual Health

Adolescent Health Project

Charles Drew Health Center

CHI Health Center

Community Health Centers

Doctor's Offices

Douglas County

Douglas County Health Department

Douglas County STD Clinic

Essential Pregnancy Services

Federally Qualified Health Centers

Health Department

Hospitals

Methodist

Nebraska AIDS Project (NAP)

Nonprofits

North Omaha Area Health

Omah Public Library

Omaha ForUS LGBTQ Center

One World Community Health Center

Pharmacy

Planned Parenthood

Pottawattamie County Public Health

Public Health

School System

University of Nebraska Medical Center

Women's Fund

Social Determinants of Health

211

AEA Services

Anawim

Behavioral Health Connection

Bridge Out of Poverty

Canopy South Neighborhood Development

Caring for Our Communities

Catholic Charities



Charles Drew Health Center

CHI Health Center

CHI Health Immanuel

City Council

Community Alliance

Community-Based Organizations

Community Health Centers

Community Health Services

Community Health Workers

Community-Oriented Drug Enforcement

(CODE)

Community Relay

Connections Area Agency on Aging

Continued Utility Supply Despite Inability to

Pay

Department of Health and Human Services

Doctor's Offices

Douglas County Board of Commissioners

Douglas County Community Mental Health

Center

Douglas County Health Department

Eastern Nebraska Community Action

Partnership

Eastern Nebraska Office on Aging

Empowerment Network

Faith Based Supports

Family Housing Advisory Services

Food Banks/Food Pantries

Francis House

Front Porch Investments

Habitat for Humanity

Health Department

Heart Ministry

Heartland Family Services

Heartland Hope Food Pantry

Homeless Shelters

Hospitals

Housing and Urban Development

Housing Continuum

Housing Foundation of Sarpy County

Insurance Company

Intercultural Senior Center

Lasting Hope Recovery Center

Lied Center

Lift Up Sarpy

Lutheran Family Services

Medicare/Medicaid

Methodist Community Health Center

Metro Area Continuum of Care for the

Homeless

Micah House

MOBY

Nebraska Health Network

Nebraska Medicine

Nebraska Total Care

New Visions

No More Empty Pots

Nonprofits

Omaha 100

Omaha ForUS LGBTQ Center

Omaha Healthy Kids Alliance

Omaha Housing Authority

One World Community Health Center

Open Door Mission

Parks and Recreation

Pink For LGBTQ+

Pottawattamie County Sheriff's Office

Project Everlast

Project Harmony

Public Health

Refugee Empowerment Network

Regions

Reimagine

Salvation Army

Sarpy County Housing Authority

School System

Sienna Francis

Supplemental Nutrition Assistance Program

Southwest Iowa Mental Health and Disabilities

Service Region

Spark Community Development

Stephen's Center

Subsidized Housing Options

Together Inc

Unite Us

United Way

UnitedHealth

University of Nebraska Medical Center

Substance Use

988

AA/NA

Catholic Charities

Centerpointe

CHI Health Center

CHI Health Immanuel

CHI Health Mercy

Clinical Substance Abuse Treatment Services

Community Alliance

Community Health Centers

Council Bluffs Comprehensive Treatment

Center

Doctor's Offices

Douglas County Community Mental Health

Center

Douglas County Detox Center



Douglas County Outpatient and Inpatient

Services

Eastern Nebraska Community Action

Partnership

Employers

Family Access Center

Family Resource Center

Full Circle Recovery

Give Recovery

Haven Health

Heartland Family Services

Hospitals

InRoads to Recovery

Iowa Family Works

Jennie Edmundson Hospital

Lasting Hope Recovery Center

Life Recovery Groups

Lutheran Family Services

Nebraska Medicine

Nonprofits

Northpoint

NOVA Treatment Community

Open Door Mission

Public Health

Region Six

Salvation Army

Santa Monica House

Sarpy County Drug Court

Southwest Iowa Mental Health and Disabilities

Service Region

Stephen's Center

TDC Inpatient

Transitional Services of Iowa

University of Nebraska Medical Center

Valley Hope

Tobacco Use

Centerpointe

CHI Health Center

Doctor's Offices

Health Systems

Heartland Family Services

Lutheran Family Services

My Life My Quit

Nebraska Tobacco Quitline

Public Health

Women, Infants and Children





IMPACT OF ACTIONS TAKEN SINCE THE PRECEDING CHNA

Strategies and Program Activities by Health Need

Health Ne	ed #1: Behavioral Health
Goals and Anticipated Impact	Goals:
	Anticipated Impact: Improve continuum of care models to ensure access and utilization of mental health services Increase capacity and workforce to address acute behavioral health need Increase supportive environments that reduce tobacco use
Community Indicators	 CHNA 2016 10.3% of Omaha Metro adults reported their overall mental health as "fair" or "poor" 17% of Metro Area adults currently smoke cigarettes, either regularly or occasionally 11.1% of Douglas County adults who reports their typical day is "Extremely" or "Very" Stressful
	 CHNA 2019 8.3% of Omaha Metro adults reported their overall mental health as "fair" or "poor" 11.7% of Metro Area adults currently smoke cigarettes, either regularly or occasionally 10.0% of Metro Area adults (10.9% in Douglas County) who report their typical day is "Extremely" or "Very" Stressful 7.5% of Metro Area parents report that they have been told by a doctor or other healthcare provider that their schoolage child had depression 13.0% of Douglas County high school students report attempting suicide in the past year
	 CHNA 2022 17% believe that their overall mental health is "fair" or "poor" in Metro Area 14.2% of Metro Area adults currently smoke cigarettes, either regularly (every day) or occasionally (on some days) 25% of Metro Area adults have been diagnosed by a physician as having a depressive disorder (such as depression, major depression, dysthymia, or minor depression), worse than state and US percentages 20.2% Receiving Treatment for Mental Health in Metro Area, a statistically significant increase since 2018

			Strategic Obje			
Strategy	Key Activities	Hospital	Alignment & Integration	Clinical - Community Linkages	Capacity for Equitable Communities	Innovation & Impact
1.1 Expand access to mental health services for youth	1.1.1 Operate an integrated school-based mental health program FY23 Actions and Impact • Maintained integrated school-based mental health program in two schools located in Omaha with one dedicated provider. Began a partnership to provide therapy services for the DC West school system in Valley,Nebraska starting August 2022. In 2022 utilized both a virtual service model and in person model based on needs of the school and students. Students and families continued to struggle with the impact of COVID-19 including adjusting to returning to an in person school environment and/or virtual learning from home. Referrals from schools began to see an uptick towards the end of the academic year. FY23 Measures • # of new student referrals: 7 • # of students served (not unduplicated): 32 • Avg # of students served in school- based and virtual programs: 17 • # of billable office visits provided: 377 FY24 Actions and Impact • Maintained and integrated school-based mental health program in the DC West school system in Valley, NE. The program serves students and their families in grades K-12. FY24 Measures • Students served (not unduplicated): 41 • Average number of students served in school-based and virtual programs: 26 • New student referrals: 26 • Billable office visits: 533 FY25 Results Pending	CHI Health Immanuel				•

	 1.1.2 Pursue the establishment of the Lasting Hope Center for Children and Families FY23 Actions and Impact This strategy will no longer be pursued since the need is now being met in the community by Children's Nebraska. FY23 Measures No measures to report. FY24 Actions and Impact No updates to report. FY24 Measures No measures to report. FY25 Results Pending 	CHI Health Immanuel	•	•	•	•
1.2 Expand access to resources for individuals living with Alzheimer's and Dementia- Related Diseases (ADRD)	 1.2.1 Provide support for individuals with Alzheimer's/ dementia and their caregivers FY23 Actions and Impact Supported Alzheimer's Association through action plan collaboration. FY23 Measures # of individuals served through the 24/7 information and referral helpline: 262 # of individuals served through care consultation program: 145 # of individuals that participated in caregiver support groups: 252 # of individuals served through educational programming: 1354 FY24 Actions and Impact Not available at time of reporting. FY25 Results Pending 	All Hospitals	•	•	•	•

	_					
1.3 Expand access to behavioral health services for adults in crisis	 1.3.1 Continue to provide access to outpatient behavioral health services and reduce behavioral health readmissions through Lasting Hope Recovery Center Outpatient Clinic FY23 Actions and Impact Continued to operate an outpatient behavioral health clinic onsite at Lasting Hope Recovery Center. FY23 Measures # of completed outpatient visits: 7,513 FY24 Actions and Impact Expanded services offered at Lasting Hope, including an Intensive Outpatient Program. FY24 Measures Lasting Hope Outpatient Clinic visits: 7,644 FY25 Results Pending 	Lasting Hope Recovery Center	•	•	•	
	1.3.2 Invest in nonprofits addressing our priority health need of Behavioral Health	All Hospitals	•	•	•	•
	FY23 Actions and Impact					
	No measures to report.					
	FY24 Actions and Impact • Funded four nonprofits through Community Health Improvement Grants (CHIG): Immigrant Legal Center + Refugee Empowerment Center, Latino Center of the Midlands, Comunidad Maya Pixan Ixim, RISE Academy.					
	FY24 Measures					
	FY25 Results Pending					

1.4 Advocate for policy	1.3.3 Develop and pilot an integrated Peer Support Specialist (PSS) program across all areas of behavioral health services FY23 Actions and Impact	CHI Health Immanuel	•	•	•	•
change	behavioral health services FY23 Actions and Impact Through the Nebraska Hospital Association (NHA) and independent efforts, CHI Health supported numerous bills on the financial challenges facing our hospitals and	·				

- clinics as well as the skilled workforce we know we need to be successful.
- Victories: NE- LB227 Telehealth Reimbursement Parity and Cap on Insulin Cost Sharing; NE LB227 Extend Medicaid Postpartum Coverage. Secured an \$8 million appropriation for the CHI Health Central Kitchen project in Omaha, NE. Supported a \$335 million appropriation for social determinants of health projects in North and South Omaha, NE.

FY23 Measures

 # of behavioral health bills approved (of those supported by NHA and/or CHI Health): 2

FY24 Actions and Impact

- Through the Nebraska Hospital Association, CHI Health supported advocacy efforts that culminated in the passage of several laws that preserve access and sustainability of behavioral health services, such as:
 - LBB896- Change provisions relating to telehealth consultations Included in Health and Human Services Committee Priority Legislation eliminates the requirement to collect a signed consent statement for a telehealth consultation if verbal consent has already been given. The bill removes the following language: "The signed statement may be collected by paper or electronic signature and shall become a part of the patient's medical record. If the patient gives verbal consent during the initial telehealth consultation, the signed statement shall be collected within ten days after such telehealth consultation."
 - LB62 Coverage of translation and interpretation services under Medicaid requires the state to provide coverage for all necessary translation and interpretation services for individuals who receive Medicaid. It requires the Department of Health and Human Services to provide the coverage, as well as reimbursement for providers, and to "take all actions necessary" to maximize federal funding to do so.

	L DOET Creates the Nobreade Drenatel Dive		<u> </u>	
0	LB857 Creates the Nebraska Prenatal Plus			
	Program, provide for use of the Medicaid			
	Managed Care Excess Profit Fund, and change			
	provisions relating to coverage of glucose			
	monitors under the Medical Assistance Act.			
	LB857 creates the Nebraska Prenatal Plus			
	Program at DHHS for women eligible for			
	Medicaid or the Children's Health Insurance			
	Program to reduce the incidence of low birth			
	weight, pre-term birth, and adverse birth			
	outcomes while also addressing other lifestyle,			
	behavioral, and nonmedical aspects of an at-			
	risk mother's life that may affect the health and			
	well-being of the mother or the child. Services			
	eligible for reimbursement for at-risk mothers			
	include nutrition counseling, psychosocial			
	counseling and support, general client			
	education and health promotion, breastfeeding			
	support, and targeted case management.			
	LB857 limits eligibility for at-risk mothers			
	otherwise eligible for medicaid while pregnant,			
	removes CHIP eligibility and postpartum			
	coverage. It also limits nutrition counseling to 6			
	or fewer sessions.			
0	LB905 Require the Department of Health			
	and Human Services to submit a Medicaid			
	waiver or state plan amendment for medical			
	respite care, change the definition of respite			
	care, and change provisions relating to the			
	Medicaid Managed Care Excess Profit Fund.			
	LB905 requires DHHS to submit a Medicaid			
	waiver or state plan amendment to designate			
	two medical respite facilities to reimburse for			
	services provided to an individual who is			
	homeless and an adult in the expansion			
	population no later than Jan. 1, 2025.			
0	LB1355 Provide for release of certain			
	patient data by an emergency medical			
	service, change provisions of the Opioid			
	Prevention and Treatment Act, and provide for			
	state aid, research, first responder training, and			
	overdose fatality review. LB1355 requires the			
	Department of Health and Human Services to			
	provide grants to local public health			

departments, law enforcement agencies, and health care facilities, to aid programs to facilitate opioid overdose prevention efforts, education and training, recovery, problemsolving courts, and data tracking. The bill includes the intent of the Legislature to appropriate \$4,000,000 annually from the Nebraska Opioid Recovery Fund beginning in FY2024-25 for grants under the Opioid Prevention and Treatment Act. Would provide \$500,000 each to a local public health department, a law enforcement agency applicant, and to a health care facility applicant. LB1355 allows pharmacists to sell, and public health departments to distribute, fentanyl test strips. FY24 Measures • Behavioral health bills approved (of those supported by NHA and/or CHI Health): 5				
FY25 Results Pending				
 1.4.2 Support Sarpy/Cass tobacco coalition FY23 Actions and Impact Tobacco Education & Advocacy of the Midlands (TEAM) continued to work with businesses, school districts, public housing programs and city parks to implement smoke-free/ vape-free/ tobacco- free policies. Launched initiatives to decrease the number of tobacco retailers and increase the number of cessation assessment and referral system policies within health care, social service, and education agencies. FY23 Measures # of NEW coalition members: 26 # of multi-family buildings that adopted 100% smokefree/ vape- free/ tobacco- free policies within Sarpy/ Cass County: 71 	CHI Health Midlands	•	•	
FY24 Actions and Impact • Tobacco Education & Advocacy of the Midlands (TEAM) continued to work with businesses, school districts, public housing programs and city parks to implement				

	smoke-free/ vape-free/ tobacco- free policies. Continued working on new initiatives to decrease the number of tobacco retailers and increase the number of cessation assessment and referral system policies within health care, social service, and education agencies. FY24 Measures TEAM Coalition Members: 116 New Members: 9 Multi-family buildings that adopted 100% smoke-free/vape-free/tobacco-free policies within Sarpy/Cass County: 142 Jurisdictions that designate outdoor recreational areas as smoke-free/vape-free or tobacco-free in Sarpy/Cass County: 10 Cessation assessment and referral system policies within health care, social service, and education agencies in Sarpy/Cass County: 8					
	FY25 Results Pending					
1.5 Decrease behavioral health workforce shortage through educational partnerships	 1.5.1 Develop, implement and evaluate behavioral health workforce recruitment and retention strategies FY23 Actions and Impact Implemented recruitment strategies further reporting will be incorporated in FY24. FY23 Measures No measures to report. FY24 Actions and Impact Continued to implement retention strategies. Utilized funding for quarterly recognition of preceptors, supervisors, and mentors and for sponsoring conferences and continuing education. Relocated the Creighton Psychiatric Residency Program to Immanuel to accommodate additional therapists at Lasting Hope. Added a new Intensive Outpatient Program for Chemical Dependency at Lasting Hope FY24 Measures 	All Hospitals	•	•	•	•

Psychiatrists: 4 Psychiatrists: 4 Psychiatric Nurse Practitioners: 6 Chemical Dependency Program		-		1			
1.6.1 Support internal/community mental health stigma reduction campaigns FY23 Actions and Impact Engaged in Regional Health Council Community Roundtable and Regional health Council Health Systems Roundtable. Phase 2 MH Stigma Reduction campaign was implemented (Dec 2022- Jan 2023). What Makes Us is a campaign that is designed to reduce stigma by encouraging individuals to share their lived experiences with mental challenges and triumphs. FY23 Measures My workplace takes an active role in my mental health. (not campaign aware = 59.1% vs. campaign aware = 72.6%) Therapy and counseling can be an effective treatment for people with mental health conditions. (not campaign aware = 80% vs. campaign aware = 94.4%) Are you comfortable offering support to other people about their mental health conditions? (not campaign aware = 64.6% vs. campaign aware = 91.1% I would be willing to work with someone with a mental health condition. (baseline = 70.3%, end of year 2 = 70.9%) Y2 Campaign (May 2022- April 2023) # of impressions: 2,793,733 impressions # of social engagements (likes/ comments/		 Psychiatrists: 4 Psychiatric Nurse Practitioners: 6 Chemical Dependency Program Therapists: 2 Nurse: 1 Addiction Psychiatrist: 1 Vacancy Rates as of July 2024 Immanuel: 18.98% (23.67% in June 2024) Lasting Hope: 30.34% (39.33% in June 2024) 					
reduction campaigns FY23 Actions and Impact Engaged in Regional Health Council Community Roundtable and Regional health Council Health Systems Roundtable. Phase 2 MH Stigma Reduction campaign was implemented (Dec 2022- Jan 2023).What Makes Us is a campaign that is designed to reduce stigma by encouraging individuals to share their lived experiences with mental challenges and triumphs. FY23 Measures My workplace takes an active role in my mental health. (not campaign aware = 59.1% vs. campaign aware = 72.6%) Therapy and counseling can be an effective treatment for people with mental health conditions. (not campaign aware = 80% vs. campaign aware = 94.4%) Are you comfortable offering support to other people about their mental health conditions? (not campaign aware = 64.6% vs. campaign aware = 91.1% I would be willing to work with someone with a mental health condition. (baseline = 70.3%, end of year 2 =70.9%) Y2 Campaign (May 2022- April 2023) # of impressions: 2,793,733 impressions # of social engagements (likes/ comments/	1.6 Reduce mental		All Hospitals	•	•	•	•
FY24 Actions and Impact		reduction campaigns FY23 Actions and Impact Engaged in Regional Health Council Community Roundtable and Regional health Council Health Systems Roundtable. Phase 2 MH Stigma Reduction campaign was implemented (Dec 2022- Jan 2023). What Makes Us is a campaign that is designed to reduce stigma by encouraging individuals to share their lived experiences with mental challenges and triumphs. FY23 Measures My workplace takes an active role in my mental health. (not campaign aware = 59.1% vs. campaign aware = 72.6%) Therapy and counseling can be an effective treatment for people with mental health conditions. (not campaign aware = 80% vs. campaign aware = 94.4%) Are you comfortable offering support to other people about their mental health conditions? (not campaign aware = 64.6% vs. campaign aware = 91.1% I would be willing to work with someone with a mental health condition. (baseline = 70.3%, end of year 2 =70.9%) Y2 Campaign (May 2022- April 2023) # of impressions: 2,793,733 impressions # of social engagements (likes/ comments/ shares on social media): 35,672	7 III Tioophais				

	 Implemented the American Hospital Association's People Matter Words Matter Campaign by distributing monthly posters and communications and ensuring that all organization leaders knew how to order promotional materials. FY24 Measures No measures to report. FY25 Results Pending 					
1.7 Early detection of depression and connection to mental health services among pregnant people	1.7.1 Conduct perinatal depression screening during prenatal visit, while inpatient for delivery and at postpartum visit FY23 Actions and Impact	CHI Health CUMC- Bergan Mercy, CHI Health Immanuel, and CHI Health Lakeside	•	•	•	•
1.8 Prevent suicide	 1.8.1 Explore local school needs regarding suicide prevention suicide and provide training and financial resources (e.g. Question. Persuade. Respond (QPR), Mental Health First Aid, etc.) FY23 Actions and Impact No actions to report. FY23 Measures No measures to report. 	All Hospitals	•	•	•	•

	FY24 Actions and Impact No actions to report. FY24 Measures No measures to report.					
	FY25 Results Pending					
Related Activities	The following activities represent complementary efforts in which identified health need through financial support, in-kind staff control CHI Health offers integrated behavioral health services expand access to behavioral health services in a familia CHI Health Primary Care Clinics use the Screening, Brid depression, drug and alcohol abuse screening and assess SBIRT is administered annually during a wellness exam Additionally, CHI Health addresses the need for behavioral health Operation of Lasting Hope Recovery Center, a 64-bed poperation of a Pediatric Residential Treatment Facility (Participation in the Sarpy County Youth Mental Health Facility (Participation in the Sarpy County Youth Mental Health Facility (Participation in the Sarpy County Youth Mental Health Facility (Participation in the Sarpy County Youth Mental Health Facility (Participation in the Sarpy County Youth Mental Health Facility (Participation in the Sarpy County Youth Mental Health Facility (Participation in the Sarpy County Youth Mental Health Facility (Participation in the Sarpy County Youth Mental Health Facility (Participation in the Sarpy County Youth Mental Health Facility (Participation in the Sarpy County Youth Mental Health Facility (Participation in the Sarpy County Youth Mental Health Facility (Participation in the Sarpy County Youth Mental Health Facility (Participation in the Sarpy County Youth Mental Health Facility (Participation in the Sarpy County Youth Mental Health Facility (Participation in the Sarpy County Youth Mental Health Facility (Participation in the Sarpy County Youth Mental Health Facility (Participation in the Sarpy County Youth Mental Health Facility (Participation in the Sarpy County Youth Mental Health Facility (Participation in the Sarpy County Youth Mental Health Participation in the Sarpy Coun	ribution or a combin in CHI Health Prima r setting. If the setting in the ribustal in the Omerschiatric treatment on the Omerschiatric treatment of the Ome	ation there ary Care Cl Referral to ed for patie naha Metro t facility ne CHI Imn	of. inics in ord Treatment ents 12 yea through th	er to conveni (SBIRT), a urs of age and e following:	ently universal
Planned Resources	The hospital will provide staff time, philanthropic cash grants, out these initiatives.	reach communication	ons, and pr	ogram mar	nagement su	pport for
Planned Collaborators	 Omaha Metro Schools (Omaha Public Schools, including Keand Douglas County (DC) West Community Schools Heritage Services Alzheimer's Association Nebraska Hospital Association Tobacco Education & Advocacy of the Midlands (T.E.A.M.) The Wellbeing Partners Nebraska Perinatal Quality Improvement Collaborative CHI Health Behavioral Health Service Line CHI Health Forensic Nurse Examiner Program Kim Foundation 	ellom Elementary, C	UES Scho	ol System,	including Ho	ly Name